

Summary of Lec.2

Math 212a

Sept. 21, 2000

- **Completion.** Any metric space X can be completed to yield a complete metric space. More precisely, there is a metric space $X_{complete}$, unique up to isometry, which is complete, and an isometry $\phi : X \rightarrow X_{complete}$ such that $\phi(X)$ is dense in $X_{complete}$. See *completion* in the folder for Lecture 1 for details. If X is a normed vector space, then $X_{complete}$ inherits the structure of a normed vector space. A pre-Hilbert space is characterized among all normed spaces by the fact that it satisfies the parallelogram law (see below). Hence, if X is a pre-Hilbert space, then $X_{complete}$ is a Hilbert space.

- **The Cauchy-Schwartz inequality.** This says that

$$|(f, g)| \leq (f, f)^{\frac{1}{2}}(g, g)^{\frac{1}{2}}$$

for any elements f, g in a space with a positive semi-definite scalar product. See *hilbertspace* in the folder for Lecture 1 for details.

- **The Pythagorean theorem.** This says that in a pre-Hilbert space

$$\|f + g\|^2 = \|f\|^2 + \|g\|^2 \quad \text{if } (f, g) = 0.$$

See *hilbertspace* in the folder for Lecture 1 for details.

- **The parallelogram law.** This says that in a pre-Hilbert space

$$\|f + g\|^2 + \|f - g\|^2 = 2(\|f\|^2 + \|g\|^2).$$

See *hilbertspace* in the folder for Lecture 1 for details.

- **Orthogonal projection.** If M is a complete subspace of a pre-Hilbert space H , then for any $v \in H$ there is a unique $w \in M$ such that $(v - w) \perp M$. This w is characterized as being the unique element of M which minimizes $\|v - x\|$, $x \in M$. The idea of the proof is to use the parallelogram law to conclude that if $\{x_n\}$ is a sequence of elements in M for which $\|v - x_n\|$ approaches the greatest lower bound of $\|v - x\|$, $x \in M$, then $\{x_n\}$ is a Cauchy sequence. Then the assumption that M is complete guarantees that this sequence has a limit $w \in M$ which minimizes $\|v - x\|$, $x \in M$. See *hilbertspace* in the folder for Lecture 1 for details. The unique $w \in M$ so obtained is denoted by $\pi_M(v)$.

- **The Riesz representation theorem.** This says that any bounded linear function on a Hilbert space H is given by scalar product with an element of H . That is, every bounded linear function ℓ is given by

$$f \mapsto (f, g)$$

where g is a (unique) element of H . If the linear function is identically zero, we may take $g = 0$. Otherwise, the codimension of the kernel of ℓ , the space of all f such that $\ell(f) = 0$, has codimension one. Let N denote this kernel which is a closed (hence complete) subspace. Via orthogonal projection, there is a $y \in H$ with $y \perp N$ and $\|y\| = 1$. So $(f - (f, y)y) \perp y$, and the set of all elements perpendicular to y is a subspace of codimension 1 which contains N , and so must coincide with N . Thus $(f - (f, y)y) \in N$. The decomposition

$$f = (f, y)y + (f - (f, y)y)$$

shows that $\ell(f) = (f, y)\ell(y)$ so

$$\ell(f) = (f, g) \quad \text{where } g = \overline{\ell(y)}y.$$

- **Description of $L_2(\mathbf{T})$ as a space of linear functions.** $L_2(\mathbf{T})$ is defined as the completion of the pre-Hilbert space $\mathcal{C}(\mathbf{T})$, the space of continuous functions on the circle \mathbf{T} under the scalar product

$$(f, g) = \frac{1}{2\pi} \int_{\mathbf{T}} f(t)\overline{g(t)}dt.$$

Any bounded linear function on this pre-Hilbert space extends uniquely to a bounded linear function on its completion, $L_2(\mathbf{T})$. Hence by the Riesz representation theorem is given by scalar product with an element of $L_2(\mathbf{T})$. In other words, we should regard an element of $L_2(\mathbf{T})$ not as a function on the circle, but as a linear function on the space of continuous functions on the circle which is continuous relative to the norm $\|\cdot\|_2$.