

Problem set 6

Math 212a

November 4, 2004, Due Nov 11

In this problem set we present some standard counter-examples in the theory of integration and then return to the study of generalized functions.

Contents

1 Three standard counterexamples.	1
1.1 Non-dominated convergence.	1
1.2 Order of integration matters for non-integrable functions.	2
1.3 Order of integration matters in the non- σ -finite case.	2
2 The spaces \mathcal{D} and \mathcal{D}'.	2
3 Cauchy principal value.	4
4 Anti-derivatives of elements of \mathcal{D}'.	4
5 The tensor product of two generalized functions.	5

1 Three standard counterexamples.

1.1 Non-dominated convergence.

Let $f_n := n\mathbf{1}_{(0, \frac{1}{n}]}$. This is similar to one of the examples we studied in class where $\int_{\mathbf{R}} f_n dx \equiv 1$ while $\lim f_n(x) = 0$ for all x . So we get strict inequality in Fatou's lemma, and can not move the limit past the integral sign. So we know in advance that this can not be a case of dominated convergence.

1. Let g be defined by $g(x) := \sup_k f_k(x)$. We know that $g \notin \mathcal{L}_1$. But compute g and show explicitly that its integral diverges.

1.2 Order of integration matters for non-integrable functions.

2. Take $X = Y = [0, 1]$ with standard Lebesgue measure. Let

$$f(x, y) = \begin{cases} \frac{1}{x^2} & \text{if } 0 < y < x < 1 \\ -\frac{1}{y^2} & \text{if } 0 < x < y < 1 \\ 0 & \text{otherwise} \end{cases} .$$

Compute $\int_X (\int_Y f(x, y) dy) dx$ and $\int_Y (\int_X f(x, y) dx) dy$ and verify that they are not equal. This implies that $f \notin \mathcal{L}_1(X \times Y, \mathbf{R})$ for otherwise Fubini would apply. Show directly that both f^+ and f^- have infinite integrals on $X \times Y$.

1.3 Order of integration matters in the non- σ -finite case.

Let $X = Y = [0, 1]$ as before. Take \mathcal{F} to be the usual Lebesgue measurable sets and m to be Lebesgue measure. Take \mathcal{G} to be “all” subsets, and take n to be

$$n(A) = \text{number of elements in } A$$

(in particular $n(A) = \infty$ if A is not a finite set). Since $[0, 1]$ is not the countable union of finite sets, it is not σ -finite. So we expect that Fubini might fail. Let $\Delta \subset X \times Y$ be the diagonal, i.e.

$$\Delta := \{(x, y) | x = y\}.$$

3. Show that $\Delta \in \mathcal{F} \times \mathcal{G}$. Let

$$f := \mathbf{1}_\Delta.$$

Evaluate $\int_X (\int_Y f(x, y) dy) dx$ and $\int_Y (\int_X f(x, y) dx) dy$ and verify that they are not equal.

2 The spaces \mathcal{D} and \mathcal{D}' .

In the previous problem set we defined a generalized function to be a continuous linear function on the Schwartz space \mathcal{S} , and denoted the space of generalized functions by \mathcal{S}' . For various reasons (especially when we want to extend the theory to manifolds) it is convenient to study other spaces of “test functions” and “generalized functions”.

Let \mathcal{D} denote the space of infinitely differentiable functions each of which vanishes outside a compact set K (which may depend on the function). A sequence $f_n \in \mathcal{D}$ is said to converge to an $f \in \mathcal{D}$ if there is a fixed compact set K such that $\text{supp } f_n \subset K$ for all n , and then such that the f_n converge to

f uniformly, together with uniform convergence of all the derivatives. Then \mathcal{D}' consists of all linear functionals which are continuous relative to this notion of convergence.

We should really write $\mathcal{D}(\mathbf{R})$ if we are thinking of functions of one real variable, but we could equally well consider functions of n variables, in which case we would write $\mathcal{D}(\mathbf{R}^n)$ or write $\mathcal{D}(V)$ where V is any finite dimensional vector space.

For example, if Δ denotes the Laplacian on \mathbf{R}^3 (or \mathbf{R}^n), then one way of writing Green's theorem is

$$\int_G f \Delta \phi dx = \int_G (\Delta f) \phi dx + \int_{\partial G} \left(f \frac{\partial \phi}{\partial n} - \frac{\partial f}{\partial n} \phi \right) dS \quad (1)$$

where f and ϕ are smooth functions, G is a region with smooth (or piecewise smooth) boundary ∂G , where dS denotes the surface measure on the boundary and $\partial/\partial n$ denotes normal derivative. If we take both f and ϕ to be in \mathcal{D} , and take G so large that both $\text{supp } f$ and $\text{supp } \phi$ lie in the interior of G , so that there are no boundary terms, then (1) becomes

$$\langle \Delta f, \phi \rangle = \langle f, \Delta \phi \rangle. \quad (2)$$

For a generalized function f we take this as the *definition* of Δf . In other words, Δf is *defined* to be that generalized function given by

$$\langle \Delta f, \phi \rangle := \langle f, \Delta \phi \rangle \quad \forall \phi \in \mathcal{D}. \quad (3)$$

Let us illustrate this definition when f is *not* an element of \mathcal{D} : Let g be a smooth function, and let G be a domain with smooth boundary as in (1). Let

$$f = \mathbf{1}_G g.$$

As a *function*, f is equal to g inside G and to be zero outside G , and so the definition of f as a generalized function is

$$\langle f, \psi \rangle := \int_G g \psi dx, \quad \forall \psi \in \mathcal{D}.$$

Then this definition and (1) say that Δf is the sum of the function $(\Delta g)\mathbf{1}_G$ plus two other terms supported on the boundary:

$$\langle \Delta f, \phi \rangle = \int_G (\Delta g) \phi dx + \int_{\partial G} g \frac{\partial \phi}{\partial n} dS - \int_{\partial G} \frac{\partial g}{\partial n} \phi dS. \quad (4)$$

In classical electrostatics these terms supported on the boundary are thought of as a distribution of electric charges and dipole moments on the boundary. For this reason Schwartz used the term "distribution" to denote an element of \mathcal{D}' .

In three dimensions, consider the function $1/r$ (where r is the distance from the origin). Since the function $1/r$ is integrable in three dimensions at the origin,

it defines an element of \mathcal{S}' and of \mathcal{D}' and so we can use (3) to define $\Delta(1/r)$. The *function* $1/r$ is smooth away from the origin, and (by direct computation) satisfies $\Delta(1/r) = 0$ there as a *function*. But $1/r$ is *not* a smooth function on all of \mathbf{R}^3 . So we must use (3).

4. Show that

$$\Delta\left(\frac{1}{r}\right) = -4\pi\delta.$$

[Hint: Apply (1) to the region consisting of $\epsilon \leq r \leq R$ where R is chosen so large that $\text{supp } \phi$ is contained in the open ball of radius R so that there are no terms coming from the outer boundary $r = R$. Compute the terms coming from the inner boundary and let $\epsilon \rightarrow 0$.]

3 Cauchy principal value.

Back to one dimension temporarily. The function $x \mapsto 1/x$ is not locally integrable in one dimension, but we do have $\log(|x|)' = 1/x$ at all $x \neq 0$ and $\log|x|$ is locally integrable and so defines a generalized function. We may therefore try to define a generalized function by taking the derivative of $\log|x|$ in the sense of generalized functions.

5. Show that $(\log|x|)' = \text{pv}\left(\frac{1}{x}\right)$ where

$$\langle \text{pv}\left(\frac{1}{x}\right), \phi \rangle = \int_0^\infty \frac{\phi(x) - \phi(-x)}{x} dx.$$

The symbol “pv” stands for principal value, a notion that was introduced by Cauchy. Also compute the second derivative of $\log|x|$.

4 Anti-derivatives of elements of \mathcal{D}' .

We want to study the differential equation

$$g' = f. \tag{5}$$

Here f is an element of \mathcal{D}' and we look for a $g \in \mathcal{D}'$ which satisfies this equation (and differentiation is in the sense of generalized functions). We first examine the simplest case where $f = 0$:

6. Show that if $U \in \mathcal{D}'$ has derivative zero, then U is a constant, i.e. show that $\langle U, \phi \rangle = C \int_{\mathbf{R}} \phi dx$. [Hint: let \mathcal{D}_0 denote the set of all $\psi \in \mathcal{D}$ such that $\int_{\mathbf{R}} \psi dx = 0$. First show that U vanishes on \mathcal{D}_0 . Then choose an element $\theta \in \mathcal{D}$ with $\int_{\mathbf{R}} \theta dx = 1$. Write every $\phi \in \mathcal{D}$ as $\phi = \psi + a\theta$ where $a = \int_{\mathbf{R}} \phi dx$.]

We can use a similar idea to show that a solution of (5) exists for a general $f \in \mathcal{D}'$. Indeed, (5) says that

$$-\langle g, \phi' \rangle = \langle f, \phi \rangle \quad \forall \phi \in \mathcal{D}. \quad (6)$$

This defines g on the set of all $\psi \in \mathcal{D}$ which are derivatives (in the usual sense) of elements of \mathcal{D} . But these are the elements of \mathcal{D}_0 . So we wish to extend g (which is defined by (6) on \mathcal{D}_0) so as to be defined on all of \mathcal{D} . Choose θ as in Problem 6 and write a general $\phi \in \mathcal{D}$ as

$$\phi = a\theta + \phi_0, \quad a = \int_{\mathbf{R}} \phi dx.$$

The map

$$\phi \mapsto \phi_0$$

is a continuous linear map, call it π , from \mathcal{D} to \mathcal{D}_0 which is the identity map when restricted to \mathcal{D}_0 . So we can think of it as a “projection” from \mathcal{D} to \mathcal{D}_0 . (Of course, it depends on the choice of θ so we should (but won't) write π_θ .) Then define

$$\langle g, \phi \rangle := \langle g, \pi(\phi) \rangle = \langle g, \phi_0 \rangle.$$

Now g is defined on all of \mathcal{D} and, by construction, (6) is satisfied. Problem 6 now implies that the solution g of (5) is unique up to “adding a constant”.

7. Use the preceding discussion to show the following: Suppose that g is a continuous function whose derivative in the sense of generalized functions is a continuous function. I.e. assume that $-\int_{\mathbf{R}} g\phi' dx =: \langle g', \phi \rangle = \int_{\mathbf{R}} f\phi dx =: \langle f, \phi \rangle$ where f is a continuous function. Show that g is in fact continuously differentiable as a function and that $g' = f$ in the usual sense.

5 The tensor product of two generalized functions.

Let X and Y be two finite dimensional vector spaces over the real numbers, so that $X \times Y = X \oplus Y$ is again a finite dimensional vector space over the real number. We will write the typical point of $X \times Y$ as (x, y) . We can consider the space $\mathcal{D}(X \times Y)$ which consists of all infinitely differentiable functions of compact support on $X \times Y$. We can also consider the space $\mathcal{D}(X) \otimes \mathcal{D}(Y)$ which consists of all finite linear combinations of expressions of the form $\phi\psi$ where $\phi \in \mathcal{D}(X)$ and $\psi \in \mathcal{D}(Y)$. Any such expression defines a function on $X \times Y$ by the rule

$$(\phi\psi)(x, y) = \phi(x)\psi(y),$$

so we have an injection

$$\mathcal{D}(X) \otimes \mathcal{D}(Y) \mapsto \mathcal{D}(X \times Y)$$

which allows us to think of $\mathcal{D}(X) \otimes \mathcal{D}(Y)$ as a subspace of $\mathcal{D}(X \times Y)$. The Stone-Weierstrass theorem, or the original Weierstrass approximation theorem guarantees that $\mathcal{D}(X) \otimes \mathcal{D}(Y)$ is dense in $\mathcal{D}(X \times Y)$.

Now let $f \in \mathcal{D}'(X)$ and $g \in \mathcal{D}'(Y)$ be generalized functions. Let $\phi \in \mathcal{D}(X \times Y)$. For each fixed x , the function $\phi(x, \cdot) : y \mapsto \phi(x, y)$ belongs to $\mathcal{D}(Y)$ and so we can apply g to it to obtain the function $x \mapsto \langle g, \phi(x, \cdot) \rangle$. This is a function of x , and the continuity properties of g imply that this function belongs to $\mathcal{D}(X)$. We then may apply f to the function $x \mapsto \langle g, \phi(x, \cdot) \rangle$ to obtain a number. The notation is getting cumbersome, so we will shorten it and write the final result as

$$\langle f, \langle g, \phi \rangle \rangle.$$

We then define $f \otimes g$ to be this generalized function. In other words we define $f \otimes g \in \mathcal{D}'(X \times Y)$ by

$$\langle f \otimes g, \phi \rangle = \langle f, \langle g, \phi \rangle \rangle. \quad (7)$$

If $\phi = \tau \otimes \eta$ where $\tau \in \mathcal{D}(X)$ and $\eta \in \mathcal{D}(Y)$ then it is clear from the definition that

$$\langle f \otimes g, \tau \eta \rangle = \langle f, \tau \rangle \langle g, \eta \rangle.$$

This shows that on function of the form $\tau \eta$ it would not have made any difference in the definition of $f \otimes g$ had we done things in the reverse order, i.e. first apply f to the function $x \mapsto \phi(x, y)$ and then apply g to the resulting function of y . But since $\mathcal{D}(X) \otimes \mathcal{D}(Y)$ is dense in $\mathcal{D}(X \times Y)$ it follows that doing things in the reverse order yields the same answer on all of $\mathcal{D}(X \times Y)$. This is a sort of “generalized function version” of Fubini’s theorem.

Similarly, if we have three vector spaces X, Y and Z and $f \in \mathcal{D}(X)$, $g \in \mathcal{D}(Y)$, $h \in \mathcal{D}(Z)$ we can form $f \otimes (g \otimes h)$ and $(f \otimes g) \otimes h$ and verify that they give the same element of $\mathcal{D}(X \times Y \times Z)$.

It is easy to check directly from the definition that

$$\text{supp}(f \otimes g) = \text{supp}(f) \times \text{supp}(g)$$

as a subset of $X \times Y$.

Suppose that $X = Y = Z$, and to fix the ideas (and hopefully to get the powers of 2π right, although I am not all that optimistic) that they all equal \mathbf{R} . If f and g were elements of L_1 we defined their convolution $f \star g$ as

$$(f \star g)(u) = \frac{1}{\sqrt{2\pi}} \int_{\mathbf{R}} f(x)g(u-x)ds.$$

If we think of this as a generalized function and apply it to ϕ we get

$$\frac{1}{\sqrt{2\pi}} \int_{\mathbf{R}} \int_{\mathbf{R}} f(x)g(u-x)dx\phi(u)du = \frac{1}{\sqrt{2\pi}} \int_{\mathbf{R}} \int_{\mathbf{R}} f(x)g(y)\phi(x+y)dx dy.$$

Therefore we would like to define the convolution $f \star g$ of any two elements of $\mathcal{D}'(\mathbf{R})$ by

$$\langle f \star g, \phi \rangle = \frac{1}{\sqrt{2\pi}} \langle f \otimes g, \phi(x+y) \rangle \quad (8)$$

Here $\phi \in \mathcal{D}(\mathbf{R})$ and $\phi(x+y)$ denotes the function of two variables given by $(x, y) \mapsto \phi(x+y)$. The trouble is that $\phi(x+y)$ does not have compact support as a function of two variables. Indeed it is constant on any “anti-diagonal” line $x+y = a$. So the $\text{supp } \phi(x+y)$ is the anti-diagonal strip consisting of all (x, y) such that $x+y \in \text{supp}(\phi)$. So we can not use (8) in general. But by the remarks of the last section of the preceding exercise set, we can use this definition if we know that $\text{supp}(f) \times \text{supp}(g)$ intersects every anti-diagonal strip (of bounded width) in a compact set.

This will happen, for example, if

- Either f or g has compact support. For then $\text{supp}(f \otimes g)$ will be a horizontal or a vertical strip, which then meets any anti-diagonal strip of bounded width in a compact set. Or
- $\text{supp}(f) \subset [a, \infty)$ and $\text{supp}(g) \subset [b, \infty)$ for then $\text{supp}(f \otimes g)$ is contained in the (infinite) rectangle $[a, \infty) \times [b, \infty)$ which also intersects any bounded anti-diagonal strip in a compact set.

As an illustration of the first case, consider what happens if we take $f = \delta$. Then

$$\langle \delta \star g, \phi \rangle = \frac{1}{\sqrt{2\pi}} \langle \delta \otimes g, \phi(x+y) \rangle = \frac{1}{\sqrt{2\pi}} \langle g, \langle \delta, \phi(x+y) \rangle \rangle = \frac{1}{\sqrt{2\pi}} \langle g, \phi \rangle.$$

In other words,

$$(\sqrt{2\pi}\delta) \star g = g$$

for any $g \in \mathcal{D}'(\mathbf{R})$. Convolution with the element $\sqrt{2\pi}\delta$ is the identity operator.

Similarly, we can define the convolution of two generalized functions in n dimensions, the unfortunate factor $\frac{1}{\sqrt{2\pi}}$ (which was determined by our conventions for the Fourier transform) being replaced by the factor $\frac{1}{(2\pi)^{n/2}}$. The second sufficient condition for the definition of the convolution is replaced by the condition that $\text{supp}(f)$ and $\text{supp}(g)$ are both contained in the same type of “orthant”, for example in the “first quadrant” (or any translate thereof) in two variables.

Let D be any differential operator with constant coefficients. Define D^* to be the formal adjoint of D obtained by integration by parts and ignoring the boundary terms. So, for example, if D is a homogeneous differential operator of order k , then $D^* = (-1)^k D$. For instance $\Delta^* = \Delta$ for the Laplacian. The operator D is then defined on generalized functions by

$$\langle Df, \phi \rangle := \langle f, D^* \phi \rangle.$$

8. Show that

$$D(f \star g) = (Df) \star g = f \star Dg$$

whenever $f \star g$ is defined. In particular, in three dimensions, if f is any generalized function then

$$u = \left(\frac{c}{r}\right) \star f$$

is a solution to

$$\Delta u = f$$

where (I hope) $c = -\sqrt{\frac{\pi}{2}}$.