

## Answers to Second Math 21a Practice Hourly 1

Note: Problems 1-6 count 8 points each, while Problem 7 counts for 2 points.

1. Answer:

- a)  $|\mathbf{v}| = \sqrt{21}$ ,  $|\mathbf{w}| = 1$ ,  $|\mathbf{v} - 3\mathbf{w}| = \sqrt{6}$ .
- b) This number is  $\mathbf{v} \cdot \mathbf{w} = 4$ .
- c)  $(-2, 0, 1)$ .
- d)  $b = -2$  and  $c = 0$ .

2. Answer:

- a)  $(3/2, 0, 0)$ ,  $(0, -3/2, 0)$  and  $(0, 0, 3)$ . (There are infinitely many other possibilities)
- b)  $(2, -2, 1)$ , or any non-zero multiple
- c) The distance is 1.
- d) Parametric: Send  $t \rightarrow (3t, 3t, 3)$ . Nonparametric:  $x = y$  &  $z = 3 - 2x + 2y$ .  
(There are infinitely many other possibilities.)

3. Answer:

- a)  $(0, -3, 4\pi)$ .
- b)  $2\sqrt{\pi}(-3, 0, 4)$
- c)  $t \rightarrow (-3t, -3, 4(\pi + t))$ .
- d) The velocity vector at general  $t$  is  $2t(3\cos(t^2), -3\sin(t^2), 4)$  whose length is  $10t$ .  
Thus, the distance is the integral of this last function from 0 to  $\sqrt{\pi}$  which is  $5\sqrt{\pi}$ .

4. Answer:

- a)  $\mathbf{p}$  is on  $L$  and  $\mathbf{v}$  is tangent to  $L$ .
- b)  $t \rightarrow \mathbf{p} + t\mathbf{v}$ .
- c)  $(-4, -3, 0)$ ; the case of  $t = -1$  in the preceding parameterization.
- d)  $d = |\mathbf{p} \times \mathbf{v}|/|\mathbf{v}| = \sqrt{\frac{34}{35}}$ .

5. Answer:

- a) The plane where  $x - 3z = 0$ . (There are infinitely many other possibilities.)
- b)  $\mathbf{w} \cdot \mathbf{v} = 0$  requires  $b = 16$ .
- c)  $b = -6$  and  $c = -2$ .
- d) If  $\mathbf{s}$  is parallel to  $\mathbf{v}$ , then  $\mathbf{u} \times \mathbf{s}$  will be perpendicular to  $\mathbf{v}$ . For this, take  $c = -6$ .

6. Answer:

- a) True. Indeed, since  $\mathbf{r} \cdot \mathbf{r} = |\mathbf{r}|^2$  and  $\frac{d}{dt}(\mathbf{r} \cdot \mathbf{r}) = 2 \mathbf{r} \cdot \frac{d}{dt} \mathbf{r}$ , which is zero, the distance of the particle to the origin stays constant. Thus, it moves on the surface of a sphere.
- b) False: If the particle motion is given by  $\mathbf{r}(t) = t \mathbf{k}$ , then it moves on a line and not a sphere.
- c) True: See the preceding answer. In fact, any  $\mathbf{r}(t)$  of the form  $\mathbf{r}(t) = \mathbf{r}_0 + t \mathbf{k}$  will do.
- d) True: Write  $\mathbf{r}(t) = (a(t), b(t), c(t))$ , so  $\mathbf{k} \times \mathbf{r} = (-b(t), a(t), 0)$  and  $\mathbf{k} \times \frac{d}{dt} \mathbf{r} = (-\frac{d}{dt} b(t), \frac{d}{dt} a(t), 0)$ . As this is orthogonal to  $\mathbf{k} \times \mathbf{r}$ , so  $a \frac{d}{dt} a + b \frac{d}{dt} b = 0$ , which says that  $2^{-1} \frac{d}{dt} (a^2 + b^2) = 0$  so  $a^2 + b^2$  is constant. Thus, the x and y coordinates of the particle move on a circle while the z coordinate can do what it likes. This puts the motion on a circular cylinder.

7. Answer: 0. The vector  $\mathbf{u} \times \mathbf{v}$  is perpendicular to both  $\mathbf{u}$  and  $\mathbf{v}$ .