

## Homework 14: Tangent lines and planes

This homework is due Friday, 10/17 resp Thursday 10/16.

1 The equation  $f(x, y) = x^4y^2 + 5xy^5 = 26$  defines a curve in the  $xy$ -plane. Find the tangent line  $ax + by = d$  to the curve at  $(2, 1)$  by computing the gradient  $\nabla f(x, y) = \langle a, b \rangle$ , and then plugging in the point to get the constant  $d$ .

2 a) Find an equation of the tangent plane to the surface  $z = 3(x - 1)^2 + 2(y + 3)^2 + 7$  at the point  $(2, -2, 12)$ .

b) Find an equation of the tangent plane to the surface

$$z = \log(x - 2y)$$

(with  $\log = \ln$  as natural log as usual) at the point  $(3, 1, 0)$ .

3 a) Find an equation of the tangent plane to the parametric surface

$$\vec{r}(u, v) = \langle u^2, v^2, uv \rangle$$

at the point  $(u, v) = (1, 1)$ .

b) The surface satisfies the equation  $xy - z^2 = 0$ . Find the tangent plane to this surface at the same point  $(x, y, z) = (1, 1, 1)$  by computing the gradient.

4 Find an equation of the tangent plane and the normal line to the surface  $x - z - 4 \arctan(yz) = 0$  through the point  $(1 + \pi, 1, 1)$ .

5 Show that the ellipsoid  $6x^2 + 4y^2 + 2z^2 = 18$  and the sphere  $x^2 + y^2 + z^2 - 8x - 6y - 8z + 24 = 0$  are tangent to each other at the point  $(1, 1, 2)$ , meaning that they have the same tangent plane at that point.

## Main definitions

The **gradient** in two dimensions is  $\nabla f(x, y) = \langle f_x(x, y), f_y(x, y) \rangle$ . In three dimensions, it is  $\nabla f(x, y, z) = \langle f_x(x, y, z), f_y(x, y, z), f_z(x, y, z) \rangle$ .

From the chain rule, we can deduce:

**Theorem:** Gradients are orthogonal to level curves and level surfaces.

The tangent line through  $(x_0, y_0)$  to a level curve  $f(x, y) = c$  is  $ax + by = d$ , where  $\nabla f(x_0, y_0) = \langle a, b \rangle$  and  $d$  is obtained by plugging in the point.

The tangent plane through  $(x_0, y_0, z_0)$  to a level surface  $f(x, y, z) = C$  is  $ax + by + cz = d$ , where  $\nabla f(x_0, y_0, z_0) = \langle a, b, c \rangle$  and  $d$  is obtained by plugging in the point.

We can compute tangent planes also for parametrized surfaces  $\vec{r}(u, v)$  because the vectors  $\vec{r}_u, \vec{r}_v$  are velocity vectors of grid curves and so tangent to the surface. Get the normal vector  $\vec{n} = \vec{r}_u \times \vec{r}_v = \langle a, b, c \rangle$  and then get  $ax + by + cz = d$ , where  $d$  is obtained by plugging in the point  $\vec{r}(u_0, v_0)$ . Here is how to plot parametric surfaces:

```
ParametricPlot3D [{u, v^2, u v}, {u, 0, 1}, {v, 0, 1}]
```