

Name:

MWF 9 Koji Shimizu
MWF 10 Can Kozcaz
MWF 10 Yifei Zhao
MWF 11 Oliver Knill
MWF 11 Bena Tshishiku
MWF 12 Jun-Hou Fung
MWF 12 Chenglong Yu
TTH 10 Jameel Al-Aidroos
TTH 10 Ziliang Che
TTH 10 George Melvin
TTH 11:30 Jake Marcinek
TTH 11:30 George Melvin

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3, we need to see **details** of your computation.
- All functions can be differentiated arbitrarily often unless otherwise specified.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110

Mark for each of the 20 questions the correct letter. No justifications are needed.)

T

F

The
(ρ , t)

Solution:

Make a picture and look at the angles. The θ angle is false.

- 2) T F If $|\vec{v} \times \vec{w}| = 0$ then $\vec{v} = \vec{0}$ or $\vec{w} = \vec{0}$.

Solution:

No, the vectors can be parallel without being zero.

- 3) T F The surface $z^2 + 4y^2 = x^2 + 1$ is a two sheeted hyperboloid.

Solution:

It is a deformed one-sheeted hyperboloid.

- 4) T F The surface $4x^2 - 4x + y^2 - 2y - 120 = -z^2$ is an ellipsoid.

Solution:

Complete the square

- 5) T F The parametrized lines $\vec{u}(t) = \langle 1 + 2t, 2 - 5t, 1 + t \rangle$ and $\vec{v}(t) = \langle 3 - 4t, -3 + 10t, 2 - 2t \rangle$ are the same line.

Solution:

The vectors are parallel, and both lines go through the same point.

- 6) T F The surface $\sin(x) = z$ contains lines which are parallel to the y -axis.

Solution:

One can translate the surface in the y direction.

- 7) T F If $\vec{u} \cdot \vec{v} = 0$, $\vec{v} \cdot \vec{w} = 0$ and \vec{v} is not the zero vector, then $\vec{u} \cdot \vec{w} = 0$.

Solution:

The assumption means that \vec{v} is perpendicular to \vec{u} and \vec{w} . But that does not mean that \vec{u} and \vec{w} are perpendicular.

- 8) T F The curvature of a curve depends upon the speed at which one travels upon it.

Solution:

The curvature does not depend on the parametrization.

- 9) T F Two lines in space that do not intersect must be parallel.

Solution:

They can be skew.

- 10) T F A line in space can intersect an elliptic paraboloid in 4 points.

Solution:

It can only intersect it in 2 points or 1 point or avoid it at all.

- 11) T F If $\vec{u} \times \vec{v} = 0$ and $\vec{u} \cdot \vec{v} = 0$, then one of the vectors \vec{u} and \vec{v} is zero.

Solution:

A vector which is both parallel and perpendicular to another vector can only be the zero vector.

- 12) T F If the velocity vector $\vec{r}'(t)$ and the acceleration vector $\vec{r}''(t)$ of a curve are parallel at time $t = 1$, then the curvature $\kappa(t)$ of the curve is zero at time $t = 1$.

Solution:

You can see this from the formula $\kappa = |r'(t) \times r''(t)|/|r'(t)|^3$. You can also think about it as follows. Assume the curvature were $\kappa = 1/r$. Then you as well locally move on a circle with radius r . But the acceleration has now a component perpendicular to your velocity vector. But we assumed there is no such acceleration.

- 13)

T	F
---	---

 If the speed of a parametrized curve is constant over time, then the curvature of the curve $\vec{r}(t)$ is zero.

Solution:

It would be true if the velocity would be constant over time. But we can move on a circle with constant speed.

- 14)

T	F
---	---

 The length of the vector projection of a vector \vec{v} onto a vector \vec{w} is always equal to the length of the vector projection of \vec{w} onto \vec{v} .

Solution:

If the lengths of \vec{v} and \vec{w} are the same, then the statement is true. In general, it is not.

- 15)

T	F
---	---

 A quadric $ax^2 + by^2 + cz^2 = 1$ is contained in the interior of a sphere $x^2 + y^2 + z^2 < 100$, then the constants a, b, c are all positive and the quadric is an ellipsoid.

Solution:

If any of the constants would become negative, the quadric becomes unbounded.

- 16)

T	F
---	---

 There is a hyperboloid of the form $ax^2 + by^2 - cz^2 = 1$ which has a trace which is a parabola.

Solution:

Traces are either hyperbola or ellipses.

- 17)

T	F
---	---

 The set of points in space which have distance 1 from the line $x = y = z$ form a cylinder.

Solution:

Yes, if the the line is the z -axis, then $x^2 + y^2 = 1$ is the equation of the cylinder.

- 18) T F The velocity vector of a parametric curve $\vec{r}(t)$ always has constant length.

Solution:

This is only true for an arc length parametrization.

- 19) T F The volume of a parallelepiped spanned by $\vec{u}, \vec{v}, \vec{w}$ is $|(\vec{u} \times \vec{v}) \times \vec{w}|$.

Solution:

The triple scalar product contains also a dot product.



- 20) T F The equation $x^2 + y^2/4 = 1$ in space describes an ellipsoid.


Solution:

The equation describes an elliptical cylinder.


Problem 2a) (3 points)

Match the equation with their graphs. No justifications are needed.





III



IV


Enter I,II,III,IV here	Equation
	$z = \sin(5x) \cos(2y)$
	$z = \cos(y^2)$
	$z = e^{-x^2-y^2}$
	$z = e^x$

Solution:

Enter I,II,III,IV here	Equation	Justification
III	$z = \sin(5x) \cos(2y)$	two traces show waves
II	$z = \cos(y^2)$	no x dependence, periodic in y
IV	$z = e^{-x^2-y^2}$	has a maximum at (0,0)
I	$z = e^x$	no y dependence, monotone in x

Problem 2b) (4 points)

Match the contour maps with the corresponding functions $f(x, y)$ of two variables. No justifications are needed.



Enter I,II,III,IV,V or VI here	Function $f(x, y)$
	$f(x, y) = \sin(x)$
	$f(x, y) = x^2 + 2y^2$
	$f(x, y) = x + y $
	$f(x, y) = \sin(x) \cos(y)$
	$f(x, y) = xe^{-x^2-y^2}$
	$f(x, y) = x^2/(x^2 + y^2)$

Solution:

Enter I,II,III,IV,V or VI here	Function $f(x, y)$
III	$f(x, y) = \sin(x)$
II	$f(x, y) = x^2 + 2y^2$
V	$f(x, y) = x + y $
IV	$f(x, y) = \sin(x) \cos(y)$
I	$f(x, y) = xe^{-x^2-y^2}$
VI	$f(x, y) = x^2/(x^2 + y^2)$

Problem 2c) (3 points)

Match the following points in cartesian coordinates with the points in spherical coordinates:

a) $(x, y, z) = (\sqrt{2}, 0, 0)$

b) $(x, y, z) = (0, \sqrt{2}, 0)$

c) $(x, y, z) = (0, 0, \sqrt{2})$

d) $(x, y, z) = (1, 1, 0)$

e) $(x, y, z) = (1, 0, 1)$

f) $(x, y, z) = (0, 1, 1)$

1) $(\rho, \phi, \theta) = (\sqrt{2}, 0, 0)$.

2) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/2, \pi/4)$.

3) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/2, 0)$.

4) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/2, \pi/2)$.

5) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/4, \pi/2)$.

6) $(\rho, \phi, \theta) = (\sqrt{2}, \pi/4, 0)$.

Solution:

$a = 3, b = 4, c = 1, d = 2, e = 6, f = 5$

Problem 3) (10 points)

- a) (7 points) Find a parametric equation for the line which is the intersection of the two planes $2x - y + 3z = 9$ and $x + 2y + 3z = -7$.
- b) (3 points) Find a plane perpendicular to both planes given in a) which has the additional property that it passes through the point $P = (1, 1, 1)$.

Solution:

- a) We get the direction of the line by taking the cross product of $\langle 2, -1, 3 \rangle$ and $\langle 1, 2, 3 \rangle$ which is $\langle -9, -3, 5 \rangle$. To find a point in both lines, subtract one from the other to get $x - 3y = 16$. If $z = 0$, then $2x - y = 9$ and $x + 2y = -7$ so that $x = 11/5, y = -23/5$. The parametric equations are $(x, y, z) = (11/5, -23/5, 0) + t\langle -9, -3, 5 \rangle$.
- b) Plug in the coordinates $(x, y, z) = (1, 1, 1)$ of the point to get the constant $-9x - 3y + 5z = -7$.

Problem 4) (10 points)

Given the vectors $\vec{v} = \langle 1, 1, 0 \rangle$ and $\vec{w} = \langle 0, 0, 1 \rangle$ and the point $P = (2, 4, -2)$. Let Σ be the plane which goes through the origin $(0, 0, 0)$ and which contains the vectors \vec{v} and \vec{w} . Let S be the unit sphere $x^2 + y^2 + z^2 = 1$.

- a) (6 points) Compute the distance from P to the plane Σ .
- b) (4 points) Find the shortest distance from P to the sphere S .

Solution:


- a) $\Sigma : x - y = 0, n = \langle 1, -1, 0 \rangle$. The point $Q = (0, 0, 0)$ is on the plane. $\vec{PQ} \cdot \vec{n} / |\vec{n}| = \langle 2, 4, -2 \rangle \cdot \langle 1, -1, 0 \rangle / \sqrt{2} = 2 / \sqrt{2} = \sqrt{2}$ is the distance.
- b) $\sqrt{4 + 16 + 4} = \sqrt{24} = 2\sqrt{6}$ is the distance to the origin. So the distance to the sphere is 1 less. The answer is $\sqrt{24} - 1$.

Problem 5) (10 points)

- a) (6 points) Find an equation for the plane through the points $A = (0, 1, 0), B = (1, 2, 1)$ and $C = (2, 4, 5)$.
- b) (4 points) Given an additional point $P = (-1, 2, 3)$, what is the volume of the tetrahedron

which has A, B, C, P among its vertices.

A useful fact which you can use without justification in b): the volume of the tetrahedron is $1/6$ of the volume of the parallelepiped which has $AB, AC,$ and AP among its edges.



Solution:

a) The vectors $\vec{v} = \vec{AB} = \langle 1, 1, 1 \rangle$ and $\vec{w} = \vec{AC} = \langle 2, 3, 5 \rangle$ are in the plane. Their cross product is $\vec{n} = \langle 2, -3, 1 \rangle$. This vector is perpendicular to the plane. The equation of the plane is therefore $2x - 3y + z = d$. Plugging in one point like A , gives $d = -3$.

b) With the vector $\vec{u} = \vec{AP} = \langle -1, 1, 3 \rangle$, one can express the volume of the parallelepiped as $|\llbracket \vec{u}, \vec{v}, \vec{w} \rrbracket| = |\vec{u} \cdot \vec{n}| = |\langle -1, 1, 3 \rangle \cdot \langle 2, -3, 1 \rangle| = |2| = 2$. The volume of the tetrahedron is $2/6 = 1/3$.

Problem 6) (10 points)

The parametrized curve $\vec{u}(t) = \langle t, t^2, t^3 \rangle$ (known as the "twisted cubic") intersects the parametrized line $\vec{v}(s) = \langle 1 + 3s, 1 - s, 1 + 2s \rangle$ at a point P . Find the angle of intersection.

Solution:

The curves intersect at $P = (1, 1, 1)$ with $t = 1, s = 0$. So, it remains to find the angle between the velocity vectors $\vec{u}'(1) = \langle 1, 2, 3 \rangle$ and $\vec{v}'(0) = \langle 3, -1, 2 \rangle$, which is 60 degrees.

Problem 7) (10 points)

Let $\vec{r}(t)$ be the space curve $\vec{r}(t) = (\log(t), 2t, t^2)$, where $\log(t)$ is the natural logarithm (denoted by $\ln(t)$ in some textbooks).

a) What is the velocity and what is the acceleration at time $t = 1$?

b) Find the length of the curve from $t = 1$ to $t = 2$.

Solution:

$$\text{a) } \vec{v}(t) = \vec{r}'(t) = \langle 1/t, 2, 2t \rangle.$$

$$\vec{v}(1) = \langle 1, 2, 2 \rangle$$

$$\vec{a}(t) = \vec{r}''(t) = \langle -1/t^2, 0, 2 \rangle.$$

$$\vec{a}(1) = \langle -1, 0, 2 \rangle.$$


$$\text{b) } \int_1^2 \sqrt{1/t^2 + 4 + 4t^2} dt = \int_1^2 1/t + 2t dt = \log(t) + t^2 \Big|_1^2 = \log(2) + 3.$$

Problem 8) (10 points)

A planar mirror in space contains the point $P = (4, 1, 5)$ and is perpendicular to the vector $\vec{n} = \langle 1, 2, -3 \rangle$. The light ray $\vec{QP} = \vec{v} = \langle -3, 1, -2 \rangle$ with source $Q = (7, 0, 7)$ hits the mirror plane at the point P .

a) (4 points) Compute the projection $\vec{u} = \vec{P}_{\vec{n}}(\vec{v})$ of \vec{v} onto \vec{n} .

b) (6 points) Identify \vec{u} in the figure and use it to find a vector parallel to the reflected ray.



Solution:

a) $P_{\vec{n}}(\vec{v}) = \frac{(\vec{n} \cdot \vec{v})}{|\vec{n}|^2} \vec{n} = (\langle 1, 2, -3 \rangle \cdot \langle -3, 1, -2 \rangle) / 14 \vec{n} = (5/14) \langle 1, 2, -3 \rangle$.

b) With \vec{u} we can get the reflected vector \vec{w} because $\vec{w} - \vec{v} = -2\vec{u}$ so that $\vec{w} = \vec{v} - 2\vec{u}$. Note that \vec{u} points down towards the mirror.

Problem 9) (10 points)

We know the acceleration $\vec{r}''(t) = \langle 2, 1, 3 \rangle + t \langle 1, -1, 1 \rangle$ and the initial position $\vec{r}(0) = \langle 0, 0, 0 \rangle$ and initial velocity $\vec{r}'(0) = \langle 11, 7, 0 \rangle$ of an unknown curve $\vec{r}(t)$. Find $\vec{r}(6)$.

Solution:

$$\vec{r}'(t) = \int_0^t \langle 2+t, 1-t, 3+t \rangle dt + \vec{r}'(0) = \langle 11+2t+t^2/2, 7+t-t^2/2, 3t+t^2/2 \rangle$$

$$\vec{r}(t) = \int_0^t \langle 2t+t^2/2, t-t^2/2, 3t+t^2/2 \rangle dt + \vec{r}(0) = \langle 11t+t^2+t^3/6, 7t+t^2/2-t^3/6, 3t^2/2+t^3/6 \rangle$$

Plug in the time $t = 6$ gives = $\langle 138, 24, 90 \rangle$.

Problem 10) (10 points)

Intersecting the elliptic cylinder $x^2 + y^2/4 = 1$ with the plane $z = \sqrt{3}x$ gives a curve in space.

a) (3 points) Find the parametrization of the curve.

b) (3 points) Compute the unit tangent vector \vec{T} to the curve at the point $(0, 2, 0)$.

c) (4 points) Write down the arc length integral and evaluate the arc length of the curve.

Solution:

a) With $x = \sin(t)$, $y = 2 \cos(t)$, $z = \sqrt{3} \sin(t)$, we check $x^2 + y^2/4 = \sin^2(t) + \cos^2(t) = 1$. The parametrization is $\vec{r}(t) = \langle \sin(t), 2 \cos(t), \sqrt{3} \sin(t) \rangle$.

b) Compute $\vec{r}'(t) = \langle \cos(t), -2 \sin(t), \sqrt{3} \cos(t) \rangle$, the speed $|\vec{r}'(t)| = 2$ and $\vec{T}(t) = \vec{r}'(t)/|\vec{r}'(t)| = \langle \cos(t)/2, -\sin(t), \sqrt{3} \cos(t)/2 \rangle$.

c) $|\vec{r}'(t)| = 2$. The length is $\int_0^{2\pi} |\vec{r}'(t)| dt = \int_0^{2\pi} 2 dt = \boxed{4\pi}$.

