

Name:

MWF 9 Koji Shimizu
MWF 10 Can Kozcaz
MWF 10 Yifei Zhao
MWF 11 Oliver Knill
MWF 11 Bena Tshishiku
MWF 12 Jun-Hou Fung
MWF 12 Chenglong Yu
TTH 10 Jameel Al-Aidroos
TTH 10 Ziliang Che
TTH 10 George Melvin
TTH 11:30 Jake Marcinek
TTH 11:30 George Melvin

- Start by printing your name in the above box and **check your section** in the box to the left.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader cannot be given credit.
- **Show your work.** Except for problems 1-3, we need to see **details** of your computation.
- All functions can be differentiated arbitrarily often unless otherwise specified.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have 90 minutes time to complete your work.


1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
Total:		110

Problem 1) True/False questions (20 points), no justifications needed

- | | | |
|-----|--|---|
| 1) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | The length of the gradient $\nabla f(0,0)$ is the maximal directional derivative $ D_{\vec{v}}f(0,0) $ among all unit vectors $\vec{v}$. |
| 2) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | The relation $f_{xxyyxx} = f_{xyxyyx}$ holds everywhere for $f(x,y) = \cos(\exp(x^{10}) + \sin(x-y))$. |
| 3) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | $\int_0^4 \int_0^{4x} f(x,y) dydx = \int_0^{16} \int_{y/4}^{16} f(x,y) dxdy$. |
| 4) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | $g(x,y) = \int_y^0 \int_0^x f(s,t) dsdt$ satisfies $g_{xy} = -f(x,y)$. |
| 5) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If $\vec{r}(u,v)$ is a parametrization of the level surface $f(x,y,z) = c$, then $\nabla f(\vec{r}(u,v)) \cdot \vec{r}_v(u,v) = 0$. |
| 6) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If $D_{\langle 1/\sqrt{2}, 1/\sqrt{2} \rangle} f(a,b) = 3$ and $D_{\langle 1/\sqrt{2}, -1/\sqrt{2} \rangle} f(a,b) = 5$, then $D_{\vec{v}}f(a,b) \geq 0$ for all unit directions $\vec{v}$. |
| 7) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | Given a parametrization $\vec{r}(t)$ of a curve and a function $f(x,y)$ we have $\frac{d}{dt}f(\vec{r}(2t)) = 2\nabla f(\vec{r}(t)) \cdot \vec{r}'(t)$ at $t = 0$. |
| 8) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If $u(t,x)$ solves both the heat and wave equation, then $u_t = c u_{tt}$ for some constant c . |
| 9) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If the Lagrange multiplier λ at a solution to a Lagrange problem is negative then this point is neither a maximum nor a minimum. |
| 10) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | The equation $f_x^2 + f_y^2 + f_z^2 = 1$ is an example of a partial differential equation. |
| 11) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If the discriminant D of $f(x,y)$ is zero at $(0,0)$ then $\nabla f(0,0) = \langle 0,0 \rangle$. |
| 12) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If $f(x,y,z) = 0$ describes the unit sphere, then the gradient ∇f points outwards. |
| 13) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If $f(x,y)$ is a continuous function then $\int_0^2 \int_0^1 f(x,y) dxdy = \int_0^2 \int_0^1 f(y,x) dxdy$. |
| 14) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | The point $(5,5,5)$ is a critical point of $f(x,y,z) = x + y + z$. |
| 15) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | Assume $\nabla f(0,0) = \langle 0,0 \rangle$ with discriminant $D > 0$, then $-f(x,y)$ has the same critical point $(0,0)$ with discriminant $D < 0$. |
| 16) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | $\int \int_R \nabla f ^2 dxdy$ is the surface area of the cubic paraboloid $z = f(x,y) = x^3 + y^3$ defined over the region R . |
| 17) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If $D(x,y)$ is the discriminant of f at (x,y) then the following poetic formula of the directional derivative of the discriminant holds: $D_{\langle 1,0 \rangle} D = \partial_x D$. |
| 18) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | Assume $f(x,y) = -x^2 + y^4$ and a curve $\vec{r}(t)$ satisfying $\vec{r}'(t) = \nabla f(\vec{r}(t))$, then $\frac{d}{dt}f(\vec{r}(t)) \geq 0$ for all t . |
| 19) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | The Lagrange equations for extremizing $f(x,y)$ under the constraint $g(x,y) = c$ have the same solutions as the Lagrange equations for extremizing $F = f + g$ under the constraint $g = c$. |
| 20) | <div style="display: inline-block; border: 1px solid black; padding: 2px 10px;">T</div> <div style="display: inline-block; border: 1px solid black; padding: 2px 10px; margin-left: 10px;">F</div> | If f is a maximum under the constraint $g = 1$ at $(0,0)$, and $(0,0)$ is not a critical point for both f and g , then the level curves of f and g have the same tangent line at $(0,0)$. |

Problem 2) (10 points) No justifications needed

a) (6 points) Match the regions with the integrals. Each integral matches one region $A - F$.



Enter A-F	Integral
	$\int_{-1}^1 \int_{ y }^1 f(x,y) \, dx dy$
	$\int_0^1 \int_0^x f(x,y) \, dy dx$
	$\int_{-1}^0 \int_{-y}^1 f(x,y) \, dx dy$
	$\int_0^1 \int_{x-1}^0 f(x,y) \, dy dx$
	$\int_0^1 \int_{x-1}^{1-x} f(x,y) \, dy dx$
	$\int_0^1 \int_x^1 f(x,y) \, dy dx$

b) (4 points) Name the partial differential equations correctly. Each equation matches one name.



Fill in 1-4	Name
	Laplace
	Wave
	Transport
	Heat

Equation number	Formula for PDE
1	$\frac{\partial}{\partial t} u - \frac{\partial}{\partial y} u = 0$
2	$\frac{\partial}{\partial t} u - \frac{\partial^2}{\partial x^2} u = 0$
3	$\frac{\partial^2}{\partial t^2} u - \frac{\partial^2}{\partial x^2} u = 0$
4	$\frac{\partial^2}{\partial x^2} u + \frac{\partial^2}{\partial y^2} u = 0$

Problem 3) (10 points)

a) (7 points) The following contour map is inspired by a cubistic style of Picasso. Each of the points A-H fit exactly once

A point where $f_x \neq 0, f_y = 0$	
A saddle point of f	
A local maximum of f	
A critical point with $D = 0$	
A local minimum	
A point with $f_y \neq 0, f_x = 0$	
$ \nabla f $ maximal among A-H	
Point where $D_{\langle -1,1 \rangle / \sqrt{2}} f = 0$	



The painting "Pigeon with Green Peas" by Pablo Picasso was stolen in 2010. The thief got scared and disposed it to trash shortly after the theft. The garbage was emptied and taken away, the painting lost for ever. Or the thief had been clever ...

b) (3 points) Given a function $f(x, y)$ and a curve $\vec{r}(t)$. Let L be the linearization of f at $\vec{r}(0)$. Each of the following 3 vectors $\vec{a}, \vec{b}, \vec{c}$ is placed exactly twice in the puzzle to the right below.

$\vec{a} = \nabla f(\vec{r}(0))$
$\vec{b} = \vec{r}'(0)$
$\vec{c} = \vec{r}(t) - \vec{r}(0)$

$\frac{d}{dt} f(\vec{r}(t))_{t=0} =$		$\cdot$		
$L(\vec{r}(t)) =$	$f(\vec{r}(0)) +$		$\cdot$	
	$=$	$\lim_{t \rightarrow 0} \frac{1}{t}$		

Problem 4) (10 points)

a) (5 points) Find the tangent plane to the **skate board ramp**


$$z - f(x, y) = z - \sqrt{x^{30}y^3 + x} = 0$$

at the point $(1, 2, 3)$.

b) (5 points) Estimate

$$f(1.006, 1.98) = \sqrt{1.006^{30} \cdot 1.98^3 + 1.006}$$

by linearizing the function $f(x, y)$ at $(1, 2)$.



Problem 5) (10 points)


A **croissant** of length $2h$ and radius r in the shape of two cones has fixed volume

$$V(r, h) = \frac{2\pi r^2 h}{3} = 18.$$

Use Lagrange to find the values r and h for which the surface area

$$A(r, h) = 2\pi r \sqrt{r^2 + h^2}$$

is minimal. **Hint:** as you have seen in homework, it is much more convenient to minimize $f(r, h) = A(r, h)^2$ instead.




Problem 6) (10 points)

Find the local maxima, minima and saddle points of the **tadpole function**

$$f(x, y) = 3y^2 + 4x^3 + 2y^3 - 12x.$$

First tadpole: “What is your favorite book?” *Second tadpole:* “Metamorphosis by Kafka. What is your favorite year?” *First tadpole:* “Leap year!” *Both croak with laughter. Then, sick of frog jokes, they turn green.*




Problem 7) (10 points)

Find the area of the heart shaped polar region

$$(\theta - \pi)^2 \leq r \leq 2(\theta - \pi)^2$$

with $0 \leq \theta \leq 2\pi$.

*Warning: Valentine cards displaying
“You are my $r < (\theta - \pi)^2$!” do not always work.*




Problem 8) (10 points)

Mathematica 10 does not give an elementary expression for the integral

$$\int_0^1 \int_{\exp(y)}^e \frac{1}{\log(x)} dx dy ,$$

where $\log$ is the natural log. You can! “Humans are awesome 2014”! <https://www.youtube.com/watch?v=ZBCOMG2F2Zk>

The logarithmic integral $\text{Li}(x) = \int_0^x dt/\log(t)$ is important in number theory. It was Gauss who proposed first that the number $\pi(x)$ of primes smaller than x is about $\text{Li}(x)$. It is now known that $0.89 \text{Li}(x) \leq \pi(x) \leq 1.11 \text{Li}(x)$ for all large enough x . P.S. Mathematica can solve the double integral of course, but only if told to “FullSimplify”.




Problem 9) (10 points)

Compute the weighted surface area

$$\int \int_R (u^2 + v^2) |\vec{r}_u \times \vec{r}_v| du dv$$

of the monkey saddle parametrized by $\vec{r}(u, v) = \langle u, v, u^3 - 3uv^2 \rangle$ over the domain $R : u^2 + v^2 \leq 1$. This quantity is also known as the moment of inertia of the surface. Spin that monkey!



Problem 10) (10 points)

The following two integrals are called "Mad Max" integrals because they were written while watching that movie:

a) (5 points) Integrate

$$\int_0^1 \int_{\arcsin(y)}^{\pi/2} \frac{xy}{\sin(x)} dx dy .$$

b) (5 points) Integrate the double integral

$$\int \int_R \sin(x^2 + y^2) dx dy$$

where R is the disk of radius $\sqrt{\pi/2}$.

