

## Homework 22: Triple integrals

This homework is due Friday, 11/4 resp Tuesday 11/8.

- 1 a) Evaluate the iterated integral

$$\int_0^2 \int_0^z \int_0^{y^3} x \, dx \, dy \, dz .$$

- b) Which of the three following scrambled versions make sense too?

$$\int_0^2 \int_0^{y^3} \int_0^z x \, dx \, dz \, dy .$$

$$\int_0^z \int_0^2 \int_0^{y^3} x \, dx \, dz \, dy .$$

$$\int_0^2 \int_0^y \int_0^{z^3} x \, dx \, dz \, dy .$$

- 2 Evaluate the triple integral

$$\iiint_E 2yz \cos(x^5) \, dV ,$$

where

$$E = \{(x, y, z) \mid 0 \leq x \leq \pi/2, 0 \leq y \leq x, x \leq z \leq 2x\} .$$

- 3 Evaluate the triple integral

$$\iiint_E xy \, dV ,$$

where  $E$  is bounded by the parabolic cylinders  $y = 3x^2$  and  $x = 3y^2$  and the planes  $z = 0$  and  $z = x + y$ .

- 4 Use a triple integral to find the volume of the given solid enclosed by the paraboloid  $x = y^2 + z^2$  and the plane  $x = 25$ .

- 5 Find the moment of inertia

$$I = \iiint_E (x^2 + y^2) \, dz \, dx \, dy$$

about the  $z$ -axis of the solid  $E : 0 \leq z \leq r^3 \leq 8$ .

## Main definitions

If  $f(x, y, z)$  is a function and  $E$  is a **solid**, then  $\iiint_E f(x, y, z) dV$  is defined as the  $n \rightarrow \infty$  limit of the Riemann sum

$$\frac{1}{n^3} \sum_{\left(\frac{i}{n}, \frac{j}{n}, \frac{k}{n}\right) \in E} f\left(\frac{i}{n}, \frac{j}{n}, \frac{k}{n}\right).$$

Like  $dA = dx dy$  was a symbol for a small area,  $dV = dx dy dz$  indicates a small volume. Triple integrals are solved as a nested list of single integrals.

If  $f(x, y, z) = 1$  then  $\iiint_E 1 dx dy dz$  is the volume of the solid

A common situation is where the triple integral is reduced to a double integral

$$\int \int_R \left[ \int_{g(x,y)}^{h(x,y)} f(x, y, z) dz \right] dx dy .$$

This is by far the most common case. For example, if  $g(x, y) = 0$  and  $f(x, y, z) = 1$ , then

$$\int \int_R \left[ \int_0^{h(x,y)} 1 dz \right] dx dy = \int \int_R h(x, y) dx dy$$

is the signed volume of the solid under the graph of  $h$ . In variable calculus, where you were sometimes able to compute triple integrals by reducing to a single integral. You would have written  $\int_a^b A(z) dz$  to compute the volume of a solid sandwiched between  $z = a$  and  $z = b$  for which the area of the cross section at height  $z$  is  $A(z)$ . In multi variable calculus we are much more flexible as we can now also reduce to a double integral.