

Homework 17: Lagrange multipliers

This homework is due Friday, 10/20 resp Tuesday 10/24.

We look at a melon shaped candy. The outer radius is x , the inner is y . Assume we want to extremize the **sweetness function** $f(x, y) = -x^2 + 2y^2$ under the con-

- 1 constraint that $g(x, y) = x - y = 2$. Since this problem is so tasty, we require you to use the most yummy method known to mankind: the **Lagrange** method! Is your solution a minimum or maximum?




Solution:

The gradients of f and g are $\nabla f = \langle -2x, 2y \rangle$, $\nabla g = \langle 1, -1 \rangle$. The Lagrange equations are $-2x = \lambda$, $2y = -\lambda$. Elimination λ gives $x = 2y$. Plug this into the constraint. We get $\boxed{x = 4, y = 2}$ as the minimum.

The material to build a traffic light is $g(x, y) = 6 + 6\pi xy + 3\pi x^2 = 12$ is fixed (the radius of each cylinder is x and the height is y and the constant 6 is the material for the back plate). We want to build a light for which the shaded region with volume $f(x, y) = 3\pi x^2 y$ is maximal. Use the Lagrange method.

2



Solution:

The Lagrange equations $\nabla f = \lambda \nabla g, g = 12$ are

$$6\pi xy = \lambda(6\pi y + 6\pi x)$$

$$3\pi x^2 = \lambda(6\pi x)$$

$$2\pi xy + \pi x^2 = 6$$

Eliminating λ from the first two equations gives $x = y$. Plugging into the constraint gives $x = \sqrt{2/(3\pi)} = y$. The maximal value is $3\pi x^2 y = 2\sqrt{2/(3\pi)} = 0.921\dots$

- 3 The situation is the same, but we have two Lagrange multipliers (see box to the right). Use Lagrange multipliers to find the maximum and minimum f under the two constraints:

$$f(x, y, z) = 3x - y - 3z;$$

$$g(x, y, z) = x + y - z = 0$$

$$h(x, y, z) = x^2 + 2z^2 = 1 .$$

Solution:

By the method of Lagrange multipliers,

$$\langle 3, -1, -3 \rangle = \lambda \langle 1, 1, -1 \rangle + \mu \langle 2x, 0, 4z \rangle$$

The “middle” equation tells us that $\lambda = -1$. Hence, $2x\mu - 1 = 3$ and $4z\mu + 1 = -3$. In particular, $x\mu = 2$ and $-z\mu = 1$ so $x = -2z$. Plugging $x = -2z$ into the first constraint, we find $y = 3z$. Plugging $x = -2z$ in the second constraint, we see that $6z^2 = 1$. Thus the two critical points are $(\frac{2}{\sqrt{3}}, -\frac{3}{\sqrt{6}}, -\frac{1}{\sqrt{6}})$ and $(-\frac{2}{\sqrt{6}}, \frac{3}{\sqrt{6}}, \frac{1}{\sqrt{6}})$. The first gives the maximal value of $\frac{12}{\sqrt{6}}$ while the second gives the minimum value of $-\frac{12}{\sqrt{6}}$.

- 4 a) Use Lagrange multipliers to prove that the triangle with maximum area that has a given perimeter 2 is equilateral.
- b) What are the minima? Why does the Lagrange method not establish them?

Hint: Use Heron’s formula $A = \sqrt{s(s-x)(s-y)(s-z)}$, where $s = 1$ and x, y, z are the lengths of the sides.

Solution:

a) To maximize the area A , it is sufficient to maximize A^2 . We do so since the formula for A^2 does not involve a square root. If the perimeter p is fixed, then so is s . Hence, it is sufficient to maximize

$$\begin{aligned} f(x, y, z) &= s(s-x)(s-y)(s-z) \\ &= (y+z-x)(z+x-y)(x+y-z) \end{aligned}$$

provided $g(x, y, z) = x + y + z = 2s = 2$. By Lagrange multipliers,

$$\nabla f = \lambda \langle 1, 1, 1 \rangle.$$

Thus, $\frac{\partial f}{\partial x} = \frac{\partial f}{\partial y} = \frac{\partial f}{\partial z}$. We compute:

$$0 = \frac{\partial f}{\partial x} - \frac{\partial f}{\partial y} = (x+y-z) \cdot -4(x-y)$$

By the triangle inequality, $x+y > z$, so $x = y$. Similarly, $x = z$ forcing the triangle to be equilateral.

b) The minimal area is obtained if either $x = s$ or $y = s$ or $z = s$ which corresponds to a degenerate triangle with zero area. At these minima, the function f is no more differentiable. Indeed, it is the boundary where the function is defined. The constraint $g = x + y + z = 2s$

- 5 Which pyramid of height h over a square $[-a, a] \times [-a, a]$ with surface area is $4a\sqrt{h^2 + a^2} + 4a^2 = 4$ has maximal volume $V(h, a) = 4ha^2/3$? By using new variables (x, y) and multiplying V with a constant, we get to the equivalent problem to maximize $f(x, y) = yx^2$ over the constraint $g(x, y) = x\sqrt{y^2 + x^2} + x^2 = 1$. Use the later variables.

Solution:

An elegant solution can be obtained by first simplifying the constraint and write $x^2(y^2 - x^2) - (1 - x^2)^2 = x^2y^2 + 2x^2 - 1 = 0$. Now the Lagrange equations are not so bad:

$$\begin{aligned}2xy &= \lambda(4x + 2xy^2) \\x^2 &= \lambda 2x^2y \\x^2y^2 + 2x^2 &= 1.\end{aligned}$$

Eliminating λ by cross multiplying the first two equations gives $4x^3y^2 = x^2(4x + 2xy^2)$. Since $x = 0$ is not possible, we can divide both sides by x^3 and get $y^2 = 2$. Plugging into the constraint, we get $x = 1/2$. Without this simplification, the Lagrange system would have become more complicated:

$$\begin{aligned}2xy &= \lambda(\sqrt{y^2 + x^2} + x^2/\sqrt{y^2 + x^2} + 2x) \\x^2 &= \lambda yx/\sqrt{y^2 + x^2} \\1 &= x\sqrt{y^2 + x^2} + x^2\end{aligned}$$

Main definitions

The system of equations $\nabla f(x, y) = \lambda \nabla g(x, y), g(x, y) = 0$ for the three unknowns x, y, λ are the **Lagrange equations**. λ is a **Lagrange multiplier**. The **two constraint case** appears only here in homework and is not covered in section $\nabla f(x, y, z) = \lambda \nabla g(x, y, z), g(x, y, z) = 0$ are the **Lagrange equations** $\nabla f(x, y, z) = \lambda \nabla g(x, y, z) + \mu \nabla h(x, y, z), g(x, y, z) = 0, h(x, y, z) = 0$ are the **Lagrange equations** with two constraints.

Lagrange theorem: Maxima or minima of f on the constraint $g = c$ are either solutions of the Lagrange equations or critical points of g .