

LEVEL CURVES


2D: If $f(x, y)$ is a function of two variables, then $f(x, y) = c = \text{const}$ is **curve** in the plane. It is called **contour curve** or **level curve**. For example, $f(x, y) = 4x^2 + 3y^2 = 1$ is an ellipse. Level curves allow to visualize functions of two variables $f(x, y)$.

LEVEL SURFACES.



3D: If $f(x, y, z)$ is a function of three variables and c is a constant then $f(x, y, z) = c$ is a surface in space. It is called a **contour surface** or a **level surface**. For example if $f(x, y, z) = 4x^2 + 3y^2 + z^2$ then the contour surfaces are ellipsoids.

EXAMPLE. Let $f(x, y) = x^2 - y^2$. The set $x^2 - y^2 = 0$ is the union of the sets $x = y$ and $x = -y$. The set $x^2 - y^2 = 1$ consists of two hyperbola with their tips at $(-1, 0)$ and $(1, 0)$. The set $x^2 - y^2 = -1$ consists of two hyperbola with their tips at $(0, \pm 1)$.

EXAMPLE. Let $f(x, y, z) = x^2 + y^2 - z^2$. $f(x, y, z) = 0, f(x, y, z) = 1, f(x, y, z) = -1$. The set $x^2 + y^2 - z^2 = 0$ is a **cone** rotational symmetric around the z -axes. The set $x^2 + y^2 - z^2 = 1$ is a **one-sheeted hyperboloid**, the set $x^2 + y^2 - z^2 = -1$ is a **two-sheeted hyperboloid**. (To see that it is two-sheeted note that the intersection with $z = c$ is empty for $-1 \leq z \leq 1$.)



CONTOUR MAP. Drawing several contour surfaces $\{f(x, y, z) = c\}$ produces a contour map.





The example shows the graph of the function $f(x, y) = \sin(xy)$. We draw the contour map of f : The curve $\sin(xy) = c$ is $xy = C$, where $C = \arcsin(c)$ is a constant. The curves $y = C/x$ are hyperbolas except for $C = 0$, where $y = 0$ is a line. Also the line $x = 0$ is a contour curve. The contour map is a family of hyperbolas and the coordinate axes.

TOPOGRAPHY. Topographical maps often show the curves of equal height. With the contour curves as information, it is usually already possible to get a good picture of the situation.



EXAMPLE. $f(x, y) = 1 - 2x^2 - y^2$. The contour curves $f(x, y) = 1 - 2x^2 + y^2 = c$ are the ellipses $2x^2 + y^2 = 1 - c$ for $c < 1$.




SPECIAL LINES. Level curves are encountered every day:



Isobars:	pressure
Isoclines:	direction

Isothermes:	temperature
Isoheight:	height

For example, the isobars to the right show the lines of constant temperature yesterday February 27 in the north east of the US.





A SADDLE. $f(x, y) = (x^2 - y^2)e^{-x^2 - y^2}$. We can here no more find explicit formulas for the contour curves $(x^2 - y^2)e^{-x^2 - y^2} = c$. Lets try our best:



- $f(x, y) = 0$ means $x^2 - y^2 = 0$ so that $x = y, x = -y$ are contour curves.
- On $y = ax$ the function is $g(x) = (1 - a^2)x^2e^{-(1+a^2)x^2}$.
- Because $f(x, y) = f(-x, y) = f(x, -y)$, the function is symmetric with respect to reflections at the x and y axes.

A SOMBRERO. The surface $z = f(x, y) = \sin(\sqrt{x^2 + y^2})$ is in cylindrical coordintes $z = \sin(r)$. It has circles as contour lines.



ABOUT CONTINUITY. In reality, one sometimes has to deal with functions which are not smooth or not continuous: For example, when plotting the temperature of water in relation to pressure and volume, one experiences **phase transitions**, an other example are water waves breaking in the ocean. Mathematicians have also tried to explain "catastrophic" events mathematically with a theory called "catastrophe theory". Discontinuous things are useful (for example in switches), or not so useful (for example, if something breaks).

DEFINITION. A function $f(x, y)$ is **continuous** at (a, b) if $f(a, b)$ is finite and $\lim_{(x, y) \rightarrow (a, b)} f(x, y) = f(a, b)$. The later means that that along any curve $\vec{r}(t)$ with $r(0) = (a, b)$, we have $\lim_{t \rightarrow 0} f(\vec{r}(t)) = f(a, b)$. Continuity for functions of more variables is defined in the same way.

EXAMPLE. $f(x, y) = (xy)/(x^2 + y^2)$. Because $\lim_{(x, x) \rightarrow (0, 0)} f(x, x) = \lim_{x \rightarrow 0} x^2/(2x^2) = 1/2$ and $\lim_{(x, 0) \rightarrow (0, 0)} f(0, x) = \lim_{(x, 0) \rightarrow (0, 0)} 0 = 0$. The function is not continuous.

EXAMPLE. $f(x, y) = (x^2y)/(x^2 + y^2)$. In polar coordinates this is $f(r, \theta) = r^3 \cos^2(\theta) \sin(\theta)/r^2 = r \cos^2(\theta) \sin(\theta)$. We see that $f(r, \theta) \rightarrow 0$ uniformly if $r \rightarrow 0$. The fuction is continuous.