



**FUNCTIONS II**

Maths21a, O. Knill

**FUNCTIONS OF ONE VARIABLES.** A function of one variable  $f(x)$  assigns to a variable  $x$  a real number  $f(x)$ . Example:  $f(x) = x \sin(x)$ . We can visualize a function by its **graph**  $\{(x, y), y = f(x)\}$  which is a curve in space.




**FUNCTIONS OF TWO VARIABLES.** A function of two variables  $f(x, y)$  assigns to two variables  $x, y$  a real number  $f(x, y)$ . Example:  $f(x, y) = \sin(xy)$ . It could describe a temperature distribution on a plate for example.



In the same way as functions of one variable were visualized by drawing the graph  $y = f(x)$  in space, we can visualize a function of three variables as a graph  $\{(x, y, z) \mid z = f(x, y)\}$  in space.

**FUNCTIONS OF THREE VARIABLES.** A function of three variables  $g(x, y, z)$  assigns to three variables  $x, y, z$  a real number  $g(x, y, z)$ . Example:  $g(x, y, z) = \sin(xyz)$ , temperature distribution in space. We can no more draw a graph of  $g$ . But we can visualize it differently by drawing surfaces  $g(x, y, z) = c$ , where  $c$  is constant.




**SURFACES.** Many surfaces can be described as a function of three variables  $g$ . The points  $(x, y, z)$  satisfying the equation  $g(x, y, z) = c$  for a surface. Examples.

**Graphs:**  $g(x, y, z) = z - f(x, y)$ . If  $g(x, y, z) = 0$ , then  $z = f(x, y)$  and the surface is a graph of a function of two variables.

**Planes:**  $ax + by + cz = d$  is a plane orthogonal to the vector  $\vec{n} = (a, b, c)$ . The equation says  $\vec{n} \cdot \vec{x} = d$ . If a point  $\vec{x}_0$  is on the plane, then  $\vec{n} \cdot \vec{x}_0 = d$ . The equation can also be written as  $\vec{n} \cdot (\vec{x} - \vec{x}_0) = 0$  which means that every vector  $\vec{x} - \vec{x}_0$  is orthogonal to  $\vec{n}$ .


**Quadrics:** If  $f(x, y, z) = ax^2 + by^2 + cz^2 + dxy + exz + fyz + gx + hy + kz + m$  the a surface  $f(x, y, z) = 0$  is called a **quadric**. Below are some examples.

QUARTIC




$p(x, y, z)$  degree 4 polynomial in  $x, y, z$

SEXTIC




$p(x, y, z)$  degree 6 polynomial in  $x, y, z$

DECIC




$p(x, y, z)$  degree 10 polynomial in  $x, y, z$

SPHERE




$$(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2$$

PARABOLOID




$$(x - a)^2 + (y - b)^2 - c = z$$

PLANE




$$ax + by + cz = d$$

HYPERBOLOID I




$$(x - a)^2 + (y - b)^2 - (z - c)^2 = r^2$$

CYLINDER




$$(x - a)^2 + (y - b)^2 = r^2$$

HYPERBOLOID II




$$(x - a)^2 + (y - b)^2 - (z - c)^2 = -r^2$$

ELLIPSOID




$$x^2/a^2 + y^2/b^2 + z^2/c^2 = 1$$

HYPERBOLIC PARABOLOID



$$x^2 - y^2 + z = 1$$

DEFORMED HYPERBOLOID



$$x^2/a^2 + y^2/b^2 - z^2/c^2 = 1$$

**TRACES.** To draw surfaces, it helps to look at the **traces**, the intersections of the surfaces with the coordinate planes  $x = 0, y = 0$  or  $z = 0$ .

**INTERCEPTS.** Other points to look at are the **intercepts**, the intersections of the surface with the coordinate axis. The traces are shown in the pictures of the quadrics above.

For example: for the **one-sided hyperboloid**  $x^2 + y^2 - z^2 = 1$  (called HYPERBOLOID I above), the  $z$ -trace is  $x^2 + y^2 = 1$ , a circle, the  $x$ -traces  $y^2 - z^2 = 1$  is a hyperbola as well as the  $y$ -trace  $x^2 - z^2 = 1$ .