


VECTORS, DOT PRODUCT **O. Knill, Maths21a**

VECTORS. Two points $P = (x_1, y_1)$, $Q = (x_2, y_2)$ in the plane define a **vector** $\vec{v} = \langle x_2 - x_1, y_2 - y_1 \rangle$. It points from P to Q and we can write $P + \vec{v} = Q$.

COMPONENTS. Points P in space are in one to one correspondence to vectors pointing from 0 to P . The numbers \vec{v}_i in a vector $\vec{v} = \langle v_1, v_2 \rangle$ are also called **components** or of the vector.

REMARKS: vectors can be drawn **everywhere** in the plane. If a vector starts at the origin O , then the vector $\vec{v} = \langle v_1, v_2 \rangle$ points to the point $\langle v_1, v_2 \rangle$. That's why one can identify points $P = (a, b)$ with vectors $\vec{v} = \langle a, b \rangle$. Two vectors which can be translated into each other are considered **equal**. They have the same components.

ADDITION, SUBTRACTION AND SCALAR MULTIPLICATION.



$$\vec{u} + \vec{v} = \langle u_1, u_2 \rangle + \langle v_1, v_2 \rangle = \langle u_1 + v_1, u_2 + v_2 \rangle$$


$$\vec{u} - \vec{v} = \langle u_1, u_2 \rangle - \langle v_1, v_2 \rangle = \langle u_1 - v_1, u_2 - v_2 \rangle$$

$$\lambda \vec{u} = \lambda \langle u_1, u_2 \rangle = \langle \lambda u_1, \lambda u_2 \rangle$$


BASIS VECTORS. The vectors $\vec{i} = \langle 1, 0 \rangle$, $\vec{j} = \langle 0, 1 \rangle$ are called **standard basis vectors** in the plane. In space, one has the basis vectors $\vec{i} = \langle 1, 0, 0 \rangle$, $\vec{j} = \langle 0, 1, 0 \rangle$, $\vec{k} = \langle 0, 0, 1 \rangle$. Every vector $\vec{v} = \langle v_1, v_2 \rangle$ in the plane can be written as $\vec{v} = v_1 \vec{i} + v_2 \vec{j}$. Every vector $\vec{v} = \langle v_1, v_2, v_3 \rangle$ in space can be written as $\vec{v} = v_1 \vec{i} + v_2 \vec{j} + v_3 \vec{k}$.

WHERE DO VECTORS OCCUR? Here are some examples:

Velocity: vectors appear in mechanics. For example, if $(f(t), g(t))$ is a point in the plane which depends on time t , then $\vec{v} = \langle f'(t), g'(t) \rangle$ is the **velocity vector** at the point $(f(t), g(t))$. We will look at this in more detail later in the course.




Forces: Some problems in statics involve the determination of a forces acting on objects. Forces are represented as vectors




Fields: electromagnetic or gravitational fields or velocity fields in fluids are described with vectors.



Qbits: in quantum computation, rather than working with bits, one deals with **qbits**, which are vectors.



Color can be written as a vector $\vec{v} = \langle r, g, b \rangle$, where r is **red**, g is **green** and b is **blue**. An other coordinate system is $\vec{v} = \langle c, m, y \rangle = (1 - r, 1 - g, 1 - b)$, where c is **cyan**, m is **magenta** and y is **yellow**.



SVG. Scalable Vector Graphics is an emerging standard for the web for describing two-dimensional graphics in XML.



VECTOR OPERATIONS: The addition and scalar multiplication of vectors satisfy "obvious" properties. There is no need to memorize them. We write $*$ here for multiplication with a scalar but usually, the multiplication sign is left out.

$\vec{u} + \vec{v} = \vec{v} + \vec{u}$	commutativity
$\vec{u} + (\vec{v} + \vec{w}) = (\vec{u} + \vec{v}) + \vec{w}$	additive associativity
$\vec{u} + \vec{0} = \vec{0} + \vec{u} = \vec{u}$	null vector
$r * (s * \vec{v}) = (r * s) * \vec{v}$	scalar associativity
$(r + s)\vec{v} = \vec{v}(r + s)$	distributivity in scalar
$r(\vec{v} + \vec{w}) = r\vec{v} + r\vec{w}$	distributivity in vector
$1 * \vec{v} = \vec{v}$	the one element

LENGTH. The length $|\vec{v}|$ of \vec{v} is the distance from the beginning to the end of the vector.

EXAMPLES. 1) If $\vec{v} = \langle 3, 4 \rangle$, then $|\vec{v}| = \sqrt{25} = 5$. 2) $|\vec{i}| = |\vec{j}| = |\vec{k}| = 1$, $|\vec{0}| = 0$.

UNIT VECTOR. A vector of length 1 is called a **unit vector**. If $\vec{v} \neq \vec{0}$, then $\vec{v}/|\vec{v}|$ is a unit vector.
EXAMPLE: If $\vec{v} = \langle 3, 4 \rangle$, then $\vec{v} = \langle 3/5, 4/5 \rangle$ is a unit vector, $\vec{i}, \vec{j}, \vec{k}$ are unit vectors.

PARALLEL VECTORS. Two vectors \vec{v} and \vec{w} are called **parallel**, if $\vec{v} = r\vec{w}$ with some constant r . The zero vector is parallel to any other vector.

DOT PRODUCT. The **dot product** of two vectors $\vec{v} = \langle v_1, v_2, v_3 \rangle$ and $\vec{w} = \langle w_1, w_2, w_3 \rangle$ is defined as

$$\vec{v} \cdot \vec{w} = v_1 w_1 + v_2 w_2 + v_3 w_3$$

Remark: in other sciences different notations are used: $\vec{v} \cdot \vec{w} = (\vec{v}, \vec{w})$ (mathematics) $\langle \vec{v} | \vec{w} \rangle$ (quantum mechanics) $v_i w^i$ (Einstein notation) $g_{ij} v^i w^j$ (general relativity). The dot product is also called **scalar product**, or **inner product**.

LENGTH. Using the dot product one can express the length of \vec{v} as $|\vec{v}| = \sqrt{\vec{v} \cdot \vec{v}}$.

CHALLENGE. Express the dot product in terms of the length alone.

SOLUTION: $(\vec{v} + \vec{w}, \vec{v} + \vec{w}) = (\vec{v}, \vec{v}) + (\vec{w}, \vec{w}) + 2(\vec{v}, \vec{w})$ can be solved for (\vec{v}, \vec{w}) .

ANGLE. Because $|\vec{v} - \vec{w}|^2 = (\vec{v} - \vec{w}, \vec{v} - \vec{w}) = |\vec{v}|^2 + |\vec{w}|^2 - 2(\vec{v}, \vec{w})$, which is by the **cos-theorem** equal to $|\vec{v}|^2 + |\vec{w}|^2 - 2|\vec{v}| \cdot |\vec{w}| \cos(\alpha)$, with the angle α between \vec{v}, \vec{w} , we obtain the **dot product formula**

$$\vec{v} \cdot \vec{w} = |\vec{v}| \cdot |\vec{w}| \cos(\alpha)$$

CAUCHY-SCHWARZ INEQUALITY: $|\vec{v} \cdot \vec{w}| \leq |\vec{v}| |\vec{w}|$ follows from that formula because $|\cos(\alpha)| \leq 1$.
TRIANGLE INEQUALITY: $|\vec{u} + \vec{v}| \leq |\vec{u}| + |\vec{v}|$ follows from $|\vec{u} + \vec{v}|^2 = (\vec{u} + \vec{v}) \cdot (\vec{u} + \vec{v}) = \vec{u}^2 + \vec{v}^2 + 2\vec{u} \cdot \vec{v} \leq \vec{u}^2 + \vec{v}^2 + 2|\vec{u}| \cdot |\vec{v}| \leq (\vec{u} + \vec{v})^2$.

FINDING ANGLES BETWEEN VECTORS. Find the angle between the vectors $\langle 1, 4, 3 \rangle$ and $\langle -1, 2, 3 \rangle$.
ANSWER: $\cos(\alpha) = 16/(\sqrt{26}\sqrt{14}) \sim 0.839$. So that $\alpha = \arccos(0.839..) \sim 33^\circ$.

ORTHOGONAL VECTORS. Two vectors are called **orthogonal** (= **perpendicular**) if $\vec{v} \cdot \vec{w} = 0$. The zero vector $\vec{0}$ is orthogonal to any vector. **EXAMPLE:** $\vec{v} = \langle 2, 3 \rangle$ is orthogonal to $\vec{w} = \langle -3, 2 \rangle$.
PYTHAGORAS. If \vec{v} and \vec{w} are orthogonal, then $|\vec{v} - \vec{w}|^2 = |\vec{v}|^2 + |\vec{w}|^2$. **Proof:** $(\vec{v} - \vec{w}) \cdot (\vec{v} - \vec{w}) = \vec{v} \cdot \vec{v} + \vec{w} \cdot \vec{w} + 2\vec{v} \cdot \vec{w} = \vec{v} \cdot \vec{v} + \vec{w} \cdot \vec{w}$.

PROJECTION. The vector $\vec{a} = \text{proj}_{\vec{w}}(\vec{v}) = \vec{w}(\vec{v} \cdot \vec{w} / |\vec{w}|^2)$ is called the **projection** of \vec{v} onto \vec{w} .
 The **scalar projection** is defined as $|\text{comp}_{\vec{w}}(\vec{v})| = |(\vec{v} \cdot \vec{w}) / |\vec{w}||$. Its absolute value is the length of the projection of \vec{v} onto \vec{w} . The vector $\vec{b} = \vec{v} - \vec{a}$ is called the **component** of \vec{v} orthogonal to the \vec{w} -direction.

EXAMPLE. $\vec{v} = \langle 0, -1, 1 \rangle$, $\vec{w} = \langle 1, -1, 0 \rangle$, $\text{proj}_{\vec{w}}(\vec{v}) = \langle 1/2, -1/2, 0 \rangle$, $\text{comp}_{\vec{w}}(\vec{v}) = 1/\sqrt{2}$.