

HEAT AND WAVE EQUATION

Math 21b, Fall 04

FUNCTIONS OF TWO VARIABLES. We consider functions $f(x, t)$ which are for fixed t a piecewise smooth function in x . Analogously as we studied the motion of a **vector** $\vec{v}(t)$, we are now interested in the motion of a **function** $f(x, t)$. While the governing equation for a vector was an ordinary differential equation (ODE), the describing equation will now be a **partial differential equation** (PDE). The function $f(x, t)$ could denote the **temperature of a stick** at a position x at time t or the **displacement of a string** at the position x at time t . The motion of these dynamical systems can be understood using orthonormal Fourier basis $1/\sqrt{2}, \sin(nx), \cos(nx)$ treated in an earlier lecture.

The homework to this lecture is at the end of this 2 page handout.

PARTIAL DERIVATIVES. We write $f_x(x, t)$ and $f_t(x, t)$ for the **partial derivatives** with respect to x or t . The notation $f_{xx}(x, t)$ means that we differentiate twice with respect to x .

Example: for $f(x, t) = \cos(x + 4t^2)$, we have

- $f_x(x, t) = -\sin(x + 4t^2)$
- $f_t(x, t) = -8t \sin(x + 4t^2)$.
- $f_{xx}(x, t) = -\cos(x + 4t^2)$.

One also uses the notation $\frac{\partial f(x, y)}{\partial x}$ for the partial derivative with respect to x . Tired of all the "partial derivative signs", we always write $f_x(x, y)$ or $f_t(x, y)$ in this handout. This is an official abbreviation in the scientific literature.

PARTIAL DIFFERENTIAL EQUATIONS. A partial differential equation is an equation for an unknown function $f(x, t)$ in which different partial derivatives occur.

- $f_t(x, t) + f_x(x, t) = 0$ with $f(x, 0) = \sin(x)$ has a solution $f(x, t) = \sin(x - t)$.
- $f_{tt}(x, t) - f_{xx}(x, t) = 0$ has a solution $f(x, t) = \sin(x - t) + \sin(x + t)$.

THE HEAT EQUATION. The temperature distribution $f(x, t)$ in a metal bar $[0, \pi]$ satisfies the **heat equation**

$$f_t(x, t) = \mu f_{xx}(x, t)$$

This partial differential equation tells that the rate of change of the temperature at x is proportional to the second space derivative of $f(x, t)$ at x . The function $f(x, t)$ is assumed to be zero at both ends of the bar and $f(x) = f(x, t)$ is a given initial temperature distribution. The constant μ depends on the heat conductivity properties of the material. Metals for example conduct heat well and would lead to a large μ .

REWRITING THE PROBLEM. We can write the problem as

$$\frac{d}{dt}f = \mu D^2 f$$

We will solve the problem in the same way as we solved

$$\frac{d}{dt}\vec{x} = A\vec{x}$$

where A is a matrix - by diagonalization.

We use that the Fourier basis is just the diagonalization: $D^2 \cos(nx) = -n^2 \cos(nx)$ and $D^2 \sin(nx) = -n^2 \sin(nx)$ show that $\cos(nx)$ and $\sin(nx)$ are eigenfunctions to D^2 with eigenvalue n^2 . By a symmetry trick, we can focus on sin-series during the entire discussion.

SOLVING THE HEAT EQUATION WITH FOURIER THEORY. The heat equation $f_t(x, t) = \mu f_{xx}(x, t)$ with smooth $f(x, 0) = f(x)$, $f(0, 0) = f(\pi, 0) = 0$ has the solution


$$f(x, t) = \sum_{n=1}^{\infty} b_n \sin(nx) e^{-n^2 \mu t}$$

$$b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin(nx) dx$$


Proof: With the initial condition $f(x) = \sin(nx)$, we have the evolution $f(x, t) = e^{-\mu n^2 t} \sin(nx)$. If $f(x) = \sum_{n=1}^{\infty} b_n \sin(nx)$ then $f(x, t) = \sum_{n=1}^{\infty} b_n e^{-\mu n^2 t} \sin(nx)$.

A SYMMETRY TRICK. Given a function f on $[0, \pi]$ which is zero on both ends. We can extend the function to an odd function on $[-\pi, \pi]$. We do this because of laziness:


The Fourier series of an odd function is a pure sin-series. The Fourier coefficients are $b_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin(nx) dx$ (note the factor 2 and the integration on $[0, \pi]$ as you have proven in a homework.)



Besides the demotivators: "Success is a journey, not a destination. So stop running", there is an other good argument for laziness, namely **efficiency**: "We want to compute as few Fourier coefficients as possible".



EXAMPLE. Assume the initial temperature distribution is a saw-tooth function which has on the interval $[0, \pi/2]$ slope $\pi/2$, on the interval $[\pi/2, \pi]$ the slope $-\pi/2$. Let us compute the sin-Fourier coefficients of this function.




The sin-Fourier coefficients are $b_n = \frac{4}{n^2 \pi} (-1)^{(n-1)/2}$ for odd n and 0 for even n . The solution is

$$f(x, t) = \sum_n b_n e^{-\mu n^2 t} \sin(nx)$$

The exponential term containing the time makes the function $f(x, t)$ converge to 0: The body cools. The higher frequencies are damped faster: "smaller disturbances are smoothed out faster."

VISUALIZATION. We can just plot the graph of the function $f(x, t)$ or plot the temperature distribution for different times t .



THE WAVE EQUATION. The position of a string $f(x, t)$ at time t on $[0, \pi]$ satisfies the **wave equation**

$$f_{tt}(t, x) = c^2 f_{xx}(t, x)$$

where c is a constant. As we will see, c is the **speed** of the waves.

REWRITING THE PROBLEM. We can write the problem as

$$\frac{d^2}{dt^2} f = c^2 D^2 f$$

We will solve the problem in the same way as we solved

$$\frac{d^2}{dx^2} \vec{x} = A \vec{x}$$

If A is diagonal, then every basis vector x satisfies an equation of the form $\frac{d^2}{dt^2} x = -c^2 x$ which has the solution $x(t) = x(0) \cos(ct) + x(t) \sin(ct)/c$.

SOLVING THE WAVE EQUATION WITH FOURIER THEORY. The wave equation $f_{tt} = c^2 f_{xx}$ with $f(x, 0) = f(x)$, $f_t(x, 0) = g(x)$, $f(0, t) = f(\pi, t) = 0$ has the solution


$$f(x, t) = \sum_{n=1}^{\infty} a_n \sin(nx) \cos(nct) + \sum_{n=1}^{\infty} \frac{b_n}{nc} \sin(nx) \sin(nct)$$

$$a_n = \frac{2}{\pi} \int_0^{\pi} f(x) \sin(nx) dx$$


$$b_n = \frac{2}{\pi} \int_0^{\pi} g(x) \sin(nx) dx$$

Proof: With $f(x) = \sin(nx)$, $g(x) = 0$, the solution is $f(x, t) = \cos(ct) \sin(nx)$. With $f(x) = 0$, $g(x) = \sin(nx)$, the solution is $f(x, t) = \frac{1}{c} \sin(ct) \sin(nx)$. For $f(x) = \sum_n a_n \sin(nx)$ and $g(x) = \sum_n b_n \sin(nx)$, we get the formula by summation of those two solutions.


VISUALIZATION. We can just plot the graph of the function $f(x, t)$ or plot the string for different times t .



TO THE DERIVATION OF THE HEAT EQUATION. The temperature $f(x, t)$ is proportional to the kinetic energy at x . Divide the stick into n adjacent cells and assume that in each time step, a fraction of the particles moves randomly either to the right or to the left. If $f_i(t)$ is the **energy** of particles in cell i at time t , then the energy of particles at time $t + 1$ is proportional to $(f_{i-1}(t) - 2f_i(t) + f_{i+1}(t))$. This is a discrete version of the second derivative because $dx^2 f_{xx}(t, x) \sim (f(x + dx, t) - 2f(x, t) + f(x - dx, t))$.



TO THE DERIVATION OF THE WAVE EQUATION. We can model a string by n discrete particles linked by strings. Assume that the particles can move up and down only. If $f_i(t)$ is the **height** of the particles, then the right particle pulls with a force $f_{i+1} - f_i$, the left particle with a force $f_{i-1} - f_i$. Again, $(f_{i-1}(t) - 2f_i(t) + f_{i+1}(t))$ which is a discrete version of the second derivative because $dx^2 f_{xx}(t, x) \sim (f(x + dx, t) - 2f(x, t) + f(x - dx, t))$.



OVERVIEW: The heat and wave equation can be solved like ordinary differential equations:

Ordinary differential equations

$$x_t(t) = Ax(t)$$

$$x_{tt}(t) = Ax(t)$$

Diagonalizing A leads for eigenvectors $\vec{v}$

$$Av = -c^2 v$$

to the differential equations

$$v_t = -c^2 v$$

$$v_{tt} = -c^2 v$$

which are solved by

$$v(t) = e^{-c^2 t} v(0)$$

$$v(t) = v(0) \cos(ct) + v_t(0) \sin(ct)/c$$

Partial differential equations

$$f_t(t, x) = f_{xx}(t, x)$$

$$f_{tt}(t, x) = f_{xx}(t, x)$$

Diagonalizing $T = D^2$ with eigenfunctions $f(x) = \sin(nx)$

$$Tf = -n^2 f$$

leads to the differential equations

$$f_t(x, t) = -n^2 f(x, t)$$

$$f_{tt}(x, t) = -n^2 f(x, t)$$

which are solved by

$$f(x, t) = f(x, 0) e^{-n^2 t}$$

$$f(x, t) = f(x, 0) \cos(nt) + f_t(x, 0) \sin(nt)/n$$

NOTATION:

f function on $[-\pi, \pi]$ smooth or piecewise smooth.
 t time variable
 x space variable
 D the partial differential operator $Df(x) = f'(x) = f_x$
 $d/dx f(x)$.
 T linear transformation, like $Tf = D^2 f = f''$.
 c speed of the wave.

$Tf = \lambda f$ Eigenvalue equation analogously to $Av = \lambda v$.
 f_t partial derivative of $f(x, t)$ with respect to time t .
 f_x partial derivative of $f(x, t)$ with respect to space x .
 f_{xx} second partial derivative of f twice with respect to space x .
 μ heat conductivity
 $f(x) = -f(-x)$ odd function, has sin Fourier series

HOMEWORK. This homework is due until Tuesday morning December 21 in the mailboxes of your CA:

- 1) Solve the heat equation $f_t = \mu f_{xx}$ on $[0, \pi]$ for the initial condition $f(x, 0) = |\sin(3x)|$.
- 2) We want to see in this exercise how to deal with solutions to the heat equation, where the boundary values are not 0.
 - a) Verify that for any constants a, b the function $h(x, t) = (b - a)x/\pi + a$ is a solution to the heat equation.
 - b) Assume we have the problem to describe solutions $f(x, t)$ to the heat equations, where $f(0, t) = a$ and $f(\pi, t) = b$. Show that $f(x, t) - h(x, t)$ is a solution of the heat equation with $f(0, t) = 0$ and $f(\pi, t) = 0$.
 - c) Solve the heat equation with the initial condition $f(x, 0) = f(x) = \sin(3x) + x/\pi$ and satisfying $f(0, t) = 0, f(\pi, t) = 1$ for all times t . This is a situation, when the stick is kept at constant but different temperatures on the both ends.
- 3) A piano string is fixed at the ends $x = 0$ and $x = \pi$ and initially undisturbed. The piano hammer induces an initial velocity $u_t(x, t) = g(x)$ onto the string, where $g(x) = \sin(2x)$ on the interval $[0, \pi/2]$ and $g(x) = 0$ on $[\pi/2, \pi]$. Find the motion of the string.