

- The verification that  $\cos(nx), \sin(nx), 1/\sqrt{2}$  form an orthonormal family is a straightforward computation when using the identities provided. For example,  $\langle \cos(nx), \sin(mx) \rangle = \frac{1}{\pi} \int_{-\pi}^{\pi} \sin(nx) \sin(mx) dx = \frac{1}{2\pi} \int_{-\pi}^{\pi} \cos((n-m)x) - \cos((n+m)x) dx$  which is equal to 1 if  $n = m$  and equal to 0 if  $n \neq m$ . The computations can be abbreviated by noting that integrating an odd  $2\pi$  periodic function over  $[-\pi, \pi]$  is zero.
- To get the Fourier series of the function  $f(x) = |x|$ , note first that this is an **even function** so that it has a cos series. We compute

$$a_0 = \langle f, 1/\sqrt{2} \rangle = \frac{2}{\pi} \int_0^{\pi} x \frac{1}{\sqrt{2}} dx = \frac{\pi\sqrt{2}}{2}.$$

$$a_n = \langle f, \cos(nx) \rangle = \frac{2}{\pi} \int_0^{\pi} x \cos(nx) dx = \frac{2}{\pi} \left[ \frac{\cos(n\pi) - 1}{n^2} \right].$$

The Fourier coefficients of  $f(x) = 5 + |3x|$  are  $a_0 = \frac{3\pi\sqrt{2}}{2} + 5\sqrt{2}$ . and  $a_n = \frac{6}{\pi} \left[ \frac{\cos(n\pi) - 1}{n^2} \right]$ .

- The Fourier series of  $4 \cos^2(3x) + 5 \sin^2(11x) + 90$  is with

$$\cos^2(2x) = \frac{1 + \cos(2x)}{2}$$

$$\sin^2(2x) = \frac{1 - \cos(2x)}{2}$$

given as  $\boxed{4/2 + 4 \cos(6x)/2 - 5 \cos(22x)/2 + 5/2 + 90}$ . All Fourier coefficients are zero except

$$\boxed{a_0 = (189/2) \cdot \sqrt{2} \text{ and } a_6 = 2 \text{ and } a_{22} = -5/2}.$$

- To find the Fourier series of the function  $f(x) = |\sin(x)|$ , we first note that this is an **even function** so that it has a cos-series. If we integrate from 0 to  $\pi$  and multiply the result by 2, we can take the function  $\sin(x)$  instead of  $|\sin(x)|$  so that

$$a_n = \frac{2}{\pi} \int_0^{\pi} \sin(x) \cos(nx) dx.$$

We use one of the trigonometric identities provided in the text to solve this integral.

$$f(x) = \frac{2}{\pi} - \frac{4}{\pi} \left( \frac{\cos(2x)}{2^2 - 1} + \frac{\cos(4x)}{4^2 - 1} + \frac{\cos(6x)}{6^2 - 1} + \dots \right)$$

- The square of the length of the function  $f(x)$  is 1. Parseval's identity shows that

$$1 = a_0^2 + \sum_{n=1}^{\infty} a_n^2 = \left(\sqrt{2} \frac{2}{\pi}\right)^2 + \frac{16}{\pi^2} \left[ \frac{1}{(2^2 - 1)^2} + \frac{1}{(4^2 - 1)^2} + \frac{1}{(6^2 - 1)^2} + \dots \right]$$

so that the sum is  $\boxed{\pi^2/16 - 1/2}$ .

6. To solve the heat equation  $f_t = 8f_{xx}$  on  $[0, \pi]$  with the initial condition  $f(x, 0) = |\sin(3x)|$ , we make a Fourier expansion of  $|\sin(3x)|$ :

$$|\sin(3x)| = \sum_{n=1}^{\infty} b_n \sin(nx)$$

and can immediately write down the solution

$$f(x, t) = \sum_{n=1}^{\infty} b_n e^{-8n^2 t} \sin(nx) .$$

Now to the Fourier series: note that  $\sin(3x)$  is nonnegative on  $[0, \pi/3]$  and  $[2\pi/3, \pi]$  so that it agrees there with the function  $|\sin(3x)|$ . On the interval  $[\pi/3, 2\pi/3]$  however the function  $\sin(3x)$  is negative or zero so that  $-\sin(3x)$  is nonnegative there. We get therefore the Fourier coefficients as

$$b_n = \frac{2}{\pi} \left[ \int_0^{\pi/3} \sin(3x) \sin(nx) dx - \int_{\pi/3}^{2\pi/3} \sin(3x) \sin(nx) dx + \int_{2\pi/3}^{\pi} \sin(3x) \sin(nx) dx \right] .$$

We use the identity

$$2 \sin(nx) \sin(my) = \cos(nx - my) - \cos(nx + my)$$

to solve these integrals:

$$b_n = \frac{2}{\pi} \left[ \left( \frac{-3 \sin(\frac{n\pi}{3})}{n^2 - 9} \right) - \left( \frac{3 \sin(\frac{n\pi}{3})}{n^2 - 9} + \frac{3 \sin(\frac{2n\pi}{3})}{n^2 - 9} \right) + \left( \frac{-3 \sin(\frac{2n\pi}{3})}{n^2 - 9} \right) \right]$$

which is

$$\boxed{\frac{2}{\pi} \left[ \frac{-3 \sin(\frac{n\pi}{3})}{n^2 - 9} \right]} .$$

The case  $n = 3$  might look problematic at first, but the limit still exists and  $b_3 = \pi/6$ .

7. The operator  $D^6 + D^2$  has the eigenvectors  $\sin(nx)$  with eigenvalues  $-n^6 - n^2$ . With initial condition  $f(x) = b_n \sin(nx)$  we have the solution  $b_n \sin(nx) e^{-n^6 - n^2 t}$ . The function  $|\sin(x)|$  is continued as an odd function so that we have to compute the Fourier series of  $\sin(x)$  which is already the Fourier series. The general solution is  $\sum_{n=1}^{\infty} b_n \sin(nx) e^{-n^6 - n^2 t}$ , where  $b_n = 1$  for  $n = 1$  and 0 else.
8. Because the initial condition is zero on the interval  $[\pi/2, \pi]$ , we have to integrate from 0 to  $\pi/2$  only. The Fourier coefficients of the function  $g(x)$  can be computed using one of the trigonometric identities

$$\frac{2}{\pi} \int_0^{\pi/2} \sin(2x) \sin(nx) dx = \frac{-4}{\pi(n^2 - 4)} \sin(n\pi/2) .$$

The Fourier series of the initial position  $f(x) = 0$  of the string is equal to zero by assumption. The solution of the wave equation is

$$f(x, t) = \sum_{n=1}^{\infty} \frac{4(-1)^n}{\pi(n^2 - 4)} \sin(n\pi/2) \sin(nx) \sin(nt) \frac{1}{n} .$$

9. The general solution of the homogeneous equation with the function at rest initially is  $u_h(t, x) = \sum_n b_n \sin(nx) \cos(nt)$ . The particular solution is  $u_p(t, x) = -(1/4) \cos(2t) - (1/25) \cos(5t)$ . Now fix the Fourier coefficients. Note that the particular solution is zero at 0 so that

$$u(t, x) = 11 \sin(5x) \cos(5t) + 23 \sin(7x) \cos(7t) - (1/4) \cos(2t) - (1/25) \cos(5t)$$

is the solution to the differential equation.

10. a) We get

$$b_{nm} = \frac{4}{\pi^2} \int_0^{\pi/2} \int_0^{\pi/2} \sin(nx) \sin(my) \, dy \, dx$$

which is

$$\frac{4}{\pi^2} \left( -\frac{\cos(nx)}{n} \Big|_0^{\pi/2} \right) \left( -\frac{\cos(ny)}{n} \Big|_0^{\pi/2} \right) = \frac{4}{\pi^2} \frac{1}{nm} .$$

- b) The solution to the PDE is

$$\sum_{n,m=1}^{\infty} b_{nm} e^{-(n^2+m^2)t} \sin(nx) \sin(my) .$$