

### True or False

1. T, by Fact 6.1.3 (a diagonal matrix is triangular as well)
2. T, by Fact 6.2.4
3. T; See work on Page 243
4. F; We have  $\det(4A) = 4^4 \det(A)$ , by Fact 6.2.4
5. F; Let  $A = B = I_5$ , for example
6. T; We have  $\det(-A) = (-1)^6 \det(A) = \det(A)$ , by Fact 6.2.4
7. F; If fact,  $\det(A) = 0$ , since  $A$  is noninvertible
8. F; The matrix  $A$  fails to be invertible if  $\det(A) = 0$ , by Fact 6.2.5
9. T, by Fact 6.2.4
10. T, by Fact 6.2.7
11. T, by Example 6 of Section 6.2
12. F, by Fact 6.3.1. The determinant can be  $-1$ .
13. T, by Fact 6.2.7.
14. F: The second and the fourth column are linearly dependent.
15. F; If  $k = -1$  or  $k = -2$ , then the matrix has two equal columns, so that it fails to be invertible.

16. F; The determinant is  $-1$ , since there are three inversions.

17. T; The pattern with the four entries 100 guarantees a nonzero determinant.

18. F; Let  $A = 2I_2$ , for example

19. T; Matrix  $A$  is invertible

20. T; Any nonzero noninvertible matrix  $A$  will do.

21. T, by Fact 6.3.4.

22. T; We have  $\det(A) = \det(\text{rref } A) = 0$ .

23. F; Let  $A = \begin{bmatrix} 3 & 2 \\ 5 & 3 \end{bmatrix}$ , for example

24. F; Let  $A = 2I_2$ , for example

25. T; Let  $A = \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}$ . The columns of  $A$  are orthogonal; now use Fact 6.3.4.

26. F; Let  $A = \begin{bmatrix} 8 & 0 \\ 0 & \frac{1}{2} \end{bmatrix}$  and  $\vec{v} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ , for example.

27. F; In fact,  $\vec{v} \cdot (\vec{u} \times \vec{w}) = -(\det A)$ .

28. T; Let  $A = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$  and  $B = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$ , for example.

29. F; Note that  $\det \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{bmatrix} = 2$ .

30. T, by Fact 6.3.10

31. F; Let  $A = 2I_2$ , for example

32. F; Let  $A = \begin{bmatrix} 0 & -1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{bmatrix}$ , for example.

33. F; Let  $A = I_2$  and  $B = -I_2$ , for example.

34. T; Note that  $\det(B) = -\det(A) < \det(A)$ , so that  $\det(A) > 0$ .

35. T; Let's do Laplace expansion along the first row, for example (see Fact 6.2.10).

Then  $\det(A) = \sum_{j=1}^n (-1)^{1+j} a_{1j} \det(A_{1j}) \neq 0$ . Thus  $\det(A_{1j}) \neq 0$  for at least one  $j$ , so that  $A_{1j}$  is invertible.

36. T; Note that  $\det(A)$  and  $\det(A^{-1})$  are both integers, and  $(\det A)(\det A^{-1}) = 1$ . This leaves only the possibilities  $\det(A) = \det(A^{-1}) = 1$  and  $\det(A) = \det(A^{-1}) = -1$ .

37. T, since  $\text{adj}(A) = (\det A)(A^{-1})$ , by Fact 6.3.10.

38. F; Note that  $\det(A^2) = (\det A)^2$  cannot be negative, but  $\det(-I_3) = -1$ .

39. F; Note that  $\det(S^{-1}AS) = \det(A)$  but  $\det(2A) = 2^3(\det A) = 8(\det A)$ .

40. F; Note that  $\det(S^T AS) = (\det S)^2(\det A)$  and  $\det(-A) = -(\det A)$  have opposite signs.

41. T; The diagonal pattern makes an odd contribution to the determinant, while all the other contributions are even. Thus the determinant is odd; in particular, it is nonzero.

42. F; Let  $A = \begin{bmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 1 & 5 & 2 \end{bmatrix}$ , for example

43. T; Let  $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$ . If  $a \neq 0$ , let  $B = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$ ; if  $b \neq 0$ , let  $B = \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}$ ; if  $c \neq 0$ , let  $B = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ , and if  $d \neq 0$ , let  $B = \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$ .

44. T; Use Gaussian elimination for the first column only to transform  $A$  into a matrix of the form

$$B = \begin{bmatrix} 1 & \pm 1 & \pm 1 & \pm 1 \\ 0 & * & * & * \\ 0 & * & * & * \\ 0 & * & * & * \end{bmatrix}$$

Note that  $\det(B) = \det(A)$  or  $\det(B) = -(\det A)$ . The stars in matrix  $B$  all represent numbers  $(\pm 1) \pm (\pm 1)$ , so that they are 2, 0, or  $-2$ . Thus the determinant of the  $3 \times 3$  matrix  $M$  containing the stars is divisible by 8, since each pattern makes a contribution of 8, 0, or  $-8$ . Now perform Laplace expansion down the first column of  $B$  to see that  $\det(B) = \det(M)$ .