

4. $\det(\lambda I_2 - A) = \det \begin{bmatrix} \lambda & -4 \\ 1 & \lambda - 4 \end{bmatrix} = \lambda(\lambda - 4) + 4 = (\lambda - 2)^2 = 0$ so $\lambda = 2$ with algebraic multiplicity 2.

5. $\det(\lambda I_2 - A) = \det \begin{bmatrix} \lambda - 11 & 15 \\ -6 & \lambda + 7 \end{bmatrix} = \lambda^2 - 4\lambda + 13$ so $\det(\lambda I_2 - A) = 0$ for no real λ .

6. $\det(\lambda I_2 - A) = \det \begin{bmatrix} \lambda - 1 & -2 \\ -3 & \lambda - 4 \end{bmatrix} = \lambda^2 - 5\lambda - 2 = 0$ so $\lambda_{1,2} = \frac{5 \pm \sqrt{33}}{2}$.

7. $\lambda = 1$ with algebraic multiplicity 3, by Fact 7.2.2.

8. $f_A(\lambda) = \lambda^2(\lambda + 3)$ so
 $\lambda_1 = 0$ (Algebraic multiplicity 2)
 $\lambda_2 = -3$.

9. $f_A(\lambda) = (\lambda - 2)^2(\lambda - 1)$ so
 $\lambda_1 = 2$ (Algebraic multiplicity 2)
 $\lambda_2 = 1$.

10. $f_A(\lambda) = (\lambda + 1)^2(\lambda - 1)$ so $\lambda_1 = -1$ (Algebraic multiplicity 2), $\lambda_2 = 1$.

$$15. f_A(\lambda) = \lambda^2 - 2\lambda + (1 - k) = 0 \text{ if } \lambda_{1,2} = \frac{2 \pm \sqrt{4 - 4(1 - k)}}{2} = 1 \pm \sqrt{k}$$


The matrix has 2 distinct real eigenvalues when $k > 0$, no real eigenvalues when $k < 0$.

$$16. f_A(\lambda) = \lambda^2 - (a + c)\lambda + (ac - b^2)$$


24. $\lambda_1 = 0.25, \lambda_2 = 1$

25. $A \begin{bmatrix} c \\ b \end{bmatrix} = \begin{bmatrix} ac + bc \\ bc + bd \end{bmatrix} = \begin{bmatrix} (a + b)c \\ (c + d)b \end{bmatrix} = \begin{bmatrix} c \\ b \end{bmatrix}$ since $a + b = c + d = 1$; therefore, $\begin{bmatrix} c \\ b \end{bmatrix}$ is an eigenvector with eigenvalue $\lambda_1 = 1$.

Also, $A \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} a - c \\ b - d \end{bmatrix} = (a - c) \begin{bmatrix} 1 \\ -1 \end{bmatrix}$ since $a - c = -(b - d)$; therefore, $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$ is an eigenvector with eigenvalue $\lambda_2 = a - c$. Notice that $|a - c| < 1$ so a possible phase portrait is



26. Here $\begin{bmatrix} c \\ b \end{bmatrix} = \begin{bmatrix} 0.25 \\ 0.5 \end{bmatrix}$ with $\lambda_1 = 1$ and $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$ with $\lambda_2 = a - c = 0.25$.



27. a. We know $\vec{v}_1 = \begin{bmatrix} 1 \\ 2 \end{bmatrix}$, $\lambda_1 = 1$ and $\vec{v}_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}$, $\lambda_2 = \frac{1}{4}$. If $\vec{x}_0 = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$ then $\vec{x}_0 = \frac{1}{3}\vec{v}_1 + \frac{2}{3}\vec{v}_2$, so by Fact 7.1.3,


$$x_1(t) = \frac{1}{3} + \frac{2}{3} \left(\frac{1}{4}\right)^t$$

$$x_2(t) = \frac{2}{3} - \frac{2}{3} \left(\frac{1}{4}\right)^t.$$

If $\vec{x}_0 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$ then $\vec{x}_0 = \frac{1}{3}\vec{v}_1 - \frac{1}{3}\vec{v}_2$, so by Fact 7.1.3,

$$x_1(t) = \frac{1}{3} - \frac{1}{3} \left(\frac{1}{4}\right)^t$$

$$x_2(t) = \frac{2}{3} + \frac{1}{3} \left(\frac{1}{4}\right)^t$$



b. A^t approaches $\frac{1}{3} \begin{bmatrix} 1 & 1 \\ 2 & 2 \end{bmatrix}$, as $t \rightarrow \infty$. See part c for a justification.

c. Let us think about the first column of A^t , which is $A^t \bar{e}_1$. We can use Fact 7.1.3 to compute $A^t \bar{e}_1$.

Start by writing $\bar{e}_1 = c_1 \begin{bmatrix} c \\ b \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ -1 \end{bmatrix}$; a straightforward computation shows that $c_1 = \frac{1}{c+b}$ and

$$c_2 = \frac{b}{c+b}.$$

Now $A^t \bar{e}_1 = \frac{1}{c+b} \begin{bmatrix} c \\ b \end{bmatrix} + \frac{b}{c+b} (\lambda_2)^t \begin{bmatrix} 1 \\ -1 \end{bmatrix}$, where $\lambda_2 = a - c$.

Since $|\lambda_2| < 1$, the second summand goes to zero, so that $\lim_{t \rightarrow \infty} (A^t \bar{e}_1) = \frac{1}{c+b} \begin{bmatrix} c \\ b \end{bmatrix}$.

Likewise, $\lim_{t \rightarrow \infty} (A^t \bar{e}_2) = \frac{1}{c+b} \begin{bmatrix} c \\ b \end{bmatrix}$, so that $\lim_{t \rightarrow \infty} A^t = \frac{1}{c+b} \begin{bmatrix} c & c \\ b & b \end{bmatrix}$.

28. a. $w(t+1) = 0.8w(t) + 0.1m(t)$
 $m(t+1) = 0.2w(t) + 0.9m(t)$

so $A = \begin{bmatrix} 0.8 & 0.1 \\ 0.2 & 0.9 \end{bmatrix}$ which is a regular transition matrix since its columns sum to 1 and its entries are positive.

b. The eigenvectors of A are $\begin{bmatrix} 0.1 \\ 0.2 \end{bmatrix}$ or $\begin{bmatrix} 1 \\ 2 \end{bmatrix}$ with $\lambda_1 = 1$, and $\begin{bmatrix} 1 \\ -1 \end{bmatrix}$ with $\lambda_2 = 0.7$.

$$\bar{x}_0 = \begin{bmatrix} 1200 \\ 0 \end{bmatrix} = 400 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 800 \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ so } \bar{x}(t) = 400 \begin{bmatrix} 1 \\ 2 \end{bmatrix} + 800(0.7)^t \begin{bmatrix} 1 \\ -1 \end{bmatrix} \text{ or}$$

$$w(t) = 400 + 800(0.7)^t$$

$$m(t) = 800 - 800(0.7)^t.$$

c. As $t \rightarrow \infty$, $w(t) \rightarrow 400$ so Wipfs won't have to close the store.