

True or False

1. T; The matrix is $\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$.
2. F; The columns of a rotation matrix are unit vectors; see Fact 2.2.2.
3. T, by Fact 2.3.3.
4. T; Let $A = B$ in Fact 2.4.8.
5. F, by Fact 2.4.3.
6. T, by Fact 2.4.9.
7. F; Matrix AB will be 3×5 , by Definition 2.4.1b.
8. F; Note that $T \begin{bmatrix} 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$. A linear transformation transforms $\vec{0}$ into $\vec{0}$.
9. T, by Fact 2.2.3.
10. T, by Fact 2.4.5.
11. F, by Fact 2.3.6. Note that the determinant is 0.
12. T, by Fact 2.3.3.
13. T; The shear matrix $A = \begin{bmatrix} 1 & \frac{1}{2} \\ 0 & 1 \end{bmatrix}$ works.
14. T; Simplify to see that $T \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4y \\ -12x \end{bmatrix} = \begin{bmatrix} 0 & 4 \\ -12 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$.
15. T; The equation $\det(A) = k^2 - 6k + 10 = 0$ has no real solution.
16. T; The matrix fails to be invertible for $k = 5$ and $k = -1$, since the determinant is 0 for these values.

17. F; Note that $\det(A) = (k - 2)^2 + 9$ is always positive, so that A is invertible for all values of k .
18. T; Note that the columns are unit vectors, since $(-0.6)^2 + (\pm 0.8)^2 = 1$. The matrix has the form presented in Fact 2.2.2.
19. F; Consider $A = I_2$ (or any other invertible 2×2 matrix).
20. T; Note that $A = \begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}^{-1} \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 5 & 6 \\ 7 & 8 \end{bmatrix}^{-1}$ is the unique solution.
21. F; For any 2×2 matrix A , the two columns of $A \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ will be identical.
22. T; One solution is $A = \begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}$.
23. T; Note that $\begin{bmatrix} -1 & 2 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$ and $\begin{bmatrix} -1 & 2 \\ -2 & 3 \end{bmatrix} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 1 \end{bmatrix}$, so that it's a shear parallel to the line spanned by vector $\begin{bmatrix} 1 \\ 1 \end{bmatrix}$.
24. T; Just multiply it out.
25. T; The product is $\det(A)I_2$.
26. T; Writing an upper triangular matrix $A = \begin{bmatrix} a & b \\ 0 & c \end{bmatrix}$ and solving the equation $A^2 = \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix}$ we find that $A = \begin{bmatrix} 0 & b \\ 0 & 0 \end{bmatrix}$, where b is any nonzero constant.
27. T; Note that the matrix $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$ represents a rotation through $\pi/2$. Thus $n = 4$ (or any multiple of 4) works.
28. F; If a matrix A is invertible, then so is A^{-1} . But $\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$ fails to be invertible.
29. F; If matrix A has two identical rows, then so does AB , for any matrix B . Thus AB cannot be I_n , so that A fails to be invertible.
30. T, by Fact 2.3.4. Note that $A^{-1} = A$ in this case.
31. F; Consider the matrix A that represents a rotation through the angle $2\pi/17$.
32. F; Consider the reflection matrix $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$.
33. T; We have $(5A)^{-1} = \frac{1}{5}A^{-1}$.

34. T; The equation $A\vec{e}_i = B\vec{e}_i$ means that the i th columns of A and B are identical. This observation applies to all the columns.
35. T; Note that $A^2B = AAB = ABA = BAA = BA^2$.
36. T; Multiply both sides of the equation $A^2 = A$ with A^{-1} .
37. F; Consider $A = I_2$ and $B = -I_2$.
38. T; Since $A\vec{x}$ is on the line onto which we project, the vector $A\vec{x}$ remains unchanged when we project again: $A(A\vec{x}) = A\vec{x}$, or $A^2\vec{x} = A\vec{x}$, for all \vec{x} . Thus $A^2 = A$.
39. F; Consider matrix $\begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix}$, for example.
40. T; Apply Fact 2.4.9 to the equation $(A^2)^{-1}AA = I_n$, with $B = (A^2)^{-1}A$.
41. T; If you reflect twice in a row (across the same line), you will get the original vector back: $A(A\vec{x}) = \vec{x}$, or, $A^2\vec{x} = \vec{x} = I_2\vec{x}$. Thus $A^2 = I_2$ and $A^{-1} = A$.
42. F; Let $A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$, $\vec{v} = \begin{bmatrix} 1 \\ 0 \end{bmatrix}$, $\vec{w} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$, for example.
43. T; Let $A = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix}$, $B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$, for example.
44. F; By Fact 1.3.3, there is a nonzero vector \vec{x} such that $B\vec{x} = \vec{0}$, so that $AB\vec{x} = \vec{0}$ as well. But $I_3\vec{x} = \vec{x} \neq \vec{0}$, so that $AB \neq I_3$.
45. T; We can rewrite the given equation as $A^2 + 3A = -4I_3$ and $-\frac{1}{4}(A + 3I_3)A = I_3$. By Fact 2.4.9, matrix A is invertible, with $A^{-1} = -\frac{1}{4}(A + 3I_3)$.
46. T; Note that $(I_n + A)(I_n - A) = I_n^2 - A^2 = I_n$, so that $(I_n + A)^{-1} = I_n - A$.
47. T; If we shear parallel to line L , then the vector $A\vec{x} - \vec{x}$ is parallel to L , for all \vec{x} , by part b of Definition 2.2.4. By part a, we have $A(A\vec{x} - \vec{x}) = A\vec{x} - \vec{x}$, or $(A^2 - 2A + I_2)\vec{x} = \vec{0}$. Since this holds for all vectors \vec{x} in \mathbb{R}^2 , the equation $A^2 - 2A + I_2 = 0$ must hold.
48. F; Consider $T(\vec{x}) = 2\vec{x}$, $\vec{v} = \vec{e}_1$, and $\vec{w} = \vec{e}_2$.
49. F; Since there are only eight entries that are not 1, there will be at least two rows that contain only ones. Having two identical rows, the matrix will be non-invertible.
50. F; Let $A = B = \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix}$, for example.