

Math 23a: Theoretical Linear Algebra
and Multivariable Calculus I

FINAL EXAM

January 24, 2006

Your name: _____

Problem	Points	Score
1	15	
2	10	
3	10	
4	10	
5	15	
6	20	
7	20	
Total	100	

In the following problems you can use any of the results we have proved in class, if you state them clearly before using them.

Please show all your work on this exam paper. Unless otherwise stated, you must show your work and clearly indicate your line of reasoning in order to get full credit. You can write on the back of the pages if you need extra paper.

Problem 1 [15 points]

Decide whether the following statements are True or False. (Note: There is no need to justify your answers, just circle T or F. You get +3 for every correct answer and -1 for every wrong answer.)

T or F: The set of rational numbers forms an ordered field.

T or F: If A, B are $n \times n$ matrices, $A \neq 0$, and $AB = 0$, then $B = 0$.

T or F: Every orthonormal list of vectors in a finite dimensional Euclidean space V can be extended to an orthonormal basis of V .

T or F: Let $T : V \rightarrow V$ be an operator on a finite dimensional Euclidean space V . Eigenvectors of T corresponding to distinct eigenvalues are orthogonal.

T or F: If A is a unitary matrix, then $|\det A| = 1$.

Problem 2 [10 points]

Find a basis for the solution space $U \subset \mathbb{R}^4$ of the following homogeneous system of linear equations:

$$\begin{cases} x_1 - x_2 + 2x_3 - x_4 &= 0 \\ x_2 + x_3 - 3x_4 &= 0 \\ 2x_1 - x_2 + 5x_3 - 5x_4 &= 0 \end{cases}$$

Answer:

Basis U :	
-------------	--

Problem 3 [10 points]

Say which of the following functions is a linear transformation and, in case it is, write down the corresponding matrix A , the kernel $\text{Ker}\phi$ and the image $\text{Im}\phi$.

$$\begin{aligned} \text{(a)} \quad \phi: \mathbb{R}^3 &\rightarrow \mathbb{R}^2, & \phi\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) &= \begin{bmatrix} xy + z \\ yz - x \end{bmatrix} \\ \text{(b)} \quad \phi: \mathbb{R}^2 &\rightarrow \mathbb{R}^2, & \phi\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) &= \begin{bmatrix} x + 2y \\ -4y - 2x \end{bmatrix} \\ \text{(c)} \quad \phi: \mathbb{R}^3 &\rightarrow \mathbb{R}^2, & \phi\left(\begin{bmatrix} x \\ y \\ z \end{bmatrix}\right) &= \begin{bmatrix} x + y + z \\ x - y - 1 \\ z + x \end{bmatrix} \end{aligned}$$

Answers:

(a)	Y N	A=	$\text{Ker}\phi=$	$\text{Im}\phi=$
(b)	Y N	A=	$\text{Ker}\phi=$	$\text{Im}\phi=$
(c)	Y N	A=	$\text{Ker}\phi=$	$\text{Im}\phi=$

Problem 4 [10 points]

Compute the determinant of the following matrices:

$$(a) \ A = \begin{bmatrix} 0 & 0 & 5 & 1 \\ -1 & -2 & 6 & 2 \\ 0 & 0 & 7 & 3 \\ 1 & 2 & 3 & 4 \end{bmatrix}$$

$$(b) \ B = \begin{bmatrix} 1001 & 1002 & 1003 & 1004 \\ 1002 & 1003 & 1001 & 1002 \\ 1001 & 1001 & 1001 & 999 \\ 1001 & 1001 & 998 & 996 \end{bmatrix}$$

Answers:

$\det A =$	
$\det B =$	

Problem 5 [15 points]

Suppose V is a vector space over $\mathbb{C}$ of dimension n and $T : V \rightarrow V$ is a linear operator on V . Suppose that T has only one eigenvalue λ . Prove that $(T - \lambda \mathbb{I})^n = 0$.

Problem 6 [20 points]

Let A be an $n \times n$ Hermitian matrix with complex coefficients, and denote by a_{ij} the entry of A in row i and column j . Let $\lambda_1, \dots, \lambda_n$ be the eigenvalues of A , counted with multiplicities.

- (a) State the Spectral Theorem for Hermitian matrices.
- (b) Let λ be the lowest eigenvalue: $\lambda = \min\{\lambda_1, \dots, \lambda_n\}$. Prove that $a_{ii} \geq \lambda$ for every $i = 1, \dots, n$.

Problem 7 [20 points]

Consider the field of two elements $\mathbb{F} = \mathbb{Z}/2\mathbb{Z}$. Let S be any set, and let V_S be the set consisting of all subsets of S . We define on V_S two operations: addition

$$A + B = (A - B) \cup (B - A)$$

(recall that $A - B$, the difference of subsets, denotes the set of all elements of A which are not in B), and scalar multiplication (by elements of $\mathbb{Z}/2\mathbb{Z}$):

$$0 \cdot A = \emptyset, \quad 1 \cdot A = A.$$

- (a) I claim that V_S , with the above operations, is a vector space over $\mathbb{Z}/2\mathbb{Z}$. You do not need to check all the axioms. Just state and prove the following vector space axioms:

1. associativity of addition (what is $A + B + C$?),
2. existence of additive identity,
3. existence of additive inverse

- (b) Given a subset $A \subset S$, consider the following functions $I_A, U_A : V_S \rightarrow V_S$,

$$I_A(B) = A \cap B, \quad U_A(B) = A \cup B.$$

Say whether the above functions are linear transformations on V_S and, if they are, determine their kernel and their image.

(page intentionally left blank)

.