

MATH 23a, FALL 2001
THEORETICAL LINEAR ALGEBRA
AND MULTIVARIABLE CALCULUS
(Final Version) Homework Assignment # 9
Due: *Friday*, November 30, 2001

Please turn in five separate sets labelled A through E.

1. Read Sections 1.8 and 2.1–2.2 from Edwards, and re-read the Appendix.
2. (A) Finish the following proof from class (11/19) using an explicit δ - ε argument:

Let V be a real vector space with a convergent sequence $\{\mathbf{v}_n\}$ such that $\lim_{n \rightarrow \infty} \mathbf{v}_n = \mathbf{v}$, and let $\{c_n\}$ be a convergent sequence of real numbers with $\lim_{n \rightarrow \infty} c_n = c$. Show that $\{c_n \mathbf{v}_n\}$ converges to $c\mathbf{v}$.

3. (A) Suppose $f : \mathbb{R}^n \rightarrow \mathbb{R}$ and $g : \mathbb{R}^n \rightarrow \mathbb{R}$ are continuous at \mathbf{a} , and suppose $g(\mathbf{a}) \neq 0$. Show that $\frac{f}{g}$ is continuous at \mathbf{a} .
4. (B) A set S is called a **metric space** if there exists a function $d : S \times S \rightarrow \mathbb{R}$ called the *distance* such that:
 - i. $d(x, y) = d(y, x), \forall x, y \in S$
 - ii. $d(x, y) \leq d(x, z) + d(z, y), \forall x, y, z \in S$
 - iii. $d(x, x) = 0, \forall x \in S$
 - iv. $d(x, y) > 0, \forall x, y \in S$ with $x \neq y$

- (a) Let S be any set, and define $d(x, x) = 0, \forall x \in S$ and $d(x, y) = 1$ if $x \neq y$. Show that this is a metric space.
- (b) Suppose S is a metric space with distance d . Show that S is also a metric space with new distance given by:

$$d'(x, y) = \frac{d(x, y)}{1 + d(x, y)}$$

5. (B) Consider $S = \mathbb{R}^2$, with $\mathbf{v} = (a, b)$ and $\mathbf{w} = (c, e)$. We define the *Memphis metric* by $d(\mathbf{v}, \mathbf{v}) = 0$ for any \mathbf{v} , and for $\mathbf{v} \neq \mathbf{w}$,

$$d(\mathbf{v}, \mathbf{w}) = \sqrt{a^2 + b^2} + \sqrt{c^2 + e^2}$$

- (a) Show that S is a metric space. (See problem #4.)
- (b) Find $B_\varepsilon(\mathbf{0})$.
- (c) For $\mathbf{v} \neq \mathbf{0}$, find $B_\varepsilon(\mathbf{v})$, for various $\varepsilon > 0$.

6. (C) Define $f : [0, 1] \rightarrow \mathbb{R}$ as follows:

$$f(x) = \begin{cases} 0 & , \text{ if } x \notin \mathbb{Q} \\ \frac{1}{q} & , \text{ if } x \in \mathbb{Q} \text{ and } x = \frac{p}{q} \text{ in lowest terms} \end{cases}$$

- (a) Graph f .
- (b) Show that f is not continuous at any rational x .
- (c) Show that f is continuous at any irrational x .
7. (D) Given a set S in a normed vector space, we define a point $x \in S$ to be an **interior point** if $\exists \varepsilon > 0$ such that $B_\varepsilon(x) \subset S$. We define the **interior** of S to be the set of interior points, and we denote it by S° . Show that the interior of any set is open.
8. (D) Let $V = \mathbb{R}^2$, and consider the following subsets:
 $A = \mathbb{Q} \times \mathbb{Q} = \{(x, y) \in \mathbb{R}^2 \mid x, y \in \mathbb{Q}\}$ and $B = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$
 For the following, recall that \bar{S} , S° , and S^c denote the closure, interior (see #7), and complement, respectively, of S .
- (a) Find A° , $(A^c)^\circ$, \bar{A} , $\overline{A^c}$, and $\bar{A} \cap \overline{A^c}$.
- (b) Find B° , $(B^c)^\circ$, \bar{B} , $\overline{B^c}$, and $\bar{B} \cap \overline{B^c}$.
- (c) Find $\overline{A \cap B}$ and $(A \cap B)^\circ$.
9. (C) Let $S = \{(x, \sin(\frac{1}{x})) \mid x > 0\} \subset \mathbb{R}^2$. Find \bar{S} .
10. (E) A subset S in a metric space or a normed vector space is called **discrete** if, for every $x \in S$, there is some $\varepsilon > 0$ such that $B_\varepsilon(x) \cap S = \{x\}$, that is, the only intersection between the ball and the set is the point itself.
- (a) Show that every $f : S \rightarrow \mathbb{R}$ is continuous if S is discrete.
- (b) Show that every closed, bounded, and discrete set is finite, and give examples why each of these three conditions is necessary.
- (c) Show that $\mathbb{Z} \subset \mathbb{R}$ is discrete.
- (d) Show that $\mathbb{Z} \subset \mathbb{Q}_p$ is not discrete.
11. (E) Let $\{S_n\}$ be a collection of open sets in a normed vector space, and let $\{T_n\}$ be a collection of closed sets. Show that:
- (a) $S_1 \cap S_2$ is open.
- (b) $\bigcup S_n$ is open.
- (c) $T_1 \cup T_2$ is closed.
- (d) $\bigcap T_n$ is closed.