

MATH 23b, SPRING 2004  
THEORETICAL LINEAR ALGEBRA  
AND MULTIVARIABLE CALCULUS  
(Final Version) Homework Assignment # 1  
Due: February 13, 2004

Please turn in four separate sets labelled A through D. Starred problems are considered “moral homework” and do not need to be turned in, though I will consider them to be part of the course content.

1. Read Sections 1.7–1.8 from Edwards, and re-read the Appendix.

2. (A) Prove the following Lemma from class (2/6):

**Lemma.** Let  $\{c_n\}$  be a convergent sequence of real numbers with  $\lim_{n \rightarrow \infty} c_n = c$ . Show that  $\{c_n\}$  is bounded, that is, that there exists an  $M > 0$  such that  $|c_n| < M, \forall n \in \mathbb{N}$ .

3. (B) Metric spaces.

(a) Let  $S$  be any set, and define  $d(x, x) = 0, \forall x \in S$  and  $d(x, y) = 1$  if  $x \neq y$ . Show that  $(S, d)$  is a metric space. For the record, this particular metric is known as the *discrete metric*.

(b) Suppose  $S$  is a metric space with distance function  $d$ . Show that  $S$  is also a metric space with new distance function given by:

$$d'(x, y) = \frac{d(x, y)}{1 + d(x, y)}$$

4. (B) Consider  $S = \mathbb{R}^2$ , with  $\mathbf{v} = (a, b)$  and  $\mathbf{w} = (c, e)$ . We define the *Memphis metric* by  $d(\mathbf{v}, \mathbf{v}) = 0$  for any  $\mathbf{v}$ , and for  $\mathbf{v} \neq \mathbf{w}$  by:

$$d(\mathbf{v}, \mathbf{w}) = \sqrt{a^2 + b^2} + \sqrt{c^2 + e^2}$$

(a) Show that  $(S, d)$  is a metric space.

(b) Find  $B_\varepsilon(\mathbf{0})$ .

(c) For  $\mathbf{v} \neq \mathbf{0}$ , find  $B_\varepsilon(\mathbf{v})$ , for various  $\varepsilon > 0$ .

5. (C) Define  $f : [0, 1] \rightarrow \mathbb{R}$  as follows:

$$f(x) = \begin{cases} 0 & , \text{ if } x \notin \mathbb{Q} \\ \frac{1}{q} & , \text{ if } x \in \mathbb{Q} \text{ and } x = \frac{p}{q} \text{ in lowest terms} \end{cases}$$

(a) Graph  $f$ .

(Hint: Use a large scale, and plot points in a “natural” order.)

- (b) Show that  $f$  is not continuous at any rational  $x$ .
  - (c) Show that  $f$  is continuous at any irrational  $x$ .
6. (A) Suppose  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $g : \mathbb{R}^n \rightarrow \mathbb{R}$  are continuous at  $\mathbf{a}$ , and suppose  $g(\mathbf{a}) \neq 0$ . Show that  $\frac{f}{g}$  is continuous at  $\mathbf{a}$ .
7. (D) Given a set  $S$  in a normed vector space, we define a point  $x \in S$  to be an **interior point** if  $\exists \varepsilon > 0$  such that  $B_\varepsilon(x) \subset S$ . We define the **interior** of  $S$  to be the set of interior points, and we denote it by  $S^\circ$ . Show that the interior of any set is open.
8. (deferred) Let  $V = \mathbb{R}^2$ , and consider the following subsets:

$$A = \mathbb{Q} \times \mathbb{Q} = \{(x, y) \in \mathbb{R}^2 \mid x, y \in \mathbb{Q}\}$$

$$B = \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 < 1\}$$

For the following, recall that  $\bar{S}$ ,  $S^\circ$ , and  $S^c$  denote the closure, interior (see #7), and complement, respectively, of  $S$ .

- (a) Find  $A^\circ$ ,  $(A^c)^\circ$ ,  $\bar{A}$ ,  $\overline{A^c}$ , and  $\bar{A} \cap \overline{A^c}$ .
  - (b) Find  $B^\circ$ ,  $(B^c)^\circ$ ,  $\bar{B}$ ,  $\overline{B^c}$ , and  $\bar{B} \cap \overline{B^c}$ .
  - (c) Find  $\overline{A \cap B}$  and  $(A \cap B)^\circ$ .
9. (\*) Let  $S = \{(x, \sin(\frac{1}{x})) \mid x > 0\} \subset \mathbb{R}^2$ . Find  $\bar{S}$ .
10. (deferred) A subset  $S$  in a metric space or a normed vector space is called **discrete** if, for every  $x \in S$ , there is some  $\varepsilon > 0$  such that  $B_\varepsilon(x) \cap S = \{x\}$ , that is, the only intersection between the ball and the set is the point itself.
- (a) Show that every  $f : S \rightarrow \mathbb{R}$  is continuous if  $S$  is discrete.
  - (b) Show that every closed, bounded, and discrete set is finite, and give examples why each of these three conditions is necessary.
  - (c) Show that  $\mathbb{Z} \subset \mathbb{R}$  is discrete.
11. (D) Let  $\{S_n\}$  be a collection of open sets in a normed vector space, and let  $\{T_n\}$  be a collection of closed sets. Show that:
- (a)  $S_1 \cap S_2$  is open.
  - (b)  $\bigcup S_n$  is open.
  - (c)  $T_1 \cup T_2$  is closed.
  - (d)  $\bigcap T_n$  is closed.