

MATH 23a, FALL 2003
THEORETICAL LINEAR ALGEBRA
AND MULTIVARIABLE CALCULUS
(Final Version) Homework Assignment # 7
Due: November 21, 2003

1. Read Chapter 4 (especially 4.1–4.3) and 6.3–6.4 of Schneider and Barker.
2. (*) Read Chapter 2 of Schneider and Barker.
3. (A) Let $V = (\mathbb{Z}/7\mathbb{Z})^3$, and consider the linear map $L : V \longrightarrow V$ given by $L(x, y, z) = (x + y + z, 2y + 3z, 4z)$.
 - (a) Write the matrix A for L with respect to the standard basis.
 - (b) Find the eigenvalues of L , and find an eigenbasis for V . (Hint: Look for likely choices of eigenvalues—I claim that two of them are easy, and the third follows a pattern.)
 - (c) Write the matrix B for L with respect to the eigenbasis.
 - (d) Find the change of basis matrix $S : V \rightarrow V$ that takes the standard basis to the eigenbasis.
 - (e) Show directly (that is, using a computation involving S) that A and B are similar.
4. (B) Let $V = \mathbb{R}^n$, and let $\mathbf{u}, \mathbf{v} \in V$. If $A : V \longrightarrow V$ is (the matrix for) a linear transformation, then define the following bilinear form:

$$f_A(\mathbf{u}, \mathbf{v}) = \mathbf{u}^t A \mathbf{v}$$

- (a) Show that f_A is indeed a bilinear form.
 - (b) Give a necessary and sufficient condition on the matrix A that makes f_A alternating.
- (Hint #1: You might consider the $n = 2$ case first to get a feel for this bilinear form. Hint #2: Your answers just might involve the transpose!)
5. (C) Show that not every skew-symmetric multilinear form $f : V^n \longrightarrow F$ is alternating by constructing an example. (Note that the only cases where this can happen are over fields F wherein $1 + 1 = 0$. Follow Halmos' reasoning in the proof that alternating implies skew-symmetric, Section 30, Theorem 1.)
 6. (D) Let $\dim(V) = n$, and let $f : V^k \longrightarrow F$ be an alternating k -linear form with $k < n$. Show by example that it is possible to have a set of k linearly independent vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ in V such that $f(\mathbf{v}_1, \dots, \mathbf{v}_k) = 0$. (Make sure that $k \geq 2$ so that f can be alternating!)

7. (E) Let V be a vector space over F . Consider the set of k -linear forms $f : V^k \longrightarrow F$. For any two such forms f_1 and f_2 and any scalar $c \in F$, we define:

$$\begin{aligned}(f_1 + f_2)(\mathbf{v}_1, \dots, \mathbf{v}_k) &= f_1(\mathbf{v}_1, \dots, \mathbf{v}_k) + f_2(\mathbf{v}_1, \dots, \mathbf{v}_k), & \forall \mathbf{v}_1, \dots, \mathbf{v}_k \in V \\ (cf_1)(\mathbf{v}_1, \dots, \mathbf{v}_k) &= c \cdot f_1(\mathbf{v}_1, \dots, \mathbf{v}_k), & \forall \mathbf{v}_1, \dots, \mathbf{v}_k \in V\end{aligned}$$

- (a) (*) Convince yourself that the collection of k -linear forms on V form a vector space over F with addition and scalar multiplication defined as above.
- (b) Let $V = \mathbb{R}^2$. Show that the form $f : V^2 \longrightarrow \mathbb{R}$ defined by $f((a, b), (c, d)) = ad - bc$ is bilinear and alternating.
- (c) Now let $V = \mathbb{R}^3$. Construct two *linearly independent* alternating bilinear forms $f : V^2 \longrightarrow \mathbb{R}$.
- (d) Determine the dimensions of the spaces of alternating bilinear forms on $V = \mathbb{R}^2$ and $V = \mathbb{R}^3$.

8. (deferred)

Let $A : V \longrightarrow V$ be a linear transformation on a finite-dimensional vector space, and by slight abuse of notation, let A also be the matrix for this transformation with respect to a fixed basis. Using the following method, we determine the eigenvalues of A :

$$\begin{aligned}\lambda \text{ is an eigenvalue for } A &\iff V_\lambda = \text{Ker}(A - \lambda I) \text{ is non-trivial} \\ &\iff A - \lambda I \text{ is not invertible} \\ &\iff \det(A - \lambda I) = 0\end{aligned}$$

Thus we are inspired to make the following definition:

$p_A(\lambda) = \det(A - \lambda I)$ is called the **characteristic polynomial** of A

The eigenvalues of A will be the roots of the characteristic polynomial.

- (a) Prove that no scalar $\lambda_0 \in F$ is an eigenvalue for A unless it is a root of $p_A(\lambda)$.
- (b) If $p_A(\lambda) = (\lambda - \lambda_0)^k \cdot q(\lambda)$ with $q(\lambda_0) \neq 0$, then we say that the eigenvalue λ_0 has *algebraic multiplicity* equal to k . (That is, λ_0 is a root of $p_A(\lambda)$ of order k .) Show that the geometric multiplicity (which, by definition, is the dimension of the corresponding eigenspace) of an eigenvalue is less than or equal to its algebraic multiplicity.
- (c) Use this method to find all eigenvalues of the real matrix

$$A = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 0 & -1 & 0 \end{bmatrix}$$

Is the matrix A diagonalizable? Explain.