

Math 55 Problem Set 3

Neil Herriot with additions by Andrei Jorza

1. (i)  $d_1(f, f) = \int_0^1 |f(x) - f(x)| dx = \int_0^1 0 dx = 0$ . And, if  $f \neq g$ , then  $F(x) = |f(x) - g(x)|$  is not identically zero. Hence  $\exists x_0$  so  $F(x_0) = 2\epsilon > 0$ . And by continuity,  $\exists \delta$  such that  $\forall x, x_0 - \delta < x < x_0 + \delta, F(x) > \epsilon$ . So,  $d_1(f, g) = \int_0^1 F(x) dx = \int_0^{x_0 - \delta} F(x) dx + \int_{x_0 - \delta}^{x_0 + \delta} F(x) dx + \int_{x_0 + \delta}^1 F(x) dx \geq \int_{x_0 - \delta}^{x_0 + \delta} dx = 2\epsilon\delta > 0$   
 $0 = d_1(f, g) = \int_0^1 |f(x) - g(x)| dx$ , implies  $|f(x) - g(x)| = 0$  for all  $x$ , and thus  $f = g$ .  
(ii)  $d_1(f, g)$  is symmetric in definition, as  $|f - g| = |g - f|$ , and thus equals  $d_1(g, f)$ .  
(iii)  $d_1(f, h) + d_1(h, g) = \int_0^1 |f(x) - h(x)| + |h(x) - g(x)| dx \geq \int_0^1 |f(x) - g(x)| dx = d_1(f, g)$ .
2.  $\{f_n\} : \forall \epsilon > 0, \exists N (> 1/\epsilon)$  so that  $\forall n > N, x \in \mathbb{R}$  we have  $0 < \frac{n}{x^2 + n^2} \leq \frac{n}{n^2} < \epsilon$ . Thus,  $\{f_n\}$  converges uniformly (and thus also point-wise) to 0 on all of  $\mathbb{R}$ .  
 $\{g_n\} : \text{For any given } x, \text{ we can pick a sufficiently large } N (> \sqrt{\frac{x^2}{\epsilon}})$  so that,  $n > N$  implies  $g_n$  is  $\epsilon$  close to, 1. Thus,  $\{g_n\}$  converges point-wise. But, for any  $n$ , if we pick  $x = n$  to get  $g_n = \frac{1}{2}$ . This means that  $\{g_n\}$  does not converge uniformly to 1.
3. Let  $U = \{x : d(x, A) < d(x, B)\}$  and  $V$  defined analogously, with  $d$  the distance function defined in problem set 2 problem 1. Clearly,  $U$  and  $V$  are disjoint and contain  $A$  and  $B$  respectively. Now, it remains to show that they are open. Let  $x \in U, \epsilon = \frac{1}{3}(d(x, B) - d(x, A))$ , and  $y \in B_\epsilon(x)$ . Then,  $d(y, B) - d(y, A) > d(x, B) - d(x, A) - 2\epsilon > 0$ , and we have  $y \in U$  and  $U$  open. So  $U$  and  $V$  are as desired.
4. I claim that for  $U, V$  open,  $U \subset \bar{U} \subset V$ , there exists  $W$  open so that  $\bar{U} \subset W \subset \bar{W} \subset V$ . To see this, consider  $\bar{U}$  and  $X - V$ , disjoint closed sets. Then there exists  $W \supset \bar{U}, W' \supset X - V$  open and disjoint. But then  $\bar{U} \subset W \subset X - W' \subset X - V$ . Since  $X - W'$  is closed and contains  $W$ , it also contains  $\bar{W}$ . And we have  $U \subset \bar{U} \subset W \subset \bar{W} \subset V$ , as desired.

Now, letting  $S_1 \subset X - B$ , and  $S_0 \supset A$  as produced by  $X - B$  and  $\text{Int}A$  with the above lemma applied twice. I inductively create open  $S_{k/2^i}$  one level of “i” at a time. At each point if  $q < r$ , then  $\bar{S}_q \subset S_r$ .  $S_{k/2^i}$  ( $k$  odd), is generated by the above lemma between  $S_{(k-1)/2^i}$  and  $S_{(k+1)/2^i}$ . So of course, all the sets contain the closure of  $S_0$  and are contained in  $S_1$ .

I define  $f(x) = \inf(\{1\} \cup \{r : x \in S_r\})$ . This set is bounded below and non empty, so the inf exists. It is clear that  $f(x) = 0$  if  $x \in A$ ,  $f(x) = 1$  if  $x \in B$ , and has range  $[0, 1]$ . It now remains to show that  $f$  is continuous. I first show  $f(x) = \sup(\{0\} \cup \{r : x \in X - \bar{S}_r\})$ . In doing this, we only need to consider  $r$  in the sets that are terminating binary fractions, as for no other  $r$  is  $S_r$  defined and thus is it possible for  $r$  to satisfy the condition to be in the sets.

If,  $r > f(x)$  then  $\exists r_0, r > r_0 > f(x)$  with  $x \in S_{r_0} \subset S_r \subset \bar{S}_r$ , making  $x \notin X - \bar{S}_r$ . As  $0 \leq f(x)$ , we can upperbound the sup with  $f(x)$ . And,  $\forall r < f(x) \leq 1$ , then  $\exists r_0, r < r_0 < f(x)$  with  $x \notin S_{r_0} \supset \bar{S}_r$  and thus  $x \in X - \bar{S}_r$ . Since all binary fraction  $0 < r < f(x)$  (it is key here that  $f(x)$  is bounded above 1 so that all of these  $r$  produce valid  $S_r$ ) are in the set, and these are dense in the reals, we have  $f(x)$  as a lower bound for the sup as well. This fails if  $f(x) = 0$ , but then, the additional 0 element saves us. Regardless we have now proved the identity.

Next, we show  $f^{-1}([0, r))$  is open ( $r$  is now any real number). I claim  $f^{-1}([0, r)) = \cup_{s < r} S_s$ . If  $x \in \cup_{s < r} S_s$ , then  $\exists s < r$  such that  $x \in S_s$  and thus,  $f(x) = \inf(\{1\} \cup \{a : x \in S_a\}) \leq s < r$ . If  $x \notin \cup_{s < r} S_s$ , then  $\forall s < r, x \notin S_s$ . Thus, if  $x \in S_s$ , then  $s \geq r$ ; and of course  $1 \geq r$ . So,  $f(x) = \inf(\{1\} \cup \{a : x \in S_a\}) \geq r$ . So  $x \in \cup_{s < r} S_s$  if and only if  $x \in f^{-1}([0, r))$  making the sets equal; as the union of open sets is open so is  $f^{-1}([0, r))$ . The same argument, using the sup definition of  $f$ , shows that  $f^{-1}((r, 1])$  is open. As,  $f^{-1}((a, b)) = f^{-1}([0, b)) \cap f^{-1}((a, 1])$ , this set is open as well. Finally, any open set in the reals is the union of open balls (one around each point if you like),  $f^{-1}$  of any open set is the union of open sets and is thus open. Thus,  $f$  is continuous as desired.

5. Let  $X$  be a topological space and  $\mathcal{F}$  a collection of closed subsets. Define  $\mathcal{G}$  as the complements of sets in  $\mathcal{F}$ . Then a union of sets in  $\mathcal{G}$  is the complement of an intersection of the corresponding sets in  $\mathcal{F}$ . So

$\mathcal{F}$  has FIP if and only if  $\mathcal{G}$  has no finite subcover of  $X$ . Also  $\mathcal{F}$  has the total intersection property if and only if  $\mathcal{G}$  does not cover  $X$ . Therefore we are done.

6. Let  $S$  be a sequentially compact subset of a metric space.
  - (i) Suppose  $S$  is not closed. Then  $\exists x \notin S$ , such that  $\forall r > 0, B_r(x) \cap S \neq \emptyset$ . Let,  $a_n$  be a point in  $S \cap B_{1/n}(x)$ . Clearly,  $a_n \rightarrow x \notin S$ , and thus all subsequences converge to  $x \notin S$  and thus do not converge in  $S$ , contradicting the sequential compactness of  $S$ .
  - (ii) Suppose  $S$  is not totally bounded. Then let  $r$  be a radius such that there is no  $r$  net of  $S$ . Now define  $a_n$  as follows: Let  $a_0$  be any point in  $S$  and let  $a_n \in S$ , such that  $\forall m < n, d(a_m, a_n) \geq r$ . Such an  $a_n$  exists by the lack of an  $r$  net of  $S$ . But  $S$  sequentially compact implies  $\exists m, n$  such that  $d(a_m, a_n) < r$  which is impossible. So,  $S$  must be totally bounded.
7. Clearly if  $E$  is totally bounded, then it is totally bounded relative to any metric space containing it. Conversely, if  $E$  is totally bounded relative to  $X$ , then there is some finite set of point  $p_1, p_2, \dots, p_n \in X$ , so that every point in  $E$  is within  $\epsilon/2$  of these points. Without loss of generality, we may assume that there is some  $q_i \in E$  in each of these  $\epsilon/2$  balls, or else we omit the corresponding  $p_i$  from the original enumeration. But now,  $\cup_i B_\epsilon(q_i) \supset \cup_i B_{\epsilon/2}(p_i) \supset E$ , by the triangle inequality. And thus for every  $\epsilon$  we have an epsilon net of  $E$  centered around points in  $E$ , making  $E$  totally bounded.
8. Suppose no such  $r$  exists. Then for each  $r$ , in particular for each  $1/n$ , there is some  $x_n$  such that  $\forall \alpha, B_{1/n}(x_n) \not\subseteq U_\alpha$ . Now,  $X$  is compact, so exists  $n_i \in \mathbb{Z}^+, x \in X$ , so that  $x_{n_i} \rightarrow x$ . Now,  $\{U_\alpha\}$  covers  $X$ , so  $\exists \alpha$ , so that  $x \in U_\alpha$ . As these sets are open,  $\exists r$  such that  $B_r(x) \subset U_\alpha$ . And, by convergence,  $\exists k > \frac{2}{r}$ , so that  $d(x_k, x) < \frac{r}{2}$ . But then,  $B_{1/k}(x_k) \subset B_r(x) \subset U_\alpha$ , contradicting our construction of  $\{x_n\}$ , and implying the existence of such an  $r$ .