

## GREEK MATHEMATICS

### (c) SOLUTIONS BY HIGHER CURVES

#### (i.) General

Simpl. in *Cat.* 7, ed. Kalbfleisch 192. 15-25

"Ἔστιν δὲ τετραγωνισμὸς κύκλου, ὅταν τῷ δοθέντι κύκλῳ ἴσον τετράγωνον συστησώμεθα. τοῦτο δὲ Ἀριστοτέλης μὲν, ὡς ἔοικεν, οὕτω ἐγνώκει, παρὰ δὲ τοῖς Πυθαγορείοις ἠύρησθαί φησιν Ἰάμβλιχος, "ὡς δηλὸν ἔστιν ἀπὸ τῶν Σέξτου τοῦ Πυθαγορείου ἀποδείξεων, ὃς ἄνωθεν κατὰ διαδοχὴν παρέλαβεν τὴν μέθοδον τῆς ἀποδείξεως. καὶ ὕστερον δέ, φησὶν, Ἀρχιμήδης διὰ τῆς Λυκομήδους<sup>1</sup> γραμμῆς καὶ Νικομήδης διὰ τῆς ἰδίως τετραγωνιζούσης καλουμένης καὶ Ἀπολλώνιος διὰ τινος γραμμῆς, ἣν αὐτὸς μὲν κοχλιοειδοῦς ἀδελφὴν προσαγορεύει, ἣ αὐτὴ δὲ ἔστιν τῇ Νικομήδους, καὶ Κάρπος δὲ διὰ τινος γραμμῆς, ἣ ἀπλῶς ἐκ διπλῆς κινήσεως καλεῖ, ἄλλοι τε πολλοὶ ποικίλως τὸ πρόβλημα κατεσκεύασαν," ὡς Ἰάμβλιχος ἱστορεῖ.

<sup>1</sup> No meaning can be extracted from *Λυκομήδους*, which is an otherwise unknown word. The correct reading is probably *ἐλικοειδοῦς*, "spiral-shaped."

## SPECIAL PROBLEMS

### (c) SOLUTIONS BY HIGHER CURVES

#### (i.) General

Simplicius, *Commentary on Aristotle's Categories* 7, ed. Kalbfleisch 192. 15-25

The circle is squared when we construct a square equal to the given circle. Aristotle, it would appear, did not know how to do this, but Iamblichus says it was discovered by the Pythagoreans, "as is plain from the proofs of Sextus the Pythagorean,<sup>a</sup> who received the method of the proof from early tradition. And later (he says), Archimedes effected it by means of the spiral-shaped curve,<sup>b</sup> Nicomedes by means of the curve known by the special name *quadratrix*, Apollonius by means of a certain curve which he himself calls *sister of the cochloid*, but which is the same as Nicomedes' curve,<sup>c</sup> Carpus by means of a certain curve which he simply calls that *arising from a double motion*,<sup>d</sup> and many others constructed a solution of this problem in divers ways," as Iamblichus relates.

<sup>a</sup> Sextus (more properly Sextius) lived in the reign of Augustus (or Tiberius) and there is no valid reason for believing the early Pythagoreans solved the problem.

<sup>b</sup> Archimedes himself in his book *On Spirals*, which will be noticed when we come to him, merely uses the spiral to rectify the circle (Prop. 19). But the quadrature follows from *Measurement of a Circle*, Prop. 1.

<sup>c</sup> Nothing further is known of Apollonius's "sister of the cochloid," but Heath (*H.G.M.* i. 232) points out that Apollonius wrote a treatise on the *cochlias*, or cylindrical helix, that the subtangent to this curve can be used to square the circular section of the cylinder, and that the name is sufficiently akin to justify Apollonius in speaking of it as the "sister of the cochloid."

<sup>d</sup> Tannery thought this was the cycloid, but there is no evidence.

GREEK MATHEMATICS


(ii.) *The Quadratrix*

Papp. *Coll.* iv. 30. 45-32. 50, ed. Hultsch 250. 33-258. 19

*Construction of the Curve*

λ'. Εἰς τὸν τετραγωνισμόν τοῦ κύκλου παρελήφθη τις ὑπὸ Δεινοστράτου καὶ Νικομήδους γραμμὴ καὶ τινῶν ἄλλων νεωτέρων ἀπὸ τοῦ περι αὐτὴν συμπτώματος λαβοῦσα τοῦνομα· καλεῖται γὰρ ὑπ' αὐτῶν τετραγωνίζουσα καὶ γένεσιν ἔχει τοιαύτην.

Ἐκκείσθω τετράγωνον τὸ ΑΒΓΔ καὶ περι κέντρον τὸ Α περιφέρεια γεγράφθω ἡ ΒΕΔ, καὶ



κινείσθω ἡ μὲν ΑΒ οὕτως ὥστε τὸ μὲν Α σημεῖον μένειν τὸ δὲ Β φέρεσθαι κατὰ τὴν ΒΕΔ περιφέρειαν, ἡ δὲ ΒΓ παράλληλος αἰεὶ διαμένουσα τῇ ΑΔ τῷ Β σημείῳ φερομένῳ<sup>1</sup> κατὰ τῆς ΒΑ συνακολουθείτω, καὶ ἐν ἴσῳ χρόνῳ ἢ τε ΑΒ κινουμένη

SPECIAL PROBLEMS

(ii.) *The Quadratrix*

Pappus, *Collection* iv. 30. 45-32. 50, ed. Hultsch 250. 33-258. 19

*Construction of the Curve*

30. For the squaring of the circle a certain line was used by Dinostratus and Nicomedes and certain other more recent geometers, and it takes its name from its special property; for it is called by them the quadratrix,<sup>a</sup> and it is generated in this way.

Let ΑΒΓΔ be a square, and with centre Α let the arc ΒΕΔ be described, and let ΑΒ be so moved that the point Α remains fixed while Β is carried along the arc ΒΕΔ; furthermore let ΒΓ, while always remaining parallel to ΑΔ, follow the point Β in its motion along ΒΑ, and in equal times let ΑΒ, moving uni-

<sup>a</sup> Heath (*H.G.M.* i. 225-226) shows that the quadratrix was discovered by Hippias and that he may himself have used it (though this is not absolutely certain) to rectify, and so to square, the circle. He identifies Hippias with Hippias of Elis (*v. supra*, p. 149 n.a.)

<sup>1</sup> σημείου φέρον ἐν φῖ in the mss. was corrected by Torelli.

ὁμαλῶς τὴν ὑπὸ ΒΑΔ γωνίαν, τουτέστιν τὸ Β σημεῖον τὴν ΒΕΔ περιφέρειαν, διανύτω, καὶ ἡ ΒΓ τὴν ΒΑ εὐθείαν παροδεύτω, τουτέστιν τὸ Β σημεῖον κατὰ τῆς ΒΑ φερέσθω. συμβήσεται δὴλον τῇ ΑΔ εὐθείᾳ ἅμα ἐφαρμόζειν ἑκατέραν τὴν τε ΑΒ καὶ τὴν ΒΓ. τοιαύτης δὲ γνωμένης κινήσεως τεμοῦσιν ἀλλήλας ἐν τῇ φορᾷ αἱ ΒΓ, ΒΑ εὐθεῖαι κατὰ τι σημεῖον αἰεὶ συμμεριστάμενον αὐταῖς, ὑφ' οὗ σημείου γράφεται τις ἐν τῷ μεταξὺ τόπων τῶν τε ΒΑΔ εὐθειῶν καὶ τῆς ΒΕΔ περιφερείας γραμμὴ ἐπὶ τὰ αὐτὰ κοίλη, οἷα ἐστὶν ἡ ΒΖΗ, ἡ καὶ χρεώδης εἶναι δοκεῖ πρὸς τὸ τῷ δοθέντι κύκλῳ τετράγωνον ἴσον εὔρειν. τὸ δὲ ἀρχικὸν αὐτῆς σύμπτωμα τοιοῦτόν ἐστιν. ἦτις γὰρ ἂν διαχθῆ τυχοῦσα (πρὸς τὴν περιφέρειαν, ὡς ἡ ΑΖΕ, ἔσται ὡς ὅλη ἡ)¹ περιφέρεια πρὸς τὴν ΕΔ, ἡ ΒΑ εὐθεῖα πρὸς τὴν ΖΘ· τοῦτο γὰρ ἐκ τῆς γενέσεως τῆς γραμμῆς φανερόν ἐστιν.

*Sporus's Criticisms*

λα'. Δυσαρεστεῖται δὲ αὐτῇ ὁ Σπόρος εὐλόγως διὰ ταῦτα. πρῶτον μὲν γὰρ πρὸς ὃ δοκεῖ χρεώδης εἶναι πρᾶγμα, τοῦτ' ἐν ὑποθέσει λαμβάνει. πῶς γὰρ δυνατόν, δύο σημείων ἀρξαμένων ἀπὸ τοῦ Β

¹ πρὸς τὴν . . . ὅλη ἡ add. Hultsch.

formly, pass through the angle ΒΑΔ (that is, the point Β pass along the arc ΒΔ), and ΒΓ pass by the straight line ΒΑ (that is, let the point Β traverse the length of ΒΑ). Plainly then both ΑΒ and ΒΓ will coincide simultaneously with the straight line ΑΔ. While the motion is in progress the straight lines ΒΓ, ΒΑ will cut one another in their movement at a certain point which continually changes place with them, and by this point there is described in the space between the straight lines ΒΑ, ΑΔ and the arc ΒΕΔ a concave curve, such as ΒΖΗ, which appears to be serviceable for the discovery of a square equal to the given circle. Its principal property is this. If any straight line, such as ΑΖΕ, be drawn to the circumference, the ratio of the whole arc to ΕΔ will be the same as the ratio of the straight line ΒΑ to ΖΘ; for this is clear from the manner in which the line was generated.<sup>c</sup>

*Sporus's Criticisms*<sup>b</sup>

31. With this Sporus is rightly displeased for these reasons. In the first place, the end for which the construction seems to be useful is assumed in the hypothesis. For how is it possible, with two points

<sup>c</sup> If ΑΖ=ρ, ∠ΖΑΔ=φ, ΑΒ=α, then the equation of the curve is

$$\frac{\frac{1}{2}\pi}{\phi} = \frac{a}{\rho \sin \phi}$$

or

$$\pi \rho \sin \phi = 2a\phi.$$

<sup>b</sup> These acute criticisms of the quadratrix as a practical method of squaring the circle appear to be well founded. Sporus, who was not much older than Pappus himself, lived towards the end of the third century A.D. He compiled a work called *Κηρία* giving extracts on the quadrature of the circle and duplication of the cube.

κινείσθαι, τὸ μὲν κατ' εὐθείας ἐπὶ τὸ Α, τὸ δὲ κατὰ περιφερείας ἐπὶ τὸ Δ ἐν ἴσῳ χρόνῳ συναποκαταστήσῃαι<sup>1</sup> μὴ πρότερον τὸν λόγον τῆς ΑΒ εὐθείας πρὸς τὴν ΒΕΔ περιφέρειαν ἐπιστάμενον; ἐν γὰρ τούτῳ τῷ λόγῳ καὶ τὰ τάχῃ τῶν κινήσεων ἀνάγκη εἶναι. ἐπεὶ πῶς οἶόν τε συναποκαταστήναι τάχῃσιν ἀκρίτοις χρώμενα, πλὴν εἰ μὴ ἂν κατὰ τύχην ποτὲ συμβῆ; τοῦτο δὲ πῶς οὐκ ἄλογον; ἔπειτα δὲ τὸ πέρασ αὐτῆς ᾧ χρῶνται πρὸς τὸν τετραγωνισμόν τοῦ κύκλου, τουτέστιν καθ' ὃ τέμνει σημεῖον τὴν ΑΔ εὐθείαν, οὐχ εὐρίσκεται. νοείσθω δὲ ἐπὶ τῆς προκειμένης τὰ λεγόμενα καταγραφῆς· ὁπόταν γὰρ αἱ ΓΒ, ΒΑ φερόμεναι συναποκατασταθῶσιν, ἐφαρμόσουσιν τῇ ΑΔ καὶ τομὴν οὐκέτι ποιήσουσιν ἐν ἀλλήλαις· παύεται γὰρ ἡ τομὴ πρὸ τῆς ἐπὶ τὴν ΑΔ ἐφαρμογῆς ἥπερ τομὴ πέρασ αὐτῆς ἐγένετο τῆς γραμμῆς, καθ' ὃ τῇ ΑΔ εὐθείᾳ συνέπιπτεν. πλὴν εἰ μὴ λέγοι τις ἐπινοεῖσθαι προσεκβαλλομένην τὴν γραμμὴν, ὡς ὑποτιθέμεθα τὰς εὐθείας, ἕως τῆς ΑΔ. τοῦτο δ' οὐχ ἔπεται ταῖς ὑποκειμέναις ἀρχαῖς, ἀλλ' ὡς ἂν ληφθεῖη τὸ Η σημεῖον προειλημμένου τοῦ τῆς περιφερείας πρὸς τὴν εὐθείαν λόγου. χωρὶς δὲ τοῦ δοθῆναι τὸν λόγον τοῦτον οὐ χρὴ τῇ τῶν εὐρόντων ἀνδρῶν δόξῃ πιστεύοντας παραδέχεσθαι τὴν γραμμὴν μηχανικωτέραν πῶς οὐσαν [καὶ εἰς πολλὰ προβλήματα χρησιμεύουσιν τοῖς μηχανικοῖς].<sup>2</sup> ἀλλὰ πρότερον παραδεκτέον ἔστι τὸ δι' αὐτῆς δεικνύμενον πρόβλημα.


<sup>1</sup> συναποκαταστήναι coniecit Hultsch.

<sup>2</sup> καὶ . . . μηχανικοῖς interpolatori tribuit Hultsch.

beginning to move from B, to make one of them move along a straight line to A and the other along a circumference to Δ in equal time unless first the ratio of the straight line AB to the circumference BEΔ is known? For it is necessary that the speeds of the moving points should be in this ratio. And how then could one, using unadjusted speeds, make the motions end together, unless this should sometimes happen by chance? But how could this fail to be irrational? Again, the extremity of the curve which they use for the squaring of the circle, that is, the point in which the curve cuts the straight line ΑΔ, is not found. Let the construction be conceived as aforesaid. When the straight lines ΓΒ, ΒΑ move so as to end their motion together, they will coincide with ΑΔ and will no longer cut each other. In fact, the intersection ceases before the coincidence with ΑΔ, yet it was this intersection which was the extremity of the curve where it met the straight line ΑΔ. Unless, indeed, anyone should say the curve is conceived as produced, in the same way that we produce straight lines, as far as ΑΔ. But this does not follow from the assumptions made; the point Η can be found only by assuming the ratio of the circumference to the straight line. So unless this ratio is given, we must beware lest, in following the authority of those men who discovered the line, we admit its construction, which is more a matter of mechanics. But first let us deal with that problem which we have said can be proved by means of it.

Application of Quadratrix to Squaring of Circle

Τετραγώνου γὰρ ὄντος τοῦ ΑΒΓΔ καὶ τῆς μὲν περὶ τὸ κέντρον τὸ Γ περιφερείας τῆς ΒΕΔ, τῆς




δὲ ΒΗΘ τετραγωνιζούσης γινομένης, ὡς προείρηται, δέικνται, ὡς ἡ ΔΕΒ περιφέρεια πρὸς τὴν ΒΓ εὐθείαν, οὕτως ἡ ΒΓ πρὸς τὴν ΓΘ εὐθείαν. εἰ γὰρ μὴ ἔστιν, ἦτοι πρὸς μείζονα ἔσται τῆς ΓΘ ἢ πρὸς ἐλάσσονα.

Ἐστω πρότερον, εἰ δυνατόν, πρὸς μείζονα τὴν ΓΚ, καὶ περὶ κέντρον τὸ Γ περιφέρεια ἡ ΖΗΚ γεγράφθω τέμνουσα τὴν γραμμὴν κατὰ τὸ Η, καὶ κάθετος ἡ ΗΛ, καὶ ἐπιζευχθεῖσα ἡ ΓΗ ἐκβεβλήσθω ἐπὶ τὸ Ε. ἐπεὶ οὖν ἔστιν ὡς ἡ ΔΕΒ περιφέρεια πρὸς τὴν ΒΓ εὐθείαν, οὕτως ἡ ΒΓ,

Application of Quadratrix to Squaring of Circle


If ABΓΔ is a square and BEΔ the arc of a circle with centre Γ, while BHΘ is a quadratrix generated in the aforesaid manner, it is proved that the ratio of the arc ΔΕΒ towards the straight line ΒΓ is the same as that of ΒΓ towards the straight line ΓΘ. For if it is not, the ratio of the arc ΔΕΒ towards the straight line ΒΓ will be the same as that of ΒΓ towards either a straight line greater than ΓΘ or a straight line less than ΓΘ.

Let it be the former, if possible, towards a greater straight line ΓΚ, and with centre Γ let the arc ΖΗΚ be drawn cutting the curve at Η, and let the perpendicular ΗΛ be drawn, and let ΓΗ be joined and pro-



duced to Ε. Since therefore the ratio of the arc ΔΕΒ towards the straight line ΒΓ is the same as the

τουτέστιν ἡ  $\Gamma\Delta$ , πρὸς τὴν  $\Gamma\text{Κ}$ , ὡς δὲ ἡ  $\Gamma\Delta$  πρὸς τὴν  $\Gamma\text{Κ}$ , ἢ  $\text{ΒΕ}\Delta$  περιφέρεια πρὸς τὴν  $\text{ΖΗΚ}$  περιφέρειαν (ὡς γὰρ ἡ διάμετρος τοῦ κύκλου πρὸς τὴν διάμετρον, ἢ περιφέρεια τοῦ κύκλου πρὸς τὴν περιφέρειαν), φανερόν ὅτι ἴση ἐστὶν ἡ  $\text{ΖΗΚ}$  περιφέρεια τῇ  $\text{Β}\Gamma$  εὐθείᾳ. καὶ ἐπειδὴ διὰ τὸ σύμπτωμα τῆς γραμμῆς ἐστὶν ὡς ἡ  $\text{ΒΕ}\Delta$  περιφέρεια πρὸς τὴν  $\text{Ε}\Delta$ , οὕτως ἡ  $\text{Β}\Gamma$  πρὸς τὴν  $\text{Η}\Lambda$ , καὶ ὡς ἄρα ἡ  $\text{ΖΗΚ}$  πρὸς τὴν  $\text{ΗΚ}$  περιφέρειαν, οὕτως ἡ  $\text{Β}\Gamma$  εὐθεῖα πρὸς τὴν  $\text{Η}\Lambda$ . καὶ ἐδείχθη ἴση ἡ  $\text{ΖΗΚ}$  περιφέρεια τῇ  $\text{Β}\Gamma$  εὐθείᾳ· ἴση ἄρα καὶ ἡ  $\text{ΗΚ}$  περιφέρεια τῇ  $\text{Η}\Lambda$  εὐθείᾳ, ὅπερ ἄτοπον. οὐκ ἄρα ἐστὶν ὡς ἡ  $\text{ΒΕ}\Delta$  περιφέρεια πρὸς τὴν  $\text{Β}\Gamma$  εὐθείαν, οὕτως ἡ  $\text{Β}\Gamma$  πρὸς μείζονα τῆς  $\Gamma\Theta$ .  
 λβ'. Λέγω δὲ ὅτι οὐδὲ πρὸς ἐλάσσονα. εἰ γὰρ



δυνατόν, ἔστω πρὸς τὴν  $\text{Κ}\Gamma$ , καὶ περὶ κέντρον τὸ  
 344

ratio of  $\text{Β}\Gamma$ , that is  $\Gamma\Delta$ , towards  $\Gamma\text{Κ}$ , and the ratio of  $\Gamma\Delta$  towards  $\Gamma\text{Κ}$  is the same as that of the arc  $\text{ΒΕ}\Delta$  towards the arc  $\text{ΖΗΚ}$  (for the arcs of circles are in the same ratio as their diameters), it is clear that the arc  $\text{ΖΗΚ}$  is equal to the straight line  $\text{Β}\Gamma$ . And since by the property of the curve the ratio of the arc  $\text{ΒΕ}\Delta$  towards  $\text{Ε}\Delta$  is the same as the ratio of  $\text{Β}\Gamma$  towards  $\text{Η}\Lambda$ , therefore the ratio of  $\text{ΖΗΚ}$  towards the arc  $\text{ΗΚ}$  is the same as the ratio of the straight line  $\text{Β}\Gamma$  towards  $\text{Η}\Lambda$ . And the arc  $\text{ΖΗΚ}$  was proved equal to the straight line  $\text{Β}\Gamma$ ; therefore the arc  $\text{ΗΚ}$  is also equal to the straight line  $\text{Η}\Lambda$ , which is absurd. Therefore the ratio of the arc  $\text{ΒΕ}\Delta$  towards the straight line  $\text{Β}\Gamma$  is not the same as the ratio of  $\text{Β}\Gamma$  towards a straight line greater than  $\Gamma\Theta$ .

32. I say that neither is it equal to the ratio of  $\text{Β}\Gamma$  towards a straight line less than  $\Gamma\Theta$ . For, if it is possible, let the ratio be towards  $\text{Κ}\Gamma$ , and with centre

Γ περιφέρεια γεγράφθω ἡ ZMK, καὶ πρὸς ὀρθὰς τῇ ΓΔ ἢ ΚΗ τέμνουσα τὴν τετραγωνίζουσαν κατὰ τὸ Η, καὶ ἐπιζευχθεῖσα ἡ ΓΗ ἐκβεβλήσθω ἐπὶ τὸ Ε. ὁμοίως δὲ τοῖς προγεγραμμένοις δείξομεν καὶ τὴν ZMK περιφέρειαν τῇ ΒΓ εὐθείᾳ ἴσην, καὶ ὡς τὴν ΒΕΔ περιφέρειαν πρὸς τὴν ΕΔ, τουτέστιν ὡς τὴν ZMK πρὸς τὴν MK, οὕτως τὴν ΒΓ εὐθεῖαν πρὸς τὴν ΗΚ. ἐξ ὧν φανερόν ἐστι ἴση ἔσται ἡ MK περιφέρεια τῇ ΚΗ εὐθείᾳ, ὅπερ ἄτοπον. οὐκ ἄρα ἔσται ὡς ἡ ΒΕΔ περιφέρεια πρὸς τὴν ΒΓ εὐθεῖαν, οὕτως ἡ ΒΓ πρὸς ἐλάσσονα τῆς ΓΘ. ἐδείχθη δὲ ὅτι οὐδὲ πρὸς μείζονα πρὸς αὐτὴν ἄρα τὴν ΓΘ.

Ἔστι δὲ καὶ τοῦτο φανερόν ἐστι ἡ τῶν ΘΓ, ΓΒ εὐθειῶν τρίτη ἀνάλογον λαμβανομένη εὐθεῖα ἴση ἔσται τῇ ΒΕΔ περιφερείᾳ, καὶ ἡ τετραπλασίων αὐτῆς τῇ τοῦ ὅλου κύκλου περιφερείᾳ. εὐρημένης δὲ τῇ τοῦ κύκλου περιφερείᾳ ἴσης εὐθείας πρόδηλον ὡς δὴ καὶ αὐτῶ τῶ κύκλῳ ῥᾶδιον ἴσον τετράγωνον συστήσασθαι· τὸ γὰρ ὑπὸ τῆς περιμέτρου τοῦ κύκλου καὶ τῆς ἐκ τοῦ κέντρου διπλάσιόν ἐστι τοῦ κύκλου, ὡς Ἀρχιμήδης ἀπέδειξεν.

### 3. TRISECTION OF AN ANGLE

#### (a) TYPES OF GEOMETRICAL PROBLEMS

Papp. Coll. iv. 36. 57-59, ed. Hultsch 270. 1-272. 14

λς'. Τὴν δοθείσαν γωνίαν εὐθύγραμμον εἰς τρία ἴσα τεμεῖν οἱ παλαιοὶ γεωμέτραι θελήσαντες ἠπόρησαν δι' αἰτίαν τοιαύτην. τρία γένη φαιμέν εἶναι

Γ let the arc ZMK be described, and let KH at right angles to ΓΔ cut the quadratrix at H, and let ΓH be joined and produced to E. In similar manner to what has been written above, we shall prove also that the arc ZMK is equal to the straight line ΒΓ, and that the ratio of the arc ΒΕΔ towards ΕΔ, that is, the ratio of ZMK towards MK, is the same as that of the straight line ΒΓ towards ΗΚ. From this it is clear that the arc MK is equal to the straight line ΚΗ, which is absurd. The ratio of the arc ΒΕΔ towards the straight line ΒΓ is therefore not the same as the ratio of ΒΓ towards a straight line less than ΓΘ. Moreover it was proved not the same as the ratio of ΒΓ towards a straight line greater than ΓΘ; therefore it is the same as the ratio of ΒΓ towards ΓΘ itself.

This also is clear, that if a straight line is taken as a third proportional to the straight lines ΘΓ, ΓΒ it will be equal to the arc ΒΕΔ, and four times this straight line will be equal to the circumference of the whole circle. A straight line equal to the circumference of the circle having been found, a square can easily be constructed equal to the circle itself. For the rectangle contained by the perimeter of the circle and the radius is double of the circle, as Archimedes demonstrated.<sup>a</sup>

### 3. TRISECTION OF AN ANGLE

#### (a) TYPES OF GEOMETRICAL PROBLEMS

Pappus, Collection iv. 36. 57-59, ed. Hultsch 270. 1-272. 14

36. When the ancient geometers sought to divide a given rectilinear angle into three equal parts they were at a loss for this reason. We say that there

<sup>a</sup> See *supra*, pp. 316-321.