

Dec. 6, 99
Solutions to 2nd Math Xa Midterm


- ① (a) false - inflection points can only occur where $g''(x) = 0$, here $g''(e) = 4 \neq 0$
- (b) true - we know that critical points include points where $g' = 0$, here we're told $g'(e) = 0$
- (c) false - g would have to be concave down for this to be true at $x=e$, and $g''(e) = 4$, (which is positive \Rightarrow concave up)
- (d) true - for the same reasons (c) is false. Since $g'(e) = 0$ and $g''(e) = 4$ then the second derivative test implies $x=e$ is a local minimum
- (e) false - we already know $g(x)$ has a local minimum at $x=e$
- (f) false - since the polynomial has a leading coefficient which is negative then $g(x) \rightarrow -\infty$ as $x \rightarrow \infty$, so $g(x)$ has no absolute minimum, only local minimums

② try $\frac{(x-2)^4}{(x+1)x^2(x-1)} = Q(x)$

- has vertical asymptotes at $x = -1, 0$ and 1 because of the factors in the denominator,
- has just the one x -intercept at $x = 2$ because that's the only root for the numerator
- is negative exactly where $(x+1)(x-1)$ is negative, since the other terms, x^2 and $(x-2)^4$ are always positive, and this is precisely when $-1 < x < 1$
- note $\lim_{x \rightarrow \infty} Q(x) = \lim_{x \rightarrow -\infty} Q(x) = 1$ because the ratio of the leading terms for numerator and denominator equals 1.

Solutions continued

③ (a)



(b) $f'(x) = k(x - (-\pi))(x - 2\pi)$
and $f'(0) = -2 = k(0 + \pi)(0 - 2\pi)$
 $= k(-2\pi^2)$
so $k = \frac{-2}{-2\pi^2} = \frac{1}{\pi^2}$


so $f'(x) = \frac{1}{\pi^2}(x + \pi)(x - 2\pi)$

(c) x -coordinate of vertex is halfway between the two roots at $-\pi$ and 2π , at $\frac{2\pi + (-\pi)}{2} = \frac{\pi}{2}$

$f'(\frac{\pi}{2}) = \frac{1}{\pi^2}(\frac{\pi}{2} + \pi)(\frac{\pi}{2} - 2\pi)$
 $= \frac{1}{\pi^2}(\frac{3}{2}\pi)(-\frac{3}{2}\pi) = -\frac{9}{4}$

(d) $f'(x) = \frac{1}{\pi^2}(x + \pi)(x - 2\pi)$
 $= \frac{1}{\pi^2}x^2 - \frac{\pi}{\pi^2}x - \frac{2\pi^2}{\pi^2}$
so $f''(x) = \frac{2}{\pi^2}x - \frac{1}{\pi}$

(e) (i) use the first derivative test: f' goes from positive to negative at $x = -\pi \Rightarrow$ local maximum



(ii) and likewise the only place that f' goes from negative to positive is at $x = 2\pi \Rightarrow$ local minimum

(iii) since f' is a parabola, degree 2, then F must be a cubic polynomial, and cubics don't have abs. max or min (range is $(-\infty, \infty)$)

④ (a) $W(100)$ is the amount of water in the reservoir after 100 seconds have passed

8. A, B, E, F, G

9a. $F(x) = 2000 [1999^x - x^{1999}]$

$F'(x) = 2000 [(1999 \ln 1999) 1999^x - 1999 x^{1998}]$

b. $F(x) = \pi \ln \left(\frac{x^2}{\sqrt{3x}} \right)$

use log rules to simplify:

$F(x) = \pi (\ln x^2 - \ln \sqrt{3x})$

$F(x) = \pi (2 \ln x - \frac{1}{2} \ln 3x)$

$F(x) = \pi (2 \ln x - \frac{1}{2} \ln 3 - \frac{1}{2} \ln x)$

$F'(x) = \pi \left(\frac{2}{x} - 0 - \frac{1}{2x} \right)$
 $= \frac{2\pi}{x} - \frac{\pi}{2x} = \frac{3\pi}{2x}$

Common errors: NOT simplifying the log completely before differentiating
 NOT distributing π

10a. linear equation.

slope: $\frac{139 - 271}{7 - 0} = -19$

equation: $L(t) = -19t + 271$


Exponential equation: Find b:

$139 = 271 \cdot b^7$

$\frac{139}{271} = b^7$

$\left(\frac{139}{271}\right)^{1/7} = b$

$E(t) = 271 \cdot \left(\frac{139}{271}\right)^{t/7}$



Common errors: not showing where the two graphs intersect. not ~~showing~~ providing enough detail (e.g. labelling intercepts)

C. linear model:

$0 = L(t) = -19t + 271$

$t = \frac{271}{19} = 14.26 \text{ days}$

exponential model:

the substance never completely

decays. the equation

$E(t) = 0 = 271 \left(\frac{139}{271}\right)^{t/7}$

has no solutions.

Common errors: trying to solve

$0 = 271 \left(\frac{139}{271}\right)^{t/7}$ by taking

\ln (or \log) of both sides.

This doesn't work because you can't take \ln (or \log) of 0.

