

**Conformal Dynamics Homework 6**  
Math 275 — Harvard University — Fall 2001

1. Let  $X = \mathbb{C}/\Lambda$  be a complex torus. Given an example of a smooth, locally affine map  $f : X \rightarrow X$  which is *not* a Teichmüller map.  
(This means (a) near every point  $p \in X$  there is a complex coordinate system in domain and range such that  $f(x + iy) = x + Kiy$ ; but (b) one cannot find a holomorphic quadratic differential  $\phi \in Q(X)$  such that  $\mu(f) = k\bar{\phi}/|\phi|$ .)
2. Let  $f(z)$  be a critically finite rational map. Prove that the Julia set of  $f$  is connected.
3. Prove that  $\chi(\mathcal{O}_f) \leq 0$  for any critically finite branched covering  $f : S^2 \rightarrow S^2$ .
4. Give an example of a critically finite branched cover of the sphere with orbifold of signature  $(2, 2, 2, 2)$  that is *not* combinatorially equivalent to a rational map.
5. Let  $f : S^2 \rightarrow S^2$  be a critically finite branched covering such that every branch point is periodic, and such that  $f^{-1}(p) = p$  for some  $p \in P(f)$ . Prove that  $f$  is combinatorially equivalent to a polynomial.
6. Prove that the boundary of the Mandelbrot set is not a countable union of smooth arcs.
7. Let  $\Gamma \cong \mathbb{Z} * \mathbb{Z}$  be a Kleinian group with connected domain of discontinuity  $\Omega \subset \widehat{\mathbb{C}}$ . Suppose that  $X = \Omega/\Gamma$  is a closed surface of genus two. Prove that the limit set  $\Lambda(\Gamma)$  is a Cantor set.
8. Prove that  $\Lambda(\Gamma)$  has measure zero.
9. Give a detailed proof of the puzzle theory result that ‘an only child is excellent’; include an analysis of the cases not treated by Milnor.
10. Let  $s(z) = z^2$  and let  $E \subset S^1$  be a closed set such that  $s(E) = E$  and  $s|_E$  is a bijection. Prove that  $E$  is finite.