

A DISCRETE GAUSS-BONNET TYPE THEOREM

OLIVER KNILL

ABSTRACT. We discuss a curvature theorem for subgraphs of the flat triangular tessellation of the plane. These graphs play the analogue of domains in two dimensional Euclidean space. We show that with the curvature $K(p) = 2|S_1(p)| - |S_2(p)|$ the discrete Hopf Umlaufsatz $\sum_{p \in \delta G} K(p) = 12\chi(G)$ holds, where $\chi(G)$ is the Euler characteristic of the graph, δG is the boundary of G and where $|S_r|$ is the arc length of the sphere of radius r in G .

Dedicated to Ernst Specker to his 90th birthday.

1. INTRODUCTION

For a domain G in the plane with smooth boundary C , the **Gauss-Bonnet formula** or **Umlaufsatz** $\int_C K(s) ds = 2\pi\chi(G)$ relates the curvature $K(s)$ of the boundary curve C with the Euler characteristic χ of the region G . For a simply connected smooth region G for which the boundary is a simple closed curve, the total boundary curvature is 2π . **Hopf's Umlaufsatz** relates a differential geometric quantity, the boundary curvature, with a topological invariant, the Euler characteristic. In differential geometry, curvature needs a differentiable structure, while Euler characteristic does not. It is the transcending property between different mathematical branches which makes Gauss-Bonnet type results interesting. The discrete version of the classical Hopf Umlaufsatz [4] is of combinatorial nature; curvature is an integer. The result applies to some two dimensional graphs which are part of a flat two dimensional background graph X , where dimensionality is defined inductively. We need some assumptions on G to equate the total boundary curvature with the Euler characteristic.

We use the curvature $K(p) = 2|S_1(p)| - |S_2(p)|$, where $S_r(p)$ is the arc length of the sphere $S_r(p)$ at the point p . The sphere S_r is a subgraph of G containing as vertices points of geodesic distance r and edges consisting of pairs (q, q') in $S_r(p)$ such that q and q' have distance 1. Gauss-Bonnet results are more subtle for this curvature K than for the first order curvature $K_1(p) = 6 - |S_1(p)|$ in the interior or $K_1(p) = 3 - |S_1(p)|$ at the boundary. While $\sum_{p \in G} K_1(p) = 6\chi(G)$ holds for any two dimensional graph G with one dimensional boundary, the sum $\sum_G K(p)$ of all curvatures $K(p) = 2|S_1(p)| - |S_2(p)|$ is only equal to $30\chi(D)$ if more smoothness is present for the two dimensional graph G . In this article, we see this subtlety to appear in a simpler Umlaufsatz situation, where the graph is a subgraph of a fixed flat background lattice.

Date: Feb. 11, 2010, final revised Nov. 8, 2010.

1991 Mathematics Subject Classification. Primary: 05C10, 57M15 .

Key words and phrases. Graph theory, Gauss-Bonnet, Curvature.




FIGURE 1. The numbers near each vertex indicate the curvature of the point. At each of a few chosen points, the spheres of radius 1 and 2 in G are drawn. Adding up the curvatures over the boundary gives 12. If a point has as a neighborhood a disc of radius 2, the curvature is zero.

The main result in this paper is the formula $\sum_{p \in \partial G} K(p) = 12\chi(G)$ which holds for discrete domains G and for the **second order** curvature K . To do so, we need to specify precisely what a smooth domain is. The background lattice X plays the role of the two dimensional plane. Its vertices are the set of points $\{k(1,0) + l(1,\sqrt{3})/2 \mid k,l \text{ integers}\}$. The edges consist of pairs for which the Euclidean distance is 1. In the infinite graph X , every point p has 6 neighbors. Together with edges formed by neighboring vertices, these points form the **unit sphere** $S_1(p)$, a subgraph of X . Similarly, any sphere $S_2(p)$ of radius 2 in this discrete plane has length $|S_2| = 12$. The **curvature** $K = 2|S_1| - |S_2|$ is zero at every point of the background lattice X .

2. TOPOLOGY OF THE PLANAR TRIANGULAR LATTICE

A subset G of the triangular lattice X defines a **graph** (V, E) , where $V \subset X$ is the set of vertices in G and where E is a subset of edges (p, q) in X , pairs in V having distance 1 within X . We often equate G with the vertex set V because V defines the graph G for the situations we are interested in. We start by defining a dimension for graphs. To our best knowledge, this notion has not yet have appeared, even so in the graph theory literature, several notions of dimension exist. The inductive definition of dimension is rather general and does not require the graph to be a subset of X .

Definition. A **sphere** $S_r(p)$ is a subgraph G of X whose vertices are the set of points in G which have geodesic distance r to p normalized so that adjacent points have distance 1 within G . The edges of the sphere S_r are all pairs (p, q) with $p, q \in S_r(p)$ for which (p, q) is in E . A **disc** $B_r(p)$ in the graph G is the set of points q which have distance $d(q, p) \leq r$ in G . A **vertex** p of a graph $G = (V, E)$ is called **zero dimensional**, if p is not connected to any other vertex. A **subgraph** G of X is called **zero dimensional** if every point $p \in G$ is zero dimensional in G . Such a graph has no edges. A point p of V is called **one dimensional** if $S_1(p)$ is zero dimensional, where $S_1(p)$ is the unit sphere of p within G . A finite subset G of X is called **one dimensional** if any of the points in G is one dimensional. A point p of G is called **two dimensional**, if $S_1(p)$ is a one dimensional graph. A subset G of X is called **two dimensional**, if every vertex p of G is two dimensional.

Remarks. a) To illustrate this, we look at the Platonic solids. The cube and the dodecahedron are one dimensional. The isocahedron and octahedron are two dimensional. The tetrahedron is three dimensional. The cube and dodecahedron become two dimensional after a stellation of faces. The tetrahedron becomes two dimensional after truncating corners and then stellating the hexagonal faces.
b) Dimension could be generalized by defining inductively a **fractional dimension** by adding 1 to the average fractional dimensions of points in $S_1(p)$.

Definition. A point p in G called an **interior point** of G if the sphere $S_1(p)$ in the graph G is the same than the sphere $S_1(p)$ in the background graph X . In other words, for an interior point, the sphere $S_1(p)$ is a connected one dimensional graph without boundary, a closed circle. A point p of a two dimensional graph G is a **boundary point** of G , if it is not an interior point in G but has a neighbor in G which is an interior point. For a boundary point the sphere $S_1(p)$ is a union of one dimensional arcs. The **boundary** of G is the set of boundary points of G . The **interior** of G is the set of interior points of G .

Remarks. a) The set of subsets $\{A \subset \text{int}(G)\} \cup \{G\}$ defines a topology on G such that the interior of G is open and the boundary δG is closed.
b) The interior of a two dimensional graph G is not necessarily two dimensional. The disc of radius 1 in X for example has a single interior point so that the interior is zero dimensional.
c) Two dimensionality of a graph has no relation with being planar. The tetrahedral graph for example is three dimensional in our sense but planar. A triangularization of a torus like seen in Figure 16 is two dimensional but not planar.
d) Topologically, the triangular graph X is the only simply-connected two dimensional flat graph without boundary.

Definition. A subset G of X is a **domain** if the following 5 conditions are satisfied:

- (i) G is a two dimensional subgraph of X .
- (ii) Every point of G is either an interior point or a boundary point.
- (iii) The set of boundary points in G is a one dimensional graph.
- (iv) If two vertices p, q in G have distance 1 in X , then (p, q) is an edge in G .
- (v) Two interior points in G with a common boundary point have distance 1 or are both adjacent to a third interior point.

A domain G is called a **finite domain**, if it is a finite graph which is a domain. G is called a **smooth domain**, if G and its complement are both a domain.

12

0

-12




FIGURE 2. Examples of domains. In each of the three pictures, the integer in the upper right corner is the total boundary curvature of the domain.

These assumptions (i),(ii), (iii) are natural. Condition (iv) assures that no unnatural fissures can exist. Condition (v) assures that the connectivity topology of the domain and the connectivity topology of the interior set are the same.

Remarks. a) We could require the interior of a domain to be two dimensional but prefer not to do that because the proof of Theorem 2 simplifies if we do not make this assumption. The additional requirement would lift a difficulty on a different level. For us it will be important to look at the dimension of points with respect to the interior of G .

b) The conditions (i)-(v) have analogue statements in the continuum, where they are necessary for the classical Gauss-Bonnet to be true: we can not have hairs sticking out of the domain for example. The closure of the complement of a domain is a domain too and we can not just leave out part of the boundary. Also in the continuum, it should not happen that parts of domains are tangent to each other. We also can not allow the boundary to be two dimensional, like for the Mandelbrot set.

c) For a smooth domain, we can look at the interior $H' = \text{int}(G')$ of the complement G' of $\text{int}(G)$. Then, the boundaries satisfy $\delta(G) = \delta(G')$. The three sets $\text{int}(G')$, $\text{int}(G)$ and $\delta(G) = \delta(G')$ partition the graph X .

The following lemma allows us to deal efficiently with eligible regions and eliminate many subsets which are not regions. It says that the set of interior points determines the region as well as its boundary.

Lemma 1. *Let G be a domain and $H = \text{int}(G)$ be the set of interior points of G . Then $G = \bigcup_{q \in H} B_1(q)$, where $B_1(q)$ is the disc of radius 1 in X . Especially, the interior set $H = \text{int}(G)$ determines the domain G completely.*

Proof. If a point p is in G , then it is either an interior point or a point adjacent to an interior point. Therefore $G \subset \bigcup_{q \in H} B_1(q)$. On the other hand, if p is in $\bigcup_{q \in H} B_1(q)$, then $p \in B_1(q)$ for some q . Because $q \in G$ and $S_1(q) \in G$ by definition of being an interior point, we have $p \in G$. \square

3. CURVATURE

Definition. Let $|S_r(p)|$ denote the number of edges in the sphere $S_r(p)$. We call it the **arc length** of the sphere S_r .




FIGURE 6. The first picture is a smooth domain. It is not connected. The two parts S_1, S_2 of the regions are at first separated enough to get a curvature 24. In the second case, the sphere S_2 feels part of the other region S_1 and the total curvature of the boundary is not a multiple of 12.

Note that $|S_1|$ is not necessarily the number of vertices in S_1 . Similarly, $|S_2|$ is the number of edges in S_2 which is not always equal to the number of vertices in S_2 .

Definition. The **curvature** of a boundary vertex p in a region G is defined as

$$K(p) = 2|S_1(p)| - |S_2(p)|.$$

The **curvature** of a finite domain G is the sum of the curvatures over the boundary.

Remarks. a) This definition is motivated by differential geometry where the curvature in the interior of a smooth surface is

$$K = \lim_{r \rightarrow 0} \frac{2|S_r| - |S_{2r}|}{2\pi r^3}$$

and the curvature of a smooth curve in the plane is

$$K = \lim_{r \rightarrow 0} \frac{2|S_r| - |S_{2r}|}{2r^2}$$

for a point p on the boundary of a flat region G and where $|S_r(p)|$ is the arc-length of $S_r(p) \cap G$.

b) Every point $q \in G$ of distance 2 in X to $p \in G$ belongs to $S_2(p)$ whether there is a connection within G from p to q or not.

Definition. A **curve** γ in a smooth domain G is a sequence of points x_0, \dots, x_n in the interior of G such that $d(x_i, x_{i+1}) = 1$ and consequently (x_i, x_{i+1}) is an edge of G . A curve is a **closed curve** if $x_0 = x_n$. In graph theory, a curve is also called a chain. A closed curve is a **nontrivial closed curve** if its length is larger than 1. It is called a **simple closed curve**, if all points x_0, x_1, \dots, x_{n-1} are different and $x_0 = x_n$. A domain is called **simply connected** if every closed curve $\{x_1, \dots, x_n\}$ in the interior H of G can be deformed to a trivial closed curve within G , where

a **deformation** of a curve within G is a composition of finitely many elementary deformation steps $\{x_1, \dots, x_n\} \rightarrow \{y_1, \dots, y_m\}$ with $\sum_i d(x_i, y_{j(i)}) = 1$ and such that x_i, y_i are in H and $d(x, y)$ is the geodesic distance in the graph with $d(x, y) = 1$ if (x, y) is an edge. As in the continuum, simply connectedness means that any closed curve in the interior of G can be deformed to a point within the interior of G .

4. THE CURVATURE 12 THEOREM

The main result of this paper is a discrete version of the Umlaufsatz. It will be generalized to more general domains below.

Theorem 2 (Curvature 12 Umlaufsatz). *The total boundary curvature of a finite, smooth and simply connected domain G is 12.*

Proof. For the proof, it suffices to look at local deformations of the region. We start with an arbitrary finite, simply connected, smooth region G and find a procedure to remove interior points near the boundary while keeping it simply connected and also preserve the curvature. Removing one point affects the curvatures in a disc of radius 2 only:

Lemma 3 (Curvature is local). *Let G_1, G_2 be two regions and p be a point in both G_1 and G_2 . Let $U_1 = B_2(p) \subset G_1$ and $U_2 = B_2(p) \subset G_2$ be the discs of radius 2 in G_1 and G_2 respectively. Define $H_i = G_i \setminus \{p\}$. If $U_1 = U_2$, then*

$$\sum_{p \in H_1} K(p) - \sum_{p \in H_2} K(p) = \sum_{p \in G_1} K(p) - \sum_{p \in G_2} K(p).$$

The lemma tells that if we remove a point from a region, then the total curvature-change can be read off from the curvature-changes in a disc of radius 2. We could therefore check all possible configurations in discs of radius 2 and compare the total curvature before and after the center point is removed. We indeed did this with the help of a computer. This helped us also to identify the failure cases and so define what a domain is.

But checking all possible local deformations is not a complete proof yet. We also need to know that there is always a point which we can remove without changing the topology of G or its complement. It turns out that this question is of more global nature. Take a ring shaped region for example which has a one dimensional interior. No point can be removed without the curvature to change. See the right picture in Figure 12. The key is to look at the dimension of points in the interior of G and distinguish points which are one dimensional in $\text{int}(G)$ and points which are two dimensional in $\text{int}(G)$. A zero dimensional interior means for a simply connected region that the graph is the disc of radius 1 in X . By removing interior points, we want to reach this situation.

We can focus on the **topology of interior points** because $\text{int}(G)$ defines G by Lemma 1. It is enough to check therefore what happens if we remove interior points. Our goal is to show:

Proposition 4 (Trimming a tree). *For any simply connected smooth region G for which the interior set H has more than one point, it is possible to remove an interior point p from H , such that the new region defined by $H \setminus \{p\}$ remains*




FIGURE 7. The picture illustrates the process of pruning a tree. To shrink the region, we have to remove alternatively two or one dimensional interior points until only one interior point is left. The left picture shows a simply connected region where one branch can be trimmed. We get the tree to the right, where only ridges and bridges are left and all branches have been pruned. We would now have to start etching the ridges. There are four end ridges to choose from. The simple connectivity assures that at least one end ridge exists.

a simply connected smooth region with one interior point less and such that the curvature does not change.

Theorem 2 follows from this proposition. To prove the later, we need some more terminology:

Definition. Assume we are given a smooth and simply connected region G with interior H . Denote by H_1 the points in H which are one dimensional in H . Similarly, call H_2 the set of points in H which are two dimensional in H . Connected components of H_1 are called either **branches** or **bridges**. Connected components of H_2 are called **ridges**. A **branch** of G is a connected component of H_1 for which at least one point has only one interior neighbor. All other connected components of H_1 are called **bridges**.

The set $\text{int}(G)$ is the union of points which are two dimensional in $\text{int}(G)$ and points which are one dimensional in $\text{int}(G)$. We will use two procedures called **pruning** and **etching** to make the region smaller. The pruning procedure removes a one dimensional interior point at branches. The etching procedure removes a two dimensional interior point at ridges.

Let us start with the **pruning procedure** which removes interior points which are one dimensional in $\text{int}(G)$. It allows us to remove one dimensional branches until we can no more reduce one dimensional points in $\text{int}(G)$. Removing one dimensional parts will make sure that there will be two dimensional ridges ready for the etching procedure. If edging is no more possible, we will reach a situation where pruning needs to be done again. Figure 9 shows the situations which can occur locally at a point of a branch.






FIGURE 8. The left picture shows the process of trimming of branches, the right picture shows a sequence of edging steps. Proposition 4 claims that one of the steps can be applied to a simply connected region with more than one interior points.

12

12



12

12





FIGURE 9. The figure shows a one dimensional interior point and its possible neighbors. It can have either 1,2 or 3 neighbors. There are 4 possibilities up to a rotation. The situation of 4 or more neighbors makes the interior point two dimensional in $\text{int}(G)$.

12

12

12




FIGURE 10. A two dimensional interior point at the boundary can have either 2 or 3 or 4 interior points as neighbors. We see all the possible configurations with either 2 or 3 interior neighbors up to a rotation or a reflection.

12

12

12




FIGURE 11. Situations, where a two dimensional interior point at the boundary has 4 interior points as neighbors. In all these cases, the middle point can not be removed while keeping the region a smooth region. If a point has 5 interior points as neighbors, it automatically must have 6 neighbors and is no more connected to a boundary point of G .

After reducing one dimensional branches, the tree still can have one dimensional parts: these are two dimensional ridges connected with one dimensional bridges which can not be pruned immediately without changing the topology.

The **etching procedure** is invoked if no one dimensional branches are left any more. The region consists now of two dimensional ridges connected with bridges. Our goal is to see that we can remove a two dimensional interior point of a ridge.

The simply connectivity implies that there is a ridge which has only one bridge connected to it. To see this, look at a new graph, which contains the two dimensional ridges as vertices and one dimensional bridges as edges. This graph contains no closed curves which are not contractible, is connected and must therefore be a **tree** with at least one end point. We can consequently focus our discussion to such an end-ridge for which only one one dimensional bridge is attached. We are able to remove a boundary point on the opposite side of that region, where no branches can be and where the boundary is smooth.




FIGURE 12. The left picture shows a bridge. No interior point which is one dimensional in $\text{int}(G)$ can be removed. The middle pictures shows a situation where no interior point in $\text{int}(G)$ can be removed. We would first have to trim some branches. The right picture shows a situation where no trimming is possible any more. The region is not simply connected.

Once the etching process is over, we can again start pruning branches, or we are left with a region with only one interior point. If a region G can no more be pruned and edged then $\text{int}(G)$ consists of only one point and G consists of only 7 points and in this case, we know the total curvature is 12.

Since pruning and etching did not change the curvature, we demonstrated that one can reduce every simply connected region to a situation with only one interior point. This completes the proof of the curvature 12 theorem. \square

5. DISCRETE GAUSS-BONNET

To generalize the Umlaufsatz to domains which are not necessarily simply connected we first **define** the Euler characteristic of a region using Euler's formula:

Definition. A **face** in a domain $G = (V, E)$ is a triangle (p, q, r) of 3 points in G for the which all three points have mutual distance 1. An **edge** in G is a pair p, q of points in G of distance 1. A **vertex** is a point in G . Denote by f the number of faces in G , by $e = |E|$ the number of edges and $v = |V|$ the number of vertices. The **Euler characteristic** $\chi(G)$ of the domain G is defined as $\chi(G) = v - e + f$.

Remarks. The Euler characteristic of $\text{int}(G)$ and G is the same if G is a smooth region but we do not need this fact. For a simply connected region for example the Euler characteristic is 1.

Lemma 5. *The Euler characteristic does not change under the pruning and etching operations defined above as long as the subgraphs remain smooth domains. Filling in a simply connected hole adds 1 to the Euler characteristic.*

Proof. Both removing an end point of a one dimensional branch, as well as removing a two dimensional point from a ridge does not change χ . The number of interior points of a smooth region is $2f - e + \chi$ as can be proven inductively: Each added face corresponds to add 2 edges unless we generate a new interior point, where one edge adds a face. The second statement follows from the fact that a hole H in a region G has $\chi(H) = 1$ and that $\chi(H) + \chi(G) = \chi(H \cup G) - \chi(H \cap G)$ with $\chi(H \cap G) = 0$ because $H \cap G$ is a closed path. \square




FIGURE 13. Joining two regions changes the total curvature by 12. One can prove this by trimming both sides. The left hand side trims down to a disconnected domain consisting of two discs of radius 1, the right hand side will end up with a disc of radius 1.

Theorem 6 (Discrete Gauss-Bonnet theorem). *If G is a finite smooth domain G with boundary C , then*

$$\sum_{p \in C} K(p) = 12\chi(G).$$

We could use the same pruning-etching technique as before. However, pruning and etching can lead to final situations which have no end points like a ring. Instead of classifying all these final situations, it is easier to reduce the general situation to a simply connected situation. There are two ways to change the topology: we can either build bridges between different connected components or we can fill holes to make the region simply connected. Merging different unconnected components works as long as their complement are smooth, both the Euler characteristic as well as the total curvature add up.

We can assume the region to be connected, because both the curvature as well as the Euler characteristic are additive with respect to adding disjoint domains as long as the complement is a domain. To illustrate this more, we can join two separated regions along with a one dimensional bridge. The curvature drops by 12, the number of connected components drops by 1. See Figure 13.

Definition. A **hole** W of G is a bounded simply connected smooth region such that the interior $\text{int}(W)$ is a connected component of the complement G' of G and the interior $\text{int}(G)$ is a subset of the complement of W . Holes are assumed to be simply connected since otherwise G would be disconnected in which case the discussion can be delegated to each connected component of G . By definition, the hole W and the region G share a common one dimensional boundary.

By Theorem 2, the hole has total curvature 12. A key observation is that the point-wise curvatures at the inner boundary of G enclosing the hole are just the negative of the corresponding point-wise curvatures of the hole. Figure 14 illustrates




FIGURE 14. Filling a simply connected hole in a larger region adds to the curvature exactly the same amount than the total curvature of the hole. The point-wise curvatures of the removed inside region matches the curvatures of the inner outside region. Filling a hole increases the curvature by 12 and the Euler characteristic by 1.




FIGURE 15. When cutting a ring, the curvature changes from 0 to 12. Only the last region is a smooth region. The second last is a region but not smooth because the complement is not a region.

this. This follows almost from the definition of curvature and the fact that the circles $|S_1(p) \cap W| + |S_1(p) \cap G| = 6$ and $|S_2(p) \cap W| + |S_2(p) \cap G| = 12$ so that $K_W(p) + K_G(p) = 0$. Alternatively, we could also cut rings, as Figure 15 illustrates.

6. COMPACT FLAT GRAPHS

If we introduce identifications in the hexagonal background graph X , the topology of the background space changes. Identifying points along two parallel lines for example produces a flat cylinder. With a triangular tiling, we can tessellate a torus. Because there is no boundary now, the sum of the curvatures is zero, which is the Euler characteristic. There are many different non-isometric graphs which lead to such **twisted tori**. As **graphs** they can be different even if the number of faces, edges and vertices are the same.

The notion of regular domain can be carried over to subsets of such a twisted background torus X . We assume that it is large enough so that $S_2(p)$ is a circle at every point. We still have:

Theorem 7. *If G is a domain in a twisted torus X , then $\sum_{p \in \delta G} K(p) = 12\chi(G)$.*

Proof. If the region is simply connected, the same proof as before applies: the domain can be trimmed down to a point. If G is equal to the background torus




FIGURE 16. The left picture shows a twisted flat torus X where opposite sides of a rectangular domain in a hexagonal lattice are identified and the total curvature is zero. On the right, the interior of a region G in X is shown. It has two connected components. One is simply connected with Euler characteristic 1. The closed loop has Euler characteristic 0.

X itself, the formula holds also because there is no boundary and the Euler characteristic is 0. Removing a simply connected component of G reduces the Euler characteristic by 1. Filling a hole adds 1 to the Euler characteristic. After filling up all holes, all connected components of G can be edged and trimmed to connected domains G_i for which each $\text{int}(G_i)$ is either a one dimensional closed graph winding around the torus or trimmed to a point. The Euler characteristic of a domain G_i is 0 if a trimmed down version is a closed loop or 1 if it is a point. For every connected component G_i , the Euler characteristic is therefore the same as the sum of the boundary curvatures of G_i . Summation over all components leads to the result. \square

7. COMBINATORIAL CURVATURE

In this section, we consider a more elementary Gauss-Bonnet formula. The curvature is again defined by a Puiseux discretization. Since only circles of length 1 appear in the definition, it is a first order curvature.

Definition. For a two dimensional graph with boundary, the **combinatorial Puiseux curvature** is defined as $K_1(g) = 6 - |S_1(g)|$ for interior points and $K_1(g) = 3 - |S_1(g)|$ for boundary points.

For this combinatorial curvature, Gauss-Bonnet is easier. For subgraphs G of the hexagonal lattice, the boundary curvature is almost trivially equal to 6 because the curvature is related to angles of the corresponding polygon: for a boundary point, $K_1(p) \cdot \pi/3$ is the complementary interior angle at a vertex p of the polygon. Because $\sum_p K_1(p)\pi/3 = 2\pi$ by the polygonal version of the Hopf Umlaufsatz, we have $\sum_p K_1(p) = 6\chi(G)$.

It can be generalized for two dimensional graphs and more generally to two dimensional graphs with boundary. Since it is so closely related to the Euler characteristic, we should attribute it to Euler, even so we are not aware that Euler considered $K_1(g)$, nor that he looked at the dimension of a graph.

Definition. A graph G is called a **one dimensional graph without boundary** if every sphere $S_1(p)$ in G is a zero dimensional graph with 2 points. A graph G is called a **one dimensional graph with boundary** if every unit sphere $S_1(p)$ is a zero dimensional graph with 1 or 2 points. A point in a two dimensional graph is an **interior point** if the unit sphere is a polygon, a one dimensional graph without boundary. If a two dimensional graph has only interior points it is called a **two dimensional graph without boundary**. A point p in a two dimensional graph is a **boundary point** if it is adjacent to an interior point and the unit sphere $S_1(p)$ is a connected one dimensional graph with boundary. A graph is called a **two dimensional graph with boundary** if every point is either an interior point or a boundary point.

With these definitions, the boundary of a two dimensional graph G with boundary is a one dimensional graph without boundary. The boundary of G is made of finitely many subgraphs $G_i = (V_i = \{g_{i1}, \dots, g_{in(i)}\}, E_i)$ with $i = 1, \dots, n$ of $G = (V, E)$ which are polygons and $n_i > 3$. The **stellated graph** H obtained by adding n vertices p_i and edges $(p_i, g_{ij})_{i=1, \dots, n, j=1 \dots n_i}$. The graph H is a two dimensional graph without boundary.

Definition. The Euler characteristic of a two dimensional graph G with boundary is defined as the Euler characteristic of the triangulated two dimensional graph H without boundary.

Remarks. The Euler characteristic is equal to $v - e + f + n$ where $v = |V|$, $e = |E|$ and f is the number of triangles of the original graph G and where n is the difference between the number of vertices of H and the number of vertices of G . The n polygons of the original graph can be thought of as additional faces of the old graph G .

Theorem 8 (Combinatorial Gauss-Bonnet). *For every two dimensional graph with boundary, $\sum_{g \in G} K_1(g) = 6\chi(G)$.*

Remarks. a) This result does not need the rigid requirements on the domain. The graph does not have to be part of X .

b) The result appears in a different formulation: the Princeton Companion to Mathematics [3] mentions on page 832 the formula $\sum_n (6 - n)f_n = 12$ for convex polyhedra, where f_n is the number n -hedral faces and the summation is over all face cardinalities which appear. This is an equivalent formulation but the Gauss-Bonnet character is less evident. The term polyhedron is often used in a much more restrictive way. In [2] the number $C(g) = d(g) - 6$ is called the **charge** of the graph at the point g and $\sum_g C(g) = -12$ mentioned for planar graphs. For two dimensional graphs without boundary, we have $d(g) = |S_1(g)|$ and so $C(g) = -K_1(g)$.

Theorem 8 is **graph theoretical**. Most definitions of polyhedra assume an ambient Euclidean space or more structure on a graph. The book [5] discusses some of the struggles to define polyhedra. Since two dimensional graphs with

boundary behave like polyhedra, one could more generally define a **polyhedron** as a graph in which a sequence of truncation and stellation operation produces a two dimensional graph H without boundary. All Platonic solids are polyhedra in this sense and Gauss-Bonnet holds for H . For a cube for example, which is a zero dimensional graph G in our sense, a truncation of the 8 vertices and additional stellation produces a two dimensional graph H without boundary. The stellation has produced 6 new vertices of curvature -2 . The 24 other points have curvature 1. The sum over all curvatures is $6(-2) + 24 = 12 = 6\chi(H)$. Also for a dodecahedron, a stellation produces 12 new vertices of curvature 1, while the original vertices have curvature 0 after stellation. The octahedron and the icosahedron are already two dimensional with constant curvature 2 and 1 respectively. The tetrahedron is a three dimensional graph. It becomes a two dimensional graph H without boundary after truncating its 4 vertices and then a stellation of the 6 hexagonal boundary curves. The 12 vertices have constant curvature 1, the 6 added face vertices have curvature 0.

Proof. Assume first that $G = (V, E)$ is a two dimensional graph without boundary. Every unit sphere $S_1(p)$ in G is then a polygon. The number of faces f and the number of edges e for the entire graph are then related by the **dimensionality formula**

$$3f = 2e .$$

Additionally, for any graph without self loops and without multiple edges, we have the **edge formula**

$$\sum_{g \in V} |S_1(g)| = 2e .$$

The definition of the Euler characteristic combined with these two formulas gives

$$\begin{aligned} 6\chi(G) &= 6f - 6e + 6v = -2e + 6v \\ &= -\left(\sum_{g \in G} |S_1(g)|\right) + 6v = -\sum_{g \in G} (|S_1(g)| - 6) = \sum_{g \in G} K_1(g) . \end{aligned}$$

The case of a graph G with boundary can be reduced to the boundary-less case: the boundary of G is a union of n closed cycles G_i . For each of them we add an other point P_i and add $n_i = |G_i|$ edge connections from each of the cycle boundary points of G_i to P_i . This produces a stellated graph H without boundary and which contains G as a subgraph. The formula without boundary shows that $6\chi(H)$ is the sum of curvatures of the original interior points and the sum of the curvatures of the boundary points as well as the sum of the curvatures $K_1(P_i) = 6 - n_i$ to each newly added point P_i :

$$6\chi(H) = \sum_{g \in \text{int}(G)} K(g) + \sum_{g \in \delta G} K_H(g) + \sum_{i=1}^n (6 - n_i) .$$

Since we have added n vertices, $|\delta G|$ edges and $|\delta G|$ faces so that

$$(1) \quad \chi(H) = \chi(G) + n - |\delta G| + |\delta G| = \chi(G) + n .$$

The boundary points g satisfy $K_H(g) = 6 - |S_1(g)| - 2$ and $K_G(g) = 3 - |S_1(g)|$ so that $K_H(g) - K_G(g) = 1$. In the stellated graph H each boundary point g has become an interior points for which the curvature $K_h(g)$ is by 1 larger than the




FIGURE 17. The left picture shows a two dimensional graph G with 12 points of curvature -4 and 30 points of curvature 2 leading to a total curvature of 12 . To the right, we see a graph H with boundary obtained by removing all vertices of curvature -4 and their connections. Each of the 120 boundary points which are now present has curvature $-1 = 3 - |S_1|$. The total curvature of H is $30 \cdot 2 - 120 = -60$ matching $6\chi(H)$ since $\chi(H) = 2 - 12 = -10$.

curvature $K_G(g)$ as a boundary point. From the previous boundary less case, we get

$$\begin{aligned}
 6\chi(H) &= \sum_{g \in H} K_H(g) \\
 &= \sum_{g \in \text{int}(G)} K_G(g) + \sum_{i=1}^n \sum_{g \in G_i} (K_G(g) + 1) + \sum_{i=1}^n (6 - n_i) \\
 &= \sum_{g \in G} K(g) + 6n.
 \end{aligned}$$

Combining this with formula (1) gives

$$6\chi(G) = 6\chi(H) - 6n = \sum_{g \in G} K(g) + 6n - 6n = \sum_{g \in G} K(g).$$

□

8. ABOUT HIGHER ORDER CURVATURES

The curvature definition K is motivated by the notion of **Jacobi fields** [1]. While smoothness requirements are necessary for the more sophisticated second order Gauss-Bonnet formula, the metric Gauss-Bonnet formula stated in Theorem 8 holds for any polyhedron with triangular faces. For example, every finite triangularization of a two dimensional compact manifold works. The curvature $K = 2|S_1| - |S_2|$ more differential geometric because the **Gauss-Jacobi equations** $f'' = -Kf$ in differential geometry require the second differences of a Jacobi field f .

We initially were interested in extending the notion of Jacobi fields to the discrete for numerical purposes: for a discretized Jacobi field f with $f(0) = 0$, we have $f''(0) = f(2) - 2f(1) + f(0) = f(2) - 2f(1)$. The Jacobi equations suggest to call this $-K$. Since $f(k)$ is the variation of the geodesic when changing the angle, we can integrate over the circle and we get the arc length $|S_r|$ of the circle of radius $r = 1$ or $r = 2$. Therefore K is a multiple of $2|S_1| - |S_2|$ and there is no reason to normalize this in the discrete. The first order curvature $K_1 = 6 - |S_1|$ on the other hand only requires first order differences. The curvature K has some advantages over the curvature K_1 : **1)** The curvature formula for the boundary and in the interior is the same, while for the curvature K_1 , one has to distinguish boundary and interior. **2)** There is no reference to a flat background structure for K , while K_1 refers to the flat situation with via integers six or three. **3)** It is more closely rooted to the differential geometry of manifolds and to classical notions like Jacobi fields, a notion which is of second order too. **4)** It can be adapted to higher dimension, when defining scalar curvature for graphs and where no natural flat triangulated ambient reference graph exists. While K_1 is a metric flavor, K is more differential geometric. Similarly as many metric results extend to the differential geometric setup, statement become more subtle in the discrete if higher order difference notions are used. The limitations of the results are related to similar limitations we know in the continuum.

We can combine the two results: for the Puiseux curvature with radius 2 defined by $K_2(g) = 12 - |S_2(g)|$ in the interior and $K_2(g) = 6 - |S_2(g)|$ at the boundary, we get the following corollary:

Corollary 9. *If G is a two dimensional smooth domain with boundary C in the triangular tessellation X of the plane, then $\sum_{g \in C} K_2(g) = 24\chi(G)$.*

Proof. Add $\sum_{g \in C} 6 - 2|S_1(g)| = 12\chi(G)$ from Theorem 8 with $\sum_{g \in C} 2|S_1(G)| - |S_2(g)| = 12\chi(G)$ from Theorem 6 to get $\sum_{g \in C} K_2(g) = \sum_{g \in C} 6 - |S_2(g)| = 24\chi(G)$. \square

For example, if G is a disc of radius 1, then $|S_1| = |S_2| = 2$ for all the 6 boundary points g , where $K_1(g) = 2$ and $K_2(g) = 4$.

Unlike the combinatorial curvature formula $\sum_{g \in G} K_1(g) = 6\chi(G)$, the K_2 formula is only obvious modulo the main result Theorem 6 for smooth domains.

REFERENCES

- [1] M. Berger. *A Panoramic View of Riemannian Geometry*. Springer Verlag, Berlin, 2003.
- [2] J. Bondy and U. Murty. *Graph theory*, volume 244 of *Graduate Texts in Mathematics*. Springer, New York, 2008.
- [3] T. Gowers (Editor). *The Princeton Companion to Mathematics*. Princeton University Press, 2008.
- [4] H. Hopf. Über die Drehung der Tangenten und Sehnen ebener Curven. *Compositio Math.*, 2, pages 50–62, 1935.
- [5] I. Lakatos. *Proofs and Refutations*. Cambridge University Press, 1976.