

## ON THE DYNAMICS OF A GENERAL UNITARY OPERATOR

OLIVER KNILL

Department of Mathematics, University of Arizona, Tucson, AZ 85721, USA  
(e-mail: knill@ame2.math.arizona.edu)

(Received February 12, 1997 — Revised June 12, 1997)

A unitary operator  $U$  acting on the weakly compact unit ball  $B$  of a Hilbert space  $H$  gives a topological dynamical system  $(B, U)$ . We prove in this note that for every ergodic  $U$ -invariant measure  $m$ , the dynamical system  $(B, U, m)$  is conjugated to a strictly ergodic group translation. Especially, the topological entropy of  $(B, U)$  is zero. As an application, it follows that any flow on the unit ball given by a Schrödinger equation  $i\hbar\psi = L\psi$  has zero topological entropy.

### 1. Introduction

We consider here the problem to find the set of invariant measures of a linear unitary flow  $U_t$  on a Hilbert space  $H$ . Important examples of such dynamical systems are the Schrödinger flow  $U_t\psi = e^{itL}\psi$  defined by  $i\psi = L\psi$  or the Koopman flow  $f_t(x) = U_t f = f(T_t x)$  on  $L^2(X, m)$  associated to an invariant measure  $m$  of a differential equation  $\dot{x} = f(x)$  on a manifold  $X$ . As a motivation for the problem one can say that the existence of invariant measures for  $U_t$  allows to make statements about recurrence of a wave  $\psi$  in quantum mechanics or a measurable set  $f = 1_A$  in classical mechanics. Examples in both settings show that a linear unitary dynamical system is in general non trivial.

In quantum mechanics, an orbit  $\psi_t$  of the time evolution can be observed by measuring the probabilities  $|\langle \phi, U_t \psi \rangle|^2$  that the wave  $\psi$  is in the pure state  $\phi$  at time  $t$ . For a differential equation with invariant measure  $m$ , one can observe  $m(A \cap T_t(C)) = (1_A, U_t 1_C)$ , the probability with which an orbit  $T_t(x) = x_t$  starting in the set  $C$  hits another set  $A$  at time  $t$ . It is therefore in both cases natural to look for invariant Borel measures on  $H$  with respect to the weak topology. The set of invariant measures of the flow is contained in the set of invariant measures of a time one map so that one can consider the general problem, to determine the set of invariant measures of the topological dynamical system  $(B, U)$ , where  $U$  is an arbitrary unitary operator and  $B$  is the weakly compact unit ball. We are not aware that this problem has been addressed somewhere already. In the two cases, when  $U$  has only absolutely continuous spectrum (i.e. scattering states in quantum mechanics, the mixing property in classical mechanics) or discrete spectrum (i.e. bound states in quantum mechanics, group translation in classical mechanics), the answer to the problem is quite easy.

However, both discrete and absolutely continuous spectrum are exceptional in a Baire category sense: a generic unitary operator  $U$  or a generic Koopman operator  $U_T$  has purely singular continuous spectrum [3, 9, 13, 14]. Singular continuous spectrum appears in the quantum mechanics of solid state physics [5], especially when the potential takes only finitely many values (see for example [7]). In classical mechanics, it appears, whenever a topological shift is embedded because many shift invariant measures have purely singular continuous spectrum [9].

## 2. Topological dynamics of a unitary operator

Given is a unitary operator  $U$  on a separable Hilbert space  $H$ . The unit ball  $B \subset H$  is compact in the weak topology and  $U : B \rightarrow B$  is a homeomorphism. For  $\phi \in H$ , let  $H_\phi$  be the cyclic subspace of  $\phi$  and let  $B_\phi = \{|\psi| \leq 1\}$  be the unit ball in  $H_\phi$ . For  $\phi, \psi \in H$ , the complex measure  $\mu = \mu_{\psi, \phi}$  on  $\mathbf{T} = \{|z| = 1\}$  is defined by the functional  $f \mapsto (\psi, f(U)\phi)$  on  $C(\mathbf{T})$ .

LEMMA 2.1. *The dynamics of  $U$  on  $B_\phi$  is topologically conjugated to a shift on a compact, weakly closed subset in  $l^\infty(\mathbb{Z})$ .*

*Proof:* Consider

$$F : B_\phi \rightarrow l^\infty(\mathbb{Z}, \mathbb{C}), \quad \psi \mapsto \{(\phi, U^n \psi)\}_{n \in \mathbb{Z}},$$

which maps  $\psi$  to the Fourier series of the complex measure  $\mu_{\phi, \psi}$  on  $\mathbf{T}$ .  $F$  is injective, because if  $(\phi, U^n \psi_1) = (\phi, U^n \psi_2)$  for all  $n$ , then  $(U^{-n} \phi, \psi_1) = (U^{-n} \phi, \psi_2)$  for all  $n$  so that  $\psi_1 = \psi_2$  in  $H_\phi$ . If  $l^\infty(\mathbb{Z})$  has the weak topology, then  $F$  is continuous and  $F$  is therefore a homeomorphism onto its image  $Y_\phi = F(B_\phi)$ . Moreover,  $F$  conjugates  $U : B_\phi \rightarrow B_\phi$  to the shift on  $Y_\phi$ .  $\square$

For  $\phi \in H$ , let us denote by  $\mu_\phi = \mu_{\phi, \phi}$  the spectral measure of  $\phi$ . If  $\mu_\phi$  has only atoms, then  $\phi$  is called discrete. If  $\mu_\phi$  has no atoms, then  $\phi$  is called continuous. If  $\mu_\phi$  is absolutely continuous with respect to Lebesgue measure on  $\mathbf{T}$ , then  $\phi$  is called absolutely continuous. If  $\phi$  is both orthogonal to the discrete vectors and all absolutely continuous vectors, it is called singular continuous. (For references on the spectral theory of unitary operators, see [12, Chapter II] or [4, Appendix], for the topological dynamics and ergodic theory see [6, 15, 11]).

### PROPOSITION 2.2

a) *Assume  $\phi$  is a discrete vector. Then  $U$  restricted to the weak closure of the orbit of  $\phi$  is a strictly ergodic dynamical system, which is topologically conjugated to a group translation on a compact topological group.*

b) *If  $\phi$  is absolutely continuous, then  $U^n \phi \rightarrow 0$ .*

c) *Given  $\phi \in H$ , there exists a dense  $G_\delta$  of unitary operators for which  $\phi$  is singular continuous and such that  $\phi$  is recurrent in the sense that there exists a sequence  $n_k \rightarrow \infty$  such that  $U^{n_k} \phi \rightarrow \phi$ .*

d) *There is a dense set  $\psi$  in the continuous subspace of  $U$ , such that the orbit of  $\psi$  has 0 as an accumulation point.*

*Proof:* a) The Fourier transform of  $\mu_\phi = \sum_j a_j \delta(e^{i\lambda_j})$  is the Bohr almost periodic sequence  $F(\phi) = \hat{\mu}_n \sim \sum_j a_j e^{i\lambda_j n}$ . The translates of the sequence  $F(\phi)$  have a compact closure  $X_\phi \subset Y_\phi \subset l^\infty(\mathbb{Z})$  and the shift  $T$  acts as a group translation on the compact hull  $X_\phi$ . The claim follows from Lemma 2.1.

An alternative proof is obtained by diagonalizing  $U$  so that  $U$  becomes a direct sum of operators  $U_i$  on the complex plane, where all  $U_i$  are acting by multiplication with some  $\lambda_i$ .

b) If  $\mu_\phi$  is absolutely continuous, then  $\mu_{\phi,\psi}$  is absolutely continuous [12]. The Riemann–Lebesgue lemma assures that  $\hat{\mu}_n = (\phi, U^n \psi) \rightarrow 0$ . Because this holds for all  $\psi$ , we know that  $U^n \phi \rightarrow 0$  in the weak topology.

c) A dense  $G_\delta$  of measures is rigid, that is, it has the property that for a subsequence  $\hat{\mu}_{n_k} \rightarrow 1$  [3]. A dense  $G_\delta$  set of operators has all vectors singular continuous [13]. Let  $\mathcal{U}$  be the set of all unitary operators with the strong operator topology. The map  $\mathcal{U} \rightarrow M(\mathbf{T})$  attaching to  $U$  the spectral measure  $\mu_{\psi,U}$  is continuous and surjective. A given vector  $\psi$  is therefore rigid for a generic  $U$  in the sense that  $(\psi, U^{n_k} \psi) \rightarrow 1$  for some sequence  $n_k$ , which implies  $U^{n_k} \psi$  converges to  $\psi$ .

d) A point  $\psi$  is called weakly wandering, if there exists a sequence  $k_i$  such that all vectors  $U^{k_i} \psi$  are orthogonal. Weakly wandering points are dense in the subspace of all continuous vectors by a theorem of Krengel [10].  $\square$

### 3. Ergodic theory of a unitary operator

We look now at ergodic properties of  $(B, U)$ , that is, we determine the  $U$ -invariant measures  $m$  on the compact space  $B$ .

**THEOREM 3.1.** *Given a unitary operator  $U$  on a Hilbert space  $H$  with unit ball  $B$ . Every ergodic  $U$ -invariant measure  $m$  defines a uniquely ergodic system  $(\text{supp}(m), U, m)$  which is topologically conjugated to a group translation.*

*Proof:* (i) Two vectors  $\phi, \psi \in H$  define the complex measure  $\mu = \mu_{\phi,\psi}$ . By Wiener's theorem,

$$\lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n |(\psi, U^k \phi)|^2 = \lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n |\hat{\mu}_k|^2 = \sum_{x \in \mathbf{T}} |\mu(\{x\})|^2.$$

(Wiener's theorem is usually formulated for real probability measures on  $\mathbf{T}$  (see [8]) but holds for any complex measures on  $\mathbf{T}$ .) Especially, the right-hand side is zero, for  $\phi, \psi$  in the continuous subspace  $H_c$  because then,  $\mu$  is a continuous measure ( $\mu = \mu_{\phi,\psi}$  is absolutely continuous to both  $\mu_\psi$  and  $\mu_\phi$  [12]).

(ii) For fixed  $\psi$ , the map  $f : B \rightarrow \mathbb{R}$ ,  $f(\phi) = |(\psi, \phi)|^2$  is continuous. Let  $m$  be a  $U$ -invariant measure on  $B$ . Then  $f \in L^1(B, m)$ . By Birkhoff's ergodic theorem,

$$\lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n |(\psi, U^k \phi)|^2 = \lim_{n \rightarrow \infty} n^{-1} \sum_{k=1}^n f(U^k \phi) \rightarrow \int_B f(\phi) dm(\phi) = \int_B |(\psi, \phi)|^2 dm(\phi).$$

(iii) Let  $H = H_d \oplus H_c$  be the orthogonal decomposition of  $H$  into the discrete and continuous subspace with respect to  $U$ . Every  $U$ -invariant measure  $m$  defines a  $U$ -invariant measure  $m_c$  on  $H_c$  by push-forward  $m_c(A) = m(P_c^{-1}A)$ , where  $P_c : H \rightarrow H_c$  is the projection. Let  $B_c$  be the unit ball  $H_c$ . By (i) and (ii),  $\int_{B_c} |(\psi, \phi)|^2 dm_c(\phi) = 0$  for all  $\psi \in H_c$ . This implies that  $m$ -almost all vectors  $\phi \in H_c$  are all orthogonal to the given vector  $\psi \in H_c$ . Taking a countable dense set  $\{\psi_n\}_{n \in \mathbb{N}}$  in  $B_c$ ,  $m$ -almost all vectors  $\phi$  are orthogonal to all these vectors and so to every vector in  $B_c$ . Therefore, the support of  $m_c$  is contained in  $\{0\} \subset B_c$  or  $m_c = 0$ . Especially, every invariant measure  $m$  has its support on the subspace  $H_d$  and the claim follows from Proposition 2.2 a.  $\square$

*Remark:* It follows that  $U$  has continuous spectrum, if and only if every  $U$ -invariant measure  $m$  is located on  $\{0\}$ . Because a dense set of the support  $\text{supp}(\mu)$  of  $m$  is on the sphere, every ergodic  $U$ -invariant measure has its support contained in a sphere  $S_r = \{|\phi| = r\}$ .

**COROLLARY 3.2.** *The topological entropy of  $U : B \rightarrow B$  is zero.*

*Proof:* The metric entropy of every invariant measure  $m$  is zero. By Goodman's variational theorem in ergodic theory (see [15]), the topological entropy is the supremum over all metric entropies and is vanishing too.  $\square$

*Remark:* While  $U$  is an isometry with respect to the norm (for which  $B$  is not compact), it is not an isometry in the weak topology and it is not a priori clear that the topological entropy of  $U$  is vanishing.

#### 4. Additional remarks and questions

A measure on  $B$  which is invariant under a  $\mathbb{R}$  group action  $t \mapsto U_t$ , is also invariant under the  $\mathbb{Z}$  action  $n \mapsto U_1^n$  and Theorem 3.1 shows that  $U_t$  is measure theoretically conjugated to a flow on a compact topological group. Especially "there is no chaos for the Schrödinger flow": for every measure  $m$  on  $B$  which is invariant under the Schrödinger equation  $ih\psi = \Delta\psi + V\psi$ , the measure theoretical flow  $(B, U_t)$  is measure theoretically conjugated to a group translation and the flow has zero topological entropy.

Unitary operators are not interesting for the invariant subspace problem because there are always plenty of invariant subspaces given as images of spectral projections. However, it is interesting to ask for the structure of invariant measures of  $(H, A)$ , if  $A$  is a general linear operator and  $H$  is equipped with the weak topology. In the compactification  $H \cup \{\infty\}$ , there must be some invariant measures. Do still all nontrivial invariant measures have their support on eigenspaces belonging to eigenvalues on the unit circle? The nontriviality of such dynamical systems is illustrated by the open question, whether there exists a system  $(H, A)$  for which every nonzero vector  $\psi$  has a dense orbit. (A positive answer would of course solve the invariant subspace problem). Rolevicz constructed a system  $(H, A)$ , which is transitive in the norm topology of  $H$  [2].

In quantum mechanics, one is also interested in the dynamics  $A \mapsto UAU^*$ , induced on the space of bounded operators. Because  $V : A \mapsto UAU^*$  is a unitary operator on

the Hilbert space of Hilbert–Schmidt operators  $\mathcal{B}_2$  (see [1]), Theorem 3.1 implies that for every invariant measure  $m$  on  $\mathcal{B}_2$ , the dynamical system  $(\text{sup}(m), V)$  is topologically conjugated to a group translation. If  $U$  has continuous spectrum, then  $V$  has also continuous spectrum [1]. It is also evident that if  $U$  has some point spectrum, then  $V$  contains the same point spectrum. We don't know however whether the inner automorphism  $V : A \mapsto UAU^*$  acting on the weakly compact unit ball of the algebra of all operators has zero topological entropy.

## REFERENCES

- [1] W. O. Amrein: *Non-Relativistic Quantum Dynamics*, Reidel Publishing Company, Dordrecht 1981.
- [2] B. Beauzamy: *Introduction to Operator Theory and Invariant Subspaces*, North-Holland, Amsterdam 1988.
- [3] J. R. Choksi and M. G. Nadkarni: *Invariant Subspaces and Allied Topics*, Narosa Publ. Co., New Delhi 1990.
- [4] I. P. Cornfeld, S. V. Fomin and Ya. G. Sinai: *Ergodic Theory*, Springer, Berlin 1982.
- [5] H. L. Cycon, R. G. Froese, W. Kirsch and B. Simon: *Schrödinger Operators—with Application to Quantum Mechanics and Global Geometry*, Springer, 1987.
- [6] M. Denker, C. Grillenberger and K. Sigmund: *Ergodic Theory on Compact Spaces*, Lecture Notes in Mathematics **527**, Springer, 1976.
- [7] A. Hof, O. Knill and B. Simon: *Commun. Math. Phys.* **174** (1995), 149.
- [8] Y. Katznelson: *An Introduction to Harmonic Analysis*, John Wiley and Sons, New York 1968.
- [9] O. Knill: *Discrete and Continuous Dynamical Systems*, in press (1997).
- [10] U. Krengel: *Ergodic Theorems*, Walter de Gruyter, Berlin 1985.
- [11] K. Petersen: *Ergodic Theory*, Cambridge University Press, Cambridge 1983.
- [12] M. Queffélec: *Substitution Dynamical Systems—Spectral Analysis*, Lecture Notes in Mathematics **1294**, Springer, 1987.
- [13] B. Simon: *Ann. Math.* **141** (1995), 131.
- [14] B. Simon: *CRM Proceedings and Lecture Notes* **8** (1995), 109.
- [15] P. Walters: *An Introduction to Ergodic Theory*. Graduate Texts in Mathematics **79**, Springer, New York 1982.