

LECTURE 30

FUNDAMENTAL THEOREM CALCULUS

Oliver Knill, Harvard University

November 18, 2020

PLAN

1. Poll

2. The theorem

3. Anti derivatives

4. Lesson problems

5. More problems

6. Jam: Ice Cream Problem

POLL

Which
of the
formulas
is
not
correct?

A $\frac{d}{dx} \int_0^x f(t) dt = f(x)$

B $\int_0^x f'(t) dt = f(x)$


C $\int_0^x f'(t) dt = f(x) - f(0)$

THE THEOREM

$$\frac{d}{dx} \int_a^x f(t) dt = f(x)$$

Proof:

$$\frac{F(x+h) - F(x)}{h} \sim f(x)$$



AN OTHER VERSION


$$\int_a^b f'(t) dt = f(b) - f(a)$$

Total sum of changes add up to
net change

IN WORDS

The anti derivative of the derivative is the function up to a constant

The derivative of the anti-derivative is the function.




EXAMPLES

A $\int_0^{\pi} \sin(t) dt$


B $\int_0^2 t^3 dt$

C $\int_1^3 \frac{1}{x} dx$

HISTORY



Integration




Differentiation


HISTORY II



Leibniz



Newton



Story of Maths, 2008, BBC, Marcus du Sautoy

HISTORY

NEWTON, 1666

I'VE INVENTED
CALCULUS!



LEIBNIZ, 1674

I'VE INVENT
CALCUL



ISSAAC NEWTON



LESSON PROBLEM 1

Evaluate or
or say "not
enough info"

We know: $\int_2^5 f(t) dt = 7$

A $\int_2^5 f(t) + 4 dt$

B $\int_2^5 f(t + 4) dt$

C $\int_5^2 4f(t) dt$

D $\int_2^4 f(t) + t dt + \int_4^5 f(t) dt$

LESSON PROBLEM 2

$$G(x) = \int_0^x 2t \, dt \qquad F(x) = \int_3^x 2t \, dt$$

A $G(2), G(-2), F(1), F(-2)$

B $G(x)$ for positive x

C $G(x)$ for negative x

D $F(x)$ for $x \geq 3$, and for $x < 3$

E

Graph
F,G

F

Differentiate
F,G

JAM

A

$$\int_0^1 t^3 + t^5 dt$$

B

$$\int_0^2 e^t dt$$

C

$$\int_{-1}^1 \frac{1}{1+x^2} dx$$

CA JAM OF BLUE BELL

$$F(x) = \int_3^x f(t) dt$$

At the Blue Bell ice cream factory, milk is stored in a 40,000 gallon tank. Let $f(t)$ be the rate (in gallons per hour) of milk flowing into and out of the tank one day at time t , where t is measured in hours after noon. At 3 pm, there are 15,000 gallons of milk in the tank.

(a) Define $F(x) = \int_3^x f(t) dt$. What does $F(7)$ represent?

(b) How much milk is in the tank at 7 pm?

(c) How much milk is in the tank x hours after noon?

(d) Define $G(x) = \int_0^x f(t) dt$. What does $G(7)$ represent?

(e) What does $G(x)$ represent?

(f) Let $M(x)$ be the amount of milk in the tank x hours after noon. How would you express “the instantaneous rate at which milk is flowing into/out of the tank at 6 pm” in terms of M ? In terms of f ? In terms of F ?

$f(t)$ gallons/hour
milk flowing
into tank



THE END