

LECTURE 34

WRAPPING UP

Oliver Knill, Harvard University

December 2, 2020

PLAN

1. Poll

2. How to review

2. Motion, Continuity, Functions, Music

3. Extrema, Catastrophes

4. Data

5. Statistics, Infinite integrals

6. Beyond Calculus

POLL

What was the best
topic in this course
for you?

A

Continuity

B

Extrema

C

Limits

D

Related Rates

E

Differentiation

F

Integration

POLL

What was the most
beautiful formula
for you?

$$\int f(u(x))u'(x) \, dx = F(u(x)) + C$$

SUBSTITUTION

$$\frac{d}{dx} \int_a^x f(t) \, dt = f(x)$$

THE FTC

$$\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^{xn} = e^x$$

EXPONENTIAL

$$\lim_{x \rightarrow a} \frac{f(x)}{g(x)} = \lim_{x \rightarrow a} \frac{f'(x)}{g'(x)}$$

HOSPITAL

$$(f(g))' = f'(g)g'$$

CHAIN

HOW TO REVIEW

IN 10 MINUTES




50:00

©Thinkwell Corp.

IN 400 SECONDS




IN 5 SECONDS



IN 5 SECONDS

THE
LION KING

IN 20 SECONDS



MATH 1A		Oliver		
<u>Lecture 1: Hello</u>		knill@math.harvard.edu		
① Hello		McKayley	JAM Section mckayley.green@college.	
② Technology (pull)	40			
③ Netiquette (jam)	Min 10	Jaxon	10:30 Section jaxson.hill@college	
④ Modelling (Slides)	5			
⑤ Bouncing ball (jam) (pull)	10	www.math.harvard.edu/~knill		
⑥ Coffee cooling (jam)	35 min	Section page		
⑦ Questions and Answers		Functions of the day 		
⑧ Deadlines				

SHORT SUMMARIES

Twitter summary:

Calculus links the rate of change of a function with the area under the curve. The derivative F' of the integral F is equal to f .

DIFFERENT ANGLES



MINDMAP



DANCING FUNCTIONS

Beautiful Dance Moves



$\sin(x)$



$\cos(x)$



$\tan(x)$



$\cot(x)$



$|x|$



x



x^2



x^2+y^2



$\sqrt{x}$



$\sqrt{-x}$




$\frac{1}{x}$



crap.

FROM DRAW TO PLAY


```
Play[Sin[5 x]^2 Sin[5200 x], {x, 0, 2}]
```



2 s | 8000 Hz

DISCONTINUITIES

```
Play[ArcTan[Tan[5 x]], {x, 0, 2}]
```



Discontinuities
do not sound nice!

EXPLORE FUNCTIONS

$$f(x) = \sin(10000 \sqrt{x})$$

LARGER SCALE



$T = \{1, 2, 4, 8, 16\};$

$f[x_] := \text{SoundNote}[x, T[\text{Mod}[x, 5] + 1]/10, \text{"Oboe"}];$

$A = \text{Table}[\text{Floor}[13 * \text{Sin}[k/2]], \{k, 20\}];$


$\text{Sound}[\text{Map}[f, A], \{0, 8\}]$

MELODIES

```
f = {1, 1, 3, 3, 5, 5, 1, 1, 1, 1, 3, 3, 5, 5, 1, 1, 5, 5, 6, 6, 8, 8, 8, 8, 5, 5, 6, 6,  
8, 8, 8, 8, 8, 10, 8, 6, 5, 5, 1, 1, 8, 10, 8, 6, 5, 5, 1, 1, 3, 3, -4, -4, 1, 1, 1, 1};  
A = Map[SoundNote, f];  
s1 = Sound[{"Piano", A}, {0, 24}];  
s2 = Sound[{"Flute", RotateRight[A, 8]}, {0, 24}];  
s3 = Sound[{"Guitar", RotateRight[A, 16]}, {0, 24}];  
Sound[{s1, s2, s3}, {0, 12}]
```

1, 1, 3, 3, 5, 5, 1, 1

Bro-ther-Ja-cob



SONGS

```
rise[x_] := Table[{x[[k, 1]] + 12, x[[k, 2]]}, {k, Length[x]};
SN[{x_, t_}] := SoundNote[x, t]; T = 6;
down[x_] := Table[{x[[k, 1]] - 12, x[[k, 2]]}, {k, Length[x]};
d = {1, 5, 8}; g = {1, 6, 10}; a = {0, 3, 8}; h = {-2, 1, 5}; e = {3, 6, 10};
a1 = {{{1}, 4}, {{8}, 4}, {{10}, 1}, {{8}, 1}, {{6}, 2}, {{8}, 4}, {{6}, 1}, {{5}, 1}, {{3}, 2}, {{5}, 2}, {{3}, 1}, {{1}, 1}, {{3}, 3}, {{}, 1}, {{3}, 4}};
b1 = {{{1}, 4}, {{1}, 4}, {{3}, 4}, {{-4}, 4}, {{1}, 2}, {{0}, 1}, {{-2}, 1}, {{-2}, 2}, {{-4}, 1}, {{-6}, 1}, {{-4}, 3}, {{}, 1}, {{-4}, 4}};
a2 = {{d, 4}, {d, 4}, {g, 4}, {d, 4}, {a, 4}, {d, 4}, {a, 4}, {a, 4}};
b2 = {{g, 4}, {h, 4}, {e, 4}, {a, 4}, {d, 4}, {g, 4}, {a, 4}, {a, 4}};
s1 = Sound["Flute", SN /@ rise[a1]], {0 T, 1 T}];
s3 = Sound["Viola", SN /@ rise[a1]], {1 T, 2 T}];
s5 = Sound["Violin", SN /@ rise[b1]], {2 T, 3 T}];
s7 = Sound["Viola", SN /@ rise[b1]], {3 T, 4 T}];
S = Sound[{s1, s2, s3, s4, s5, s6, s7, s8}]

s2 = Sound["Guitar", SN /@ down[a2]], {0 T, 1 T}];
s4 = Sound["Harp", SN /@ down[a2]], {1 T, 2 T}];
s6 = Sound["Piano", SN /@ down[b2]], {2 T, 3 T}];
s8 = Sound["Cello", SN /@ down[b2]], {3 T, 4 T}];
```



SONGS

With
random
components.

```
(* Oliver Knill, May 5, 2020 *)
R[x_] := RandomChoice[Permutations[x]];
f[{x_, t_}] := Module[{a=R[x]}, Table[{a[[k]]}, t/3], {k, 3}];
rise[x_] := Table[{x[[k, 1]]+12, x[[k, 2]]}, {k, Length[x]}];
down[x_] := Table[{x[[k, 1]]-12, x[[k, 2]]}, {k, Length[x]}];
SN[{x_, t_}] := SoundNote[x, t]; T=3;
d={1, 5, 8, 13}; g={1, 6, 10, 13}; a={0, 3, 8, 12};
d1={1}; e1={3}; g1={6}; f1={5}; k1={8}; g2={18};
d2={13}; h1={10}; c1={12}; e2={15}; k2={20}; f2={17};
a1={{d, 1}, {d, 1}, {g, 2}, {d, 2}, {d, 2}, {a, 4}};
a2={{a, 1}, {a, 1}, {a, 2}, {a, 2}, {a, 2}, {d, 4}};
a3={{d, 1}, {d, 1}, {d, 2}, {d, 2}, {d, 2}, {g, 4}};
a4={{g, 1}, {g, 1}, {d, 2}, {d, 2}, {a, 2}, {d, 4}};
b1={{k1, 1}, {k1, 1}, {h1, 2}, {k1, 2}, {d2, 2}, {c1, 4}};
b2={{k1, 1}, {k1, 1}, {h1, 2}, {k1, 2}, {e2, 2}, {d2, 4}};
b3={{k1, 1}, {k1, 1}, {k2, 2}, {f2, 2}, {d2, 2}, {c1, 2}, {h1, 2}};
b4={{g2, 1}, {g2, 1}, {f2, 2}, {d2, 2}, {e2, 2}, {d2, 4}};
c1=Join[f[{d, 1}], f[{d, 1}], f[{g, 2}], f[{d, 2}], f[{d, 2}], f[{a, 4}]];
c2=Join[f[{a, 1}], f[{a, 1}], f[{a, 2}], f[{a, 2}], f[{a, 2}], f[{d, 4}]];
c3=Join[f[{d, 1}], f[{d, 1}], f[{d, 2}], f[{d, 2}], f[{d, 2}], f[{g, 4}]];
c4=Join[f[{g, 1}], f[{g, 1}], f[{d, 2}], f[{d, 2}], f[{a, 2}], f[{d, 4}]];
s1=Sound[{"Piano", SN/@down[a1]}, {0T, 1T}];
s2=Sound[{"Piano", SN/@down[a2]}, {1T, 2T}];
s3=Sound[{"Piano", SN/@down[a3]}, {2T, 3T}];
s4=Sound[{"Piano", SN/@down[a4]}, {3T, 4T}];
t1=Sound[{"Flute", SN/@rise[b1]}, {0T, 1T}];
t2=Sound[{"Flute", SN/@rise[b2]}, {1T, 2T}];
t3=Sound[{"Flute", SN/@rise[b3]}, {2T, 3T}];
t4=Sound[{"Flute", SN/@rise[b4]}, {3T, 4T}];
u1=Sound[{"Guitar", SN/@rise[c1]}, {0T, 1T}];
u2=Sound[{"Guitar", SN/@rise[c2]}, {1T, 2T}];
u3=Sound[{"Guitar", SN/@rise[c3]}, {2T, 3T}];
u4=Sound[{"Guitar", SN/@rise[c4]}, {3T, 4T}];
S=Sound[{s1, s2, s3, s4, t1, t2, t3, t4, u1, u2, u3, u4}];
Export["birthday.mp3", S, "MP3"]
```


CREATIVITY



```
A = Table[RandomChoice[Range[80] - 40], {30}, {6}];  
f[x_] := SoundNote[x, RandomChoice[{1, 2}]];  
Sound[{"Piano", Map[f, A]}, {0, 6}]
```

TRY IT OUT!



play $\sin(200 t) \sin(10000 t)$

Extended Keyboard

Upload

Examples

Random

Input interpretation:

waveform $\sin(200 t) \sin(10000 t)$


Waveform:

[Play Sound](#)



(audible to the human ear)

Frequency power spectrum:



Download Page

POWERED BY THE WOLFRAM LANGUAGE

MOTION




INTERMEDIATE VALUE


$f(x)$ continuous on $[a,b]$

$f(a), f(b)$ different signs

$\Rightarrow f$ has a root in $[a,b]$



CONTINUITY




CONTINUITY





Experiment of Sunday, May 8, 2011

DERIVATIVES

$f(x)$ = DJI data at day x
from 1918-2020




DERIVATIVES




DJI change
last year, 230
trading days so far

EQUILIBRIA


Basic Principle: A system wants to be in a minimal energy state.




EQUILIBRIUM CHANGE




$$f(x) = x^4 - x^2$$



$$f(x) = x^4 - x^2 - 0.6x$$





EQUILIBRIA IN PSYCHOLOGY




FLIPPING PERCEPTION



PERCEPTION





MATHEMATICAL LINK



CONFLICTS

ECONOMICS



APPROPOS PSYCHOLOGY

While sitting at your desk, lift your right foot off the floor and make clockwise circles. Now, while doing this draw the number "6" in the air with your right hand. Your foot will change direction and there's nothing you can do about it.

DATA TO FUNCTIONS

Data

Histogram

smoothed PDF




Data

Histogram


smoothed PDF



INFINITE INTEGRALS



ALTHOUGH THE NOTATION MAY BE UNFAMILIAR, ALL IT MEANS IS AN AREA.. THE INTEGRAL SIGN ITSELF IS A STRETCHED "S," FOR SUM, WHICH THE INTEGRAL, IN SOME SENSE, IS.



AS A SUMLIKE SOMETHING, THE INTEGRAL SERVES TO DEFINE THE **MEAN AND VARIANCE of a continuous random variable.**

$$\mu = \int_{-\infty}^{\infty} xf(x)dx$$


BY ANALOGY
WITH THE
DISCRETE
FORMULAS:

$$\mu = \sum_{\text{all } x} xp(x)$$

$$\sigma^2 = \int_{-\infty}^{\infty} (x-\mu)^2 f(x)dx$$

$$\sigma^2 = \sum_{\text{all } x} (x-\mu)^2 p(x)$$

ALTHOUGH IT MAY NOT BE OBVIOUS FROM THE FORMULAS, THESE DEFINITIONS OF MEAN AND VARIANCE ARE ENTIRELY CONSISTENT WITH THEIR ROLE AS CENTER AND AVERAGE SPREAD OF THE PROBABILITIES GIVEN BY THE DENSITY $f(x)$. THE PICTURE TO KEEP IN MIND IS THIS:



STATISTICS

$$\int_{-\infty}^{\infty} f(x) \, dx = 1$$

Assumption

$$\int_{-\infty}^{\infty} xf(x) \, dx = m$$

Mean

$$\int_{-\infty}^{\infty} (x - m)^2 f(x) \, dx$$



Variance

$$\sigma(f) = \sqrt{\text{Var}(f)}$$

Standard Deviation

LIE WITH STATS


1954
Darrell Huff
1913-2001



*CALCULUS AND
HUMOR*

How do you make  burn?

Differentiate a log



BEYOND 1A

Math 1B

Series and Diff equations

Math 2Ia


Multivariable Calculus

Math 2Ib

Linear Algebra and Diff Equ

Stats 101

Probability and Statistics



Computer
Science

Physics
Astronomy

Business
Economics

BioChem
LifeScience

Psychology
Philosophy



THE END