

# CALCULUS AND DIFFERENTIAL EQUATIONS

MATH 1B

## Lecture 25: Differential equations, 11/03/2021

### DIFFERENTIAL EQUATIONS

**25.1.** A differential equation is an equation for an unknown function involving derivatives of the function. For example,  $y'(t) = y(t)$  is a differential equation for an unknown function  $y(t)$ . Think of  $t$  as “time”.

**25.2.** Unlike for usual equations like  $3x = 4$ , where we look for a **number** as a solution, we now look for a **function**. A **solution** to the differential equation is a function  $y(t)$  which satisfies the equation. You might notice that there is more than one solution to the equation  $y'(t) = y(t)$ . Can you see the general solution?

**25.3.** We are already dealing with differential equations when integrating: the equation  $y'(t) = t^2$  has the solution  $y(t) = \int_0^t x^2 dx + C$ , where  $C$  is a constant. A **specific solution** to the above equation  $y'(t) = t^2$  is  $t^3/3$ . The **general solution** is  $t^3/3 + C$ .

### GROWTH MODELS

**25.4.** Here is a typical **population growth** problem:

**Example:** If  $M(t)$  is the number of Canadian geese on the Charles river. Each geese has 2 off springs a year in average while 1 geese dies. How many geese are there in 5 years if there are 5000 geese initially?



FIGURE 1. A Canadian goose. Photo by Oliver on August 23, 2020.

We can model this as  $M'(t) = 2M(t) - M(t) = M(t)$ . The solution of  $M(t) = 5000e^t$  gives for  $t = 5$  the number  $5000e^5$ .

## DECAY MODELS

**25.5.** An other situation, where differential equations appear are **decay models**. Here is a typical example

**Example:** The amount  $N(t)$  of Carbon 14 in a sample satisfies

$$N'(t) = -0.0001216N(t)$$

The negative sign means that the number of Carbon 14 isotopes **decay** with time. If we have initially  $N_0$  atoms, then after time  $t$  we have  $M(t) = e^{-0.0001216t}M(0)$ . Since the decay number is  $\log(2)/5700 = -0.0001216$  we know that in 5700 years the amount of Carbon 14 is half. We could also write  $M(t) = e^{-t/5700}M(0)$ .

## BANKING

**25.6.**

**Example:** If  $M(t)$  is the bank account, then under a continuous compounding assumption, the balance  $M'(t)$  is  $rM(t)$ , where  $r = 0.06$  is the **interest rate**. If there are  $M(0) = 100'000$  dollars initially, the equation how much do we have in  $t = 10$  years?

The equation  $M' = rM$  is solved by  $M(t) = 100'000e^{0.06t}$ .

**Example:** If additionally, 50'000 dollars is continuously transferred to the account we can model this with  $M'(t) = M(t) + 50'000$ . Can you see why  $M(t) = 100'000e^{0.06t} + 50000(e^{0.06t} - 1)$  satisfies the differential equation?

Solution: subtract  $M(t)$  from

$$M'(t) = 0.06 * 100'000e^{0.06t} + 0.06 * 50'000e^{0.06t}$$

This gives 50'000.

**25.7.** Compare that if  $r$  is the interest rate and we would compound annually in  $n$  steps, the bank account grows like

$$(1 + r/n)^n .$$

This is a compound interest formula and for  $n \rightarrow \infty$  it approaches  $e^r$ .

**25.8.** Differential equations can be much more general and model much more interesting situations. While  $y' = ry$  models exponential growth for positive  $r$  or exponential decay for negative  $r$ . We will also look at more complicated systems later in this course.

## REMINDERS

- Homework PS 23 is due on Friday.
- There are partial point recovery problems for the midterms coming up.