



Lecture 26

11/05/2021

*Slope
fields*

8/30/2021 near Mather house

Table of Contents

1) Review: $y' = r y - c$ problems. Egg nogg!

2) Slope fields and examples

3) Uniqueness of $y' = F(y, t)$

4) Worksheet

1) Review

A

$y' = r y$ has the general
solution

$$y(t) = C e^{rt}$$

B

$y' = f(t)$ has the general
solution

$$y(t) = F(t) + C$$

$$F' = f$$

C

$y' = r y + a$ has the general
solution

$$y(t) = C e^{rt} - \frac{a}{r}$$

Eggnoy

from worksheet



input : sugar
 4 Gallons/min
 1/2 concentration

**100 Gallon
 tank**

$S(t)$ amount
 of sugar

milk
 cream
 egg whites

output:
 Eggnog
 4 Gallons/min




Sugar concentration

in worksheet:

$$S'(t) = 2 - \frac{S(t)}{25}$$

$$S(0) = 0$$



$$S(t) = 50 - 50e^{-t/25}$$


Eggnog equation

$S' = a - rS$ has the general
solution $y(t) = Ce^{-rt} + \frac{a}{r}$

This is the same as the Banking problem.

We will come back to this next week!


2) Slope Fields



$$y' = r y$$

has solutions

$$y(t) = C e^{rt}$$



$$y' = f(t)$$


has solutions

$$y(t) = F(t) + C$$

if $F' = f$.

Here $f = \sin(t)$.


$$y(t) = -\cos(t) + C$$



logistic equation

$$y' = y(2-y)$$


example of
autonomous
equation



see homework

3) Uniqueness of solutions

If F is
differentiable,
 $y'(t) = F(y,t)$
has a unique
solution $y(t)$,
given the initial
condition
 $y(0) = b$



6) *Worksheet*

The End