



Lecture 28

11/10/2021

*Compartmental
analysis*

8/30/2021 near Mather house

Table of Contents

1) Eggnog review: 2 pictures!


2) Input-Container-Output

3) Example


4) Worksheet

Weather in real time

The US Weather service runs an interesting website that allows you to view weather forecast data at any place in the United States. Here is the url for this website: <https://digital.weather.gov/>. Go to this page and here is what you see:



I am writing this on July 7 in Cambridge MA, so the weather there interests me. I can focus in on the forecasts for that day using the arrows and ± on the left side near the top of the map. You can see the screen shot of the temperature forecasts for 8pm in



Notice that this depicts the *maximum* temperature for the Boston area up until 8pm tonight. It says this under the temperature map--it also says there when the forecast was issued.

Temperatures in the 90's are nice!! Most people don't like it that hot but I do because I save the memory of hot days in a special place for the upcoming grim days of January when I am out with the flying snow, the temperature below zero (Fahrenheit) and a gale blowing long and evil from the northeast. That's when I use up those memories of 90° days.


The weather service is very good at explaining how they produce their graphics. You get this information and the raw data too by clicking on the button under the map that says 'NDFD Data' (see the screen shot below).



Eggnoy



milk
cream
egg whites




input : sugar milk
4 Gallons/min
1/2 concentration



100 Gallon
tank

$S(t)$ amount
of sugar




output:
Eggnog
4 Gallons/min



Slopefield picture (t - S plane)

$$S'(t) = 2 - \frac{S(t)}{25}$$




$$S(t) = 50 - 50e^{-t/25}$$

t

S-S' picture

$$\begin{aligned} S'(t) &= f(S(t)) \\ &= 2 - \frac{S(t)}{25} \end{aligned}$$



The equilibrium is stable because $f'(50) = -1/25$ is negative!

Input-Container-Output

$$y'(t) = \text{input} - \text{output}(y)$$



both input and output depend on concentration. Only output depends on the quantity y .


Worksheet problems

$$M' = 2 - \frac{M}{5} = f(M, t)$$

$$M(0) = 2$$

Draw the slope field picture in the t - M plane

Equilibrium: $M = 10$
Slope field:




Find the solution $M(t)$ satisfying $M(0) = 2$
 $10 + ce^{-t/5}$
 How do we find c ?
 $M(0) = 10 + ce^{-0t} = 2$
 $= 10 + c = 2$
 $c = -8$


Assume: Output changes to 8 l/hour.
How does the diff equ. change?

$$M' = 2 - \frac{M}{20-4t}$$

no more autonomous!



Mango - Cranberry - Orange



- a) $M(0) = 2$
- b) Input: 2
- c) $\frac{\text{Mango in Mixer}}{\text{Total of mix in Mixer}} = \frac{M(t)}{20} = \frac{M(t)}{5}$

d) Output:

$$4 \frac{M(t)}{20} = \frac{M(t)}{5}$$

$M(t)$ = Mango in Liters

$$M'(t) = \text{Input} - \text{Output}$$


$$= 2 - \frac{M(t)}{5}$$

$$M(0) = 2$$

$$y' = a - ry$$

input output

Review: 2 pictures



general solution

$$y(t) = \frac{a}{r} + ce^{-rt}$$

Lecture 28

Reminders:
 • Point recovery due tomorrow.
 • QRD Friday.

Compartmental Analysis
 Input-mixer-output analysis

- 1) Review: $y' = f(y, t)$
 stability, autonomous, $(y-y')$ and $(t-y)$ pictures
- 2) Orange-Cranberry-Mango problem
- 3) General strategy to solve input-output prob.
- 4) Salt water problem (worked)
- 5) **Huuuuhs!** problem.

Reminders

1) Recovery Exam 1 due Tomorrow

2) Homework 26 due Friday

3) QRD project on Weather due Friday

The End