

# DIFFERENTIAL GEOMETRY

MATH 136

## Unit 15: Local Gauss-Bonnet

**15.1.** We now prove the Gauss-Bonnet theorem in the situation when  $U \subset R$  is a polygon. The parametrization  $r : R \rightarrow M$  plants the polygon  $r(U) \subset r(R)$  into the surface  $M$ .




FIGURE 1. The local Gauss-Bonnet theorem tells that face, edge and vertex curvatures of a polygon  $r(U)$  in a manifold  $M$  add up to  $2\pi$ .

**15.2.** A **simple polygon** in  $M$  is the image  $r(U)$  of a simple polygon  $U \subset \mathbb{R}^2$  such that  $r$  is smooth and injective on  $U$ . Its **Euler characteristic** is  $\chi(U) = |V| - |E| + |F| = 3 - 3 + 1 = 1$ . As in the discrete Hopf Umlaufsatz, the vertex curvatures are defined as  $\kappa_i = \pi - \alpha_i$ , where  $\alpha_i$  are the polygon angles. The **angles  $\alpha_i$  of the polygon** are defined by  $\cos(\alpha_i) = \dot{x}_i(1) \cdot \dot{x}_{i+1}(0)$ , the dot product of the velocity vectors of the arcs at the end of the incoming and the beginning of the outgoing arc.

**15.3.** Let  $U$  be a simple polygon on  $M$ . There are three contributions to curvature: the **face curvature** is the integral of  $K$  over the interior, the **geodesic curvature** integrates sectional curvature  $\kappa_g$  over the edges  $C_j$  and then there are the **vertex curvatures**  $\kappa_j = \pi - \alpha_j$  attached to the vertices.

**Theorem 1** (Local Gauss-Bonnet).  $\iint_U K dV + \sum_j \int_{C_j} \kappa_g(x_j(t)) dt + \sum_j \kappa_j = 2\pi$ .

**15.4.** If  $x(t) = r(u(t), v(t))$  parametrizes the boundary of the surface  $M = r(U)$ , we can assume that it is parametrized by arc-length. The velocity vector  $\dot{x}$  is a 3-vector tangent to the surface. We look at the orthonormal frame field  $(z, w)$  from last time. The **geodesic curvature** of a curve  $x$  is defined at points where  $x$  is smooth and given as  $\kappa_g = (n \times \dot{x}) \cdot \ddot{x}$ . Unlike  $\kappa = |\dot{x} \times \ddot{x}|$ , it is signed. So is the **normal curvature**  $\kappa_n = n \cdot \ddot{x}$ . Since  $\dot{x} \cdot \ddot{x} = 0$ , Pythagoras gives  $\kappa_g^2 + \kappa_n^2 = \kappa^2$ . The velocity vector of

the curve can be expressed as an angle so that  $\dot{x} = \cos(\theta)z + \sin(\theta)w$ . We write  $\dot{w}$  for  $\frac{d}{dt}w(x(t))$ .

**Lemma 1** (Geodesic lemma).  $\kappa_g = \dot{\theta} - (z \cdot \dot{w})$ .

*Proof.* Fill in the parts of the definition  $\kappa_g = (n \times \dot{x}) \cdot \ddot{x}$ :

(i)  $n \times \dot{x} = \cos(\theta)w - \sin(\theta)z$ .

(ii)  $\ddot{x} = \dot{\theta}(-\sin(\theta)z + \cos(\theta)w) + \cos(\theta)\dot{z} + \sin(\theta)\dot{w}$ .

(iii) So,  $\kappa_g = (n \times \dot{x}) \cdot \ddot{x} = \dot{\theta} - z \cdot \dot{w}$  □

**15.5.** We can now prove the local Gauss-Bonnet theorem:

*Proof.* (i) Integrating the geodesic lemma gives

$$\int_0^L \kappa_g dt = \int_0^L \dot{x} dt - \int X dr$$

(ii) Green's theorem assures that  $\int X dr = \iint_U K dV$  as  $KdV = dX$ .

(iii) The Hopf Umlaufsatz for curved polygons gives  $\int_0^L \dot{\theta}(t) dt + \sum_j(\pi - \alpha_j) = 2\pi$ .

(iv) Putting (i),(ii),(iii) together gives the proof. □

**15.6. Example 1)** If  $K$  is constant 0 and  $U$  is a triangle, Gauss Bonnet is  $\kappa_1 + \kappa_2 + \kappa_3 = 2\pi$ . This is equivalent to  $\alpha_1 + \alpha_2 + \alpha_3 = \pi$  you know from elementary school geometry. For an **n-gon**, the identity  $\sum_{i=1}^n \kappa_i = 2\pi$  is equivalent to  $\sum_{i=1}^n \alpha_i = (n - 2)\pi$ .

**15.7. Example 2)** If  $M = \mathbb{S}^2$  is a sphere of radius 1, then curvature is  $K = 1$ . The integral  $\iint_U K dV$  is the **area**  $|U|$  **of the triangle**. The formula becomes  $|U| + \sum(\pi - \alpha_i) = 2\pi$  and so  $\alpha_1 + \alpha_2 + \alpha_3 = |r(U)| + \pi$ . This is **Girard's theorem** or **Harriot's theorem** in spherical geometry, named after Albert Girard or Thomas Harriot.




FIGURE 2. A triangle in the plane has  $\alpha_1 + \alpha_2 + \alpha_3 = \pi$ . For a spherical triangle of area  $A$ , Harriot's theorem gives  $\alpha_1 + \alpha_2 + \alpha_3 = \pi + A$ . On a hyperbolic space, Lambert's theorem is  $\alpha_1 + \alpha_2 + \alpha_3 = \pi - A$ .

**15.8. Example 3)** If  $M$  is a surface of constant curvature  $-1$ , a triangle is called **hyperbolic**. Now,  $\iint_U KdV = -|U|$  and  $\alpha_1 + \alpha_2 + \alpha_3 = \pi - |U|$ , a formula found by Johann Heinrich Lambert. The right hand side  $\pi - |U|$  is called **spherical defect**.

**15.9. Example 4)** Take a sphere with a simple closed geodesic on it, integral of  $K$  on each half is  $2\pi$ . The total integral is  $4\pi$ .

**15.10. Example 5)** If  $K = 0$  and  $r(U)$  is a region in the plane bound by a simple smooth curve, we have the **Hopf Umlaufsatz**  $\int \kappa_g(t) dt = 2\pi$ .