

INTRODUCTION TO CALCULUS

MATH 1A

Unit 15: Review

MAJOR POINTS

f is **continuous** at a if there is $b = f(a)$ such that $\lim_{x \rightarrow a} f(x) = b$ for every a . The intermediate value theorem: $f(a) > 0, f(b) < 0$ implies f having a root in (a, b) .

$f'(x) = 0, f''(x) > 0$ then x is **local min**. $f'(x) = 0, f''(x) < 0$ then x is **local max**. For **global minima or maxima**, compare local extrema and boundary values.

If f changes sign we have a **root** $f = 0$, if f' changes sign, we have a **critical point** $f' = 0$ if f'' changes sign, we have an **inflection points**. A function is **even** if $f(-x) = f(x)$, and **odd** if $f(-x) = -f(x)$.

If $f' > 0$ then f is increasing, if $f' < 0$ it is decreasing. If $f''(x) > 0$ it is **concave up**, if $f''(x) < 0$ it is **concave down**. If $f'(x) = 0$ then f has a horizontal tangent.

Hospital's theorem applies for $0/0$ or ∞/∞ situations. In that case, $\lim_{x \rightarrow p} f(x)/g(x)$, where $f(p) = g(p) = 0$ or $f(p) = g(p) = \infty$ with $g'(p) \neq 0$ are given by $f'(p)/g'(p)$.

With $Df(x) = (f(x+h) - f(x))/h$ and $S(x) = h(f(0) + f(2h) + \dots + f((k-1)h))$ we have a **preliminary fundamental theorem of calculus** $SDf(kh) = f(kh) - f(0)$ and $DS(f(kh)) = f(kh)$.

Roots of $f(x)$ with $f(a) < 0, f(b) > 0$ can be obtained by the dissection method by applying the **Newton map** $T(x) = x - f(x)/f'(x)$ again and again.

Algebra reminders

Healing: $(a+b)(a-b) = a^2 - b^2$ or $1 + a + a^2 + a^3 + a^4 = (a^5 - 1)/(a - 1)$
Denominator: $1/a + 1/b = (a+b)/(ab)$
Exponential: $(e^a)^b = e^{ab}$, $e^a e^b = e^{a+b}$, $a^b = e^{b \log(a)}$
Logarithm: $\log(ab) = \log(a) + \log(b)$. $\log(a^b) = b \log(a)$
Trig functions: $\cos^2(x) + \sin^2(x) = 1$, $\sin(2x) = 2 \sin(x) \cos(x)$, $\cos(2x) = \cos^2(x) - \sin^2(x)$
Square roots: $a^{1/2} = \sqrt{a}$, $a^{-1/2} = 1/\sqrt{a}$

Important functions

Polynomials	$x^3 + 2x^2 + 3x + 1$	Exponential	$5e^{3x}$
Rational functions	$(x + 1)/(x^3 + 2x + 1)$	Logarithm	$\log(3x)$
Trig functions	$2 \cos(3x)$	Inverse trig functions	$\arctan(x)$

Important derivatives

$f(x)$	$f'(x)$	$f(x)$	$f'(x)$
$f(x) = x^n$	nx^{n-1}	$f(x) = \sin(ax)$	$a \cos(ax)$
$f(x) = e^{ax}$	ae^{ax}	$f(x) = \tan(x)$	$1/\cos^2(x)$
$f(x) = \cos(ax)$	$-a \sin(ax)$	$f(x) = \log(x)$	$1/x$
$f(x) = \arctan(x)$	$1/(1 + x^2)$	$f(x) = \sqrt{x}$	$1/(2\sqrt{x})$

Differentiation rules

Addition rule	$(f + g)' = f' + g'$	Quotient rule	$(f/g)' = (f'g - fg')/g^2$
Scaling rule	$(cf)' = cf'$	Chain rule	$(f(g(x)))' = f'(g(x))g'(x)$
Product rule	$(fg)' = f'g + fg'$	Easy rule	simplify before deriving

Extremal problems

To maximize or minimize f on an interval $[a, b]$, find all critical points inside the interval, evaluate f on the boundary $f(a), f(b)$ and then compare the values to find the global maximum. Do not make the second derivative test at the boundary.

Limit examples

$\lim_{x \rightarrow 0} \sin(x)/x$	l'Hopital 0/0	$\lim_{x \rightarrow 1} (x^2 - 1)/(x - 1)$	heal
$\lim_{x \rightarrow 0} (1 - \cos(x))/x^2$	l'Hopital 0/0 twice	$\lim_{x \rightarrow \infty} \exp(x)/(1 + \exp(x))$	l'Hopital
$\lim_{x \rightarrow 0} (1/x)/\log(x)$	l'Hopital ∞/∞	$\lim_{x \rightarrow 0} (x + 1)/(x + 5)$	no work necessary

Important things

Summation and rate of change are at the heart of calculus.
 The 3 major types of discontinuities are jump, oscillation, infinity.
 Dissection and Newton methods are algorithms to find roots.
 The fundamental theorem of trigonometry is $\lim_{x \rightarrow 0} \sin(x)/x = 1$.
 The derivative is the limit $Df(x) = [f(x + h) - f(x)]/h$ as $h \rightarrow 0$.
 The rule $D(1 + h)^{x/h} = (1 + h)^{x/h}$ leads to $\exp'(x) = \exp(x)$.
 If you forget a derivative like of $\arctan(x)$, use the chain rule.