

INTRODUCTION TO CALCULUS

MATH 1A

Unit 12: Maxima and Minima

12.1. Pierre Fermat made a simple but profound observation: if $f'(x)$ is not zero, then x can not be a maximum nor be a minimum. His reasoning was: if you make a step h then you end up at $f(x+h) \sim L(x+h) = f(x) + hf'(x)$. Indeed, we all know that if there is a slope and do a step we end up a bit higher.

12.2. Lets call a point x a **local maximum** of f if $f(y) \leq f(x)$ for all y near enough to x . The function $f(x) = x^3 - 2x$ for example has a local minimum at $x = 1$ and a local maximum at $x = -1$. The observation of Fermat is equivalent to:

Fermat's principle: If a differentiable function f has a local maximum or minimum at x , then $f'(x) = 0$.

12.3. The function $f(x) = x^2$ for example has the derivative $f'(x) = 2x$. This is zero at $x = 0$, the minimum of f . Note that the converse of Fermat's statement is not necessarily true: if $f'(x) = 0$, then x does not need to be a maximum or minimum. The standard example is $f(x) = x^3$. We have $f'(x) = 3x^2$ which is zero at $x = 0$. But $x = 0$ is neither a maximum nor minimum of f .

12.4. A point x is called a **critical point** of f , if $f'(x) = 0$. Critical points are important because they are **candidates for maxima and minima**.

12.5. The next test allows to see whether we have a maximum or minimum. The derivative should exists near a but not necessarily at a . Like for $f(x) = |x|$ and $a = 0$.

First derivative test: If a is a critical point of f and the slope $f'(x)$ changes from negative to positive at a then a is a local minimum. If $f'(x)$ changes from positive to negative at a , then a is a local maximum. If $f'(x)$ does not change sign, the a is neither a local maximum nor local minimum.

12.6. Second derivatives help. It assumes that the second derivative exists at a .

Second derivative test: If a is a critical point of f and $f''(a) > 0$, then f is a local minimum. If $f''(a) < 0$, then f is a local maximum. If $f''(a) = 0$, the test is inconclusive.

Homework

This PSet is due Wednesday February 21, 2024.

Problem 12.1: Find the critical points of the following two functions

a) $f(x) = x^4 - 4x^3 + 4x^2$.

b) $f(x) = x^4(x - 3)^2$.

Problem 12.2: Use the second derivative test to determine the nature of the critical points in the same two functions:

a) $f(x) = x^4 - 4x^3 + 4x^2$.

b) $f(x) = x^4(x - 3)^2$.

Problem 12.3: What does the first derivative test tell you about the behavior at the critical points in the two cases

a) $f(x) = x^4 - 4x^3 + 4x^2$.

b) $f(x) = x^4(x - 3)^2$.

In each case, is there a point, where the first derivative test gives more information than the second derivative test?

Problem 12.4: Find all the critical points and determine whether it is a local maximum, a local minimum or neither. You can use either test. You will see that some of the cases are a bit unusual.

a) $f(x) = e^2x - e^x$

b) $f(x) = e^x + x$

c) $f(t) = t^4 + t^3$

d) $f(t) = |2t - 8|$

e) $f(x) = 5$.

Problem 12.5: a) Both the first and second derivative test do not work for the **tamed devil function** $f(x) = x \sin(1/x)$ at $x = 0$. Why not? (Since we have no chain rule yet, you can certainly look up the first and second derivative using a tool like Wolfram alpha.)

b) Function $f(x) = \arcsin(\sin(x))$ has appeared in our ground hog movie. Where are the maxima and minima? To do so, plot the function $f(x)$ and its derivative $f'(x)$ and use one of the derivative tests at the maxima and minima.