

## Lecture 15: Review for first midterm

### Major points

A function is **continuous**, if the closeness of  $x, y$  implies the closeness of  $f(x), f(y)$ . Intermediate value theorem:  $f(a) > 0, f(b) < 0$  implies  $f$  having a root in  $(a, b)$ .

At a local **extremum**,  $f'(x) = 0$ . If  $f''(x) > 0$ , it is a local minimum. If  $f''(x) < 0$ , it is a local maximum. **Global extrema**: compare local extrema and boundary values.

If  $f' > 0$  then  $f$  is increasing, if  $f' < 0$  it is decreasing. If  $f''(x) > 0$  it is **concave up**, if  $f''(x) < 0$  it is **concave down**. If  $f'(x) = 0$  then  $f$  has a horizontal tangent.

Hoptial tells that limits  $\lim_{x \rightarrow p} f(x)/g(x)$ , where  $f(p) = g(p) = 0$  or  $f(p) = g(p) = \infty$  with  $g'(p) \neq 0$  are given by  $f'(p)/g'(p)$ .

With  $Df(x) = (f(x+h) - f(x))/h$  and  $S(x) = h(f(h) + f(2h) + \dots f(kh))$  we have  $SDf(kh) = f(kh) - f(0)$  and  $DS(f(kh)) = f(kh)$ . This is a preliminary fundamental theorem of calculus.

Roots of  $f(x)$  with  $f(a) < 0, f(b) > 0$  can be obtained numerically by dissection or by applying the **Newton map**  $T(x) = x - f(x)/f'(x)$  again and again.

### Algebra reminders

Healing:  $(a+b)(a-b) = a^2 - b^2$  or  $1 + a + a^2 + a^3 + a^4 = (a^5 - 1)/(a - 1)$   
 Denominator:  $1/a + 1/b = (a+b)/(ab)$   
 Exponential:  $(e^a)^b = e^{ab}, e^a e^b = e^{a+b}, a^b = e^{b \log(a)}$   
 Logarithm:  $\log(ab) = \log(a) + \log(b), \log(a^b) = b \log(a)$   
 Trig functions:  $\cos^2(x) + \sin^2(x) = 1, \sin(2x) = 2 \sin(x) \cos(x), \cos(2x) = \cos^2(x) - \sin^2(x)$   
 Square roots:  $a^{1/2} = \sqrt{a}, a^{-1/2} = 1/\sqrt{a}$

### Important functions

Polynomials	$x^3 + 2x^2 + 3x + 1$	Exponential	$5e^{3x}$
Rational functions	$(x+1)/(x^3 + 2x + 1)$	Logarithm	$\log(3x)$
Trig functions	$2 \cos(3x)$	Inverse trig functions	$\arctan(x)$

### Important derivatives

$f(x) = x^n$	$f'(x) = nx^{n-1}$	$f(x) = \sin(ax)$	$f'(x) = a \cos(a \cdot x)$
$f(x) = e^{ax}$	$f'(x) = ae^{ax}$	$f(x) = \tan(x)$	$f'(x) = 1/\cos^2(x)$
$f(x) = \cos(ax)$	$f'(x) = -a \sin(a \cdot x)$	$f(x) = \log(x)$	$f'(x) = 1/x$

### Differentiation rules

Addition rule	$(f+g)' = f' + g'$	Quotient rule	$(f/g)' = (f'g - fg')/g^2$
Scaling rule	$(cf)' = cf'$	Chain rule	$(f(g(x)))' = f'(g(x))g'(x)$
Product rule	$(fg)' = f'g + fg'$	Easy rule	simplify before deriving

### Extremal problems

- Build a fence of length  $x+2y = 12$  with dimensions  $x$  and  $y$  with maximal area  $A = xy$ .
- Find the largest area  $A = 4xy$  of a rectangle with vertices  $(x, y), (-x, y), (-x, -y), (x, -y)$  inscribed in the ellipse  $x^2 + 2y^2 = 1$ .
- Which isosceles triangle of height  $h$  and base  $2x$  and area  $xh = 1$  has minimal circumference  $2x + 2\sqrt{x^2 + h^2}$ ?
- Where is the distance  $\sqrt{x^2 + y^2}$  of the parabola  $y = x^2 = 2$  to the point  $(0, 0)$  minimal?
- A cone of height  $h = 1 + x$  and radius  $r = \sqrt{1 - x^2}$  is tightly enclosed by a unit sphere centered at height  $x$ . Maximize the volume  $r^2\pi h/3$  of the cone.
- Maximize  $f(x) = \sin(x)$  on  $[0, \pi]$ .

### Limit examples

$\lim_{x \rightarrow 0} \sin(x)/x$	l'Hopital 0/0	$\lim_{x \rightarrow 1} (x^2 - 1)/(x + 1)$	heal directly
$\lim_{x \rightarrow 0} (1 - \cos(x))/x^2$	l'Hopital 0/0 twice	$\lim_{x \rightarrow \infty} \exp(x)/(1 + \exp(x))$	l'Hopital
$\lim_{x \rightarrow 0} x \log(x)$	l'Hopital $\infty/\infty$	$\lim_{x \rightarrow 0} (x+1)/(x+5)$	no work necessary

### Important things

Summation and taking differences is at the hart of calculus  
 The 3 major types of discontinuities are jump, oscillation, infinity  
 The Newton method is an algorithm to find roots  
 Remember the fundamental theorem of trigonometry  $\lim_{x \rightarrow 0} \sin(x)/x = 1$ .  
 The derivative is the limit  $Df(x) = [f(x+h) - f(x)]/h$  as  $h \rightarrow 0$ . It is called rate of change.  
 The rule  $D(1+h)^{x/h} = (1+h)^{x/h}$  leads to  $\exp'(x) = \exp(x)$ .

### More Examples

- Find  $\lim_{x \rightarrow 1} (x^{1/4} - 1)/(x^{1/5} - 1)$ . Answer: 5/4.
- Find  $\lim_{x \rightarrow 1} \sin(4x - 4)/(x - 1)$ . Answer: 4.
- Find  $\lim_{x \rightarrow 2} \frac{3 - \sqrt{7+x}}{x-2}$ . Answer -1/6
- Find  $\arcsin'(5x^2)$ . Answer:  $10x(1 - 25x^4)^{-1/2}$
- Is  $1/\log|x|$  continuous at  $x = 0$ . Answer: yes
- Is  $\log(1/|x|)$  continuous at  $x = 0$ . Answer: no