

## Lecture 34: Calculus and Statistics

In this lecture, we look at an application of calculus to statistics. We have already defined the probability density function  $f$  and its anti-derivative, the cumulative distribution function.

### Probability density

Recall that a probability density function is a function  $f$  satisfying  $\int f(x) dx = 1$  and which has the property that it is  $\geq 0$  everywhere. We say  $f$  is a probability density function on an interval  $[a, b]$  if  $\int_a^b f(x) dx = 1$  and  $f(x) \geq 0$  there. In such a case, we assume that  $f$  is zero outside the interval.

Recall also that we called the antiderivative of  $f$  the cumulative distribution function  $F(x)$  (CDF).

### Expectation

The **expectation** of probability density function  $f$  is

$$m = \int_{-\infty}^{\infty} x f(x) dx .$$

In the case, when the probability density function is zero outside some interval, we have

The **expectation** of probability density function  $f$  defined on some interval  $[a, b]$  is

$$m = \int_a^b x f(x) dx .$$

### Variance and Standard deviation

The **variance** of probability density function  $f$  is

$$\int_{-\infty}^{\infty} (x - m)^2 f(x) dx ,$$

where  $m$  is the expectation.

Again, if the probability density function is defined on some interval  $[a, b]$  then

The **variance** of probability density function  $f$  is

$$\int_a^b (x - m)^2 f(x) dx ,$$

where  $m$  is the expectation of  $f$ .

The square root of the variance is the **standard deviation**.

### Examples

In the lecture, we will compute this in some examples. Here is some sample.

- 1 The expectation of the geometric distribution  $f(x) = e^{-x}$

$$\int x e^{-x} dx = 1 .$$

The variance of the geometric distribution  $f(x) = e^{-x}$  is 1 and the standard deviation 1 too.

Remember that we can compute also with Tic-Tac-Toe:

$$\int x^2 e^{-x} dx$$

$x^2$	$e^{-x}$	
$2x$	$-e^{-x}$	$\oplus$
$2$	$e^{-x}$	$\ominus$
$0$	$e^{-x}$	$\oplus$

- 2 The expectation of the standard Normal distribution  $f(x) = (2\pi)^{-1/2} e^{-x^2/2}$

$$\int_0^{\infty} x (2\pi)^{-1/2} e^{-x^2/2} dx = 0 .$$

### Homework

- 1 The function  $f(x) = \cos(x)/2$  on  $[-\pi/2, \pi/2]$  is a probability density function. Its mean is 0. Find its variance

$$\int_{-\pi/2}^{\pi/2} x^2 \cos(x) dx .$$

- 2 The **uniform distribution** on  $[a, b]$  is a distribution, where any real number between  $a$  and  $b$  is equally likely to occur. The probability density function is  $f(x) = 1/(b - a)$  for  $a \leq x \leq b$  and 0 elsewhere. Verify that  $f(x)$  is a valid probability density function.

- 3 Verify that the function which is 0 for  $x < 0$  and equal to

$$f(x) = \frac{1}{\log(2)} \frac{e^{-x}}{1 + e^{-x}}$$

for  $x \geq 0$  is a probability density function.

- 4 A particular **Cauchy distribution** has the probability density

$$f(x) = \frac{1}{\pi} \frac{1}{(x - 1)^2 + 1} .$$

Verify that  $f(x)$  is a valid probability density function.

- 5 Find the cumulative distribution function (CDF)  $F(x)$  of  $f$  in the previous problem.