

5/11/2013: Practice final A

Your Name:

- Try to answer each question on the same page as the question is asked. If needed, use the back or the next empty page for work. If you need additional paper, write your name on it.
- Do not detach pages from this exam packet or unstaple the packet.
- All functions f if not specified otherwise can be assumed to be smooth so that arbitrary many derivatives can be taken.
- Please write neatly. Answers which are illegible for the grader can not be given credit.
- Except for multiple choice problems, give computations.
- No notes, books, calculators, computers, or other electronic aids are allowed.
- You have 180 minutes time to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
12		10
13		10
Total:		140

Problem 1) TF questions (20 points). No justifications are needed.

- 1) T F The definite integral $\int_0^{2\pi} \sin^2(5x) dx$ is zero.

Solution:

The integrand is never negative and almost everywhere positive.

- 2) T F The intermediate value theorem assures that the function $\exp(\sin(x))$ has a root in the interval $(0, 2\pi)$.

Solution:

The function $\exp(\sin(x))$ is never zero.

- 3) T F $\frac{d}{dx} \cos(4x) = -4 \sin(4x)$.

Solution:

differentiate

- 4) T F If $f''(1) < 0$ then 1 is a local maximum of f .

Solution:

It also has to be a critical point.

- 5) T F The derivative of $1/x$ is $\log(x)$ for all $x > 0$.

Solution:

It is the anti-derivative, not the anti derivative

- 6) T F The limit of $\sin(3x)/(5x)$ for $x \rightarrow 0$ exists and is equal to $3/5$.

Solution:

Use Hôpital

- 7) T F The function $(e^t - 1)/t$ has the limit 1 as t goes to zero.

Solution:
Use Hopital

- 8) T F The derivative of $f(f(x))$ is $f'(f'(x))$ for any differentiable function f .

Solution:
This is not the chain rule

- 9) T F A monotonically increasing function f has no point x , where $f'(x) < 0$.

Solution:
Increasing means that the derivative is positive.

- 10) T F The function $f(x) = \exp(-x^2)$ has an inflection point x somewhere on the real line.

Solution:
The second derivative can be zero. One can see this by looking at the graph.

- 11) T F The function $f(x) = (1 - x^3)/(1 + x)$ has a limit for $x \rightarrow -1$.

Solution:
The top $1 - x^3$ is not zero at $x = -1$ so that the function has a pole

- 12) T F If we know the marginal cost for all quantities x as well as the total cost for $x = 1$ we know the total cost for all x .

- 13) T F The function f which satisfies $f(x) = 0$ for $x < 0$ and $f(x) = e^{-x}$ for $x \geq 0$ is a probability density function.

Solution:
True, it is nonnegative every where and the total integral is 1.

- 14) T F The differentiation rule $(f \cdot g)' = f'(g(x)) \cdot g'(x)$ holds for all differentiable functions f, g .

Solution:

We would need the Leibniz rule, not the chain rule.

- 15) T F Hôpital's rule assures that $\cos(x)/\sin(x)$ has a limit as $x \rightarrow 0$.

Solution:

The nominator does not go to zero for $x \rightarrow 0$.

- 16) T F A Newton step for the function f is $T(x) = x - \frac{f(x)}{f'(x)}$.

Solution:

By definition

- 17) T F The family of functions $f_c(x) = cx^2$ where c is a parameter has a catastrophe at $x = 0$.

Solution:

For $c < 0$ we have a local max, for $c > 0$ we have a local min.

- 18) T F The fundamental theorem of calculus implies $\int_{-x}^x f'(t) dt = f(x) - f(-x)$ for all differentiable functions f .

Solution:

Yes, this is the most important result in this course.

- 19) T F If f is a smooth function for which $f''(x) = 0$ everywhere, then f is constant.

Solution:

It can be linear

- 20) T F The function $f(x) = \sin(x)/(1 - \cos(x))$ can be assigned a value $f(0)$ such that $f(x)$ is continuous at 0.

Solution:

Use l'Hopital to see that the limit is the same as $\lim_{x \rightarrow 0} \cos(x)/\sin(x)$ which has no limit at $x = 0$.

Problem 2) Matching problem (10 points) Only short answers are needed.

We name some important concepts in this course. To do so, please complete the sentences with one or two words. Each question is one point.

$\lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$ is called the		of f .
$f'(x) = 0, f''(x) > 0$ implies that x is a		of f .
The sum $\frac{1}{n}[f(0) + f(1/n) + f(2/n) + \dots + f((n-1)/n) + f(1)]$ is called a		sum.
If $f(0) = -3$ and $f(4) = 8$, then f has a root on the interval $(0, 4)$ by the		theorem.
There is a point $x \in (0, 1)$ where $f'(x) = f(1) - f(0)$ by the		theorem.
The expansion rate $r'(t)$ can be obtained from $d/dtV(r(t)) = -5$ by the method of		rates.
The anti derivative $\int_{-\infty}^x f(t) dt$ of a probability density function f is called the		function.
A point x for which $f(x) = 0$ is called a		of f .
A point x for which $f''(x) = 0$ is called an		of f .
At a point x for which $f''(x) > 0$, the function is called		up.

Solution:

Derivative
 Local minimum
 Riemann sum
 Intermediate value
 Mean value
 Related
 Cumulative distribution
 Root
 Inflection Point
 Concave


Problem 3) Matching or short answer problem (10 points). No justifications are needed.


a) (4 points) Find the relation between the following functions:

function f	function g	$f = g'$	$g = f'$	none
$\log \sin(x) $	$\cot(x)$			
$1/\cos^2(x)$	$\tan(x)$			
x^5	$5x^4$			
$1/x^2$	$-1/x$			
$\sin(\log(x))$	$\cos(\log(x))/x$			

b) (3 points) Match the following functions (a-d) with a choice of **anti-derivatives** (1-4).

Function a)-d)	Fill in 1)-4)
graph a)	
graph b)	
graph c)	
graph d)	





Function f	$\lim_{x \rightarrow 0} f(x)$
$x/(e^{2x} - 1)$	
$(e^{2x} - 1)/(e^{3x} - 1)$	
$\sin(3x)/\sin(5x)$	

Solution:

a)


function f	function g	$f = g'$	$g = f'$	none
$\log \sin(x) $	$\cot(x)$		*	
$1/\cos^2(x)$	$\tan(x)$	*		
x^5	$5x^4$		*	
$1/x^2$	$-1/x$	*		
$\sin(\log(x))$	$\cos(\log(x))/x$		*	

b) 3,2,4,1

c) Use l'Hopital: 1/2,2/3,3/5

Problem 4) Area computation (10 points)

Find the area of the shield shaped region bound by the two curves $1/(1+x^2)$ and $x^2 - 1$.




Solution:

The two curves intersect at $x = \pm 2^{1/4}$.

$$\int_{-2^{1/4}}^{2^{1/4}} \frac{1}{1+x^2} - x^2 + 1 \, dx = \arctan(x) - x^3/3 + x \Big|_{-2^{1/4}}^{2^{1/4}} = 2 \arctan(2^{1/4}) - (2/3)2^{3/4} + 2 \cdot 2^{1/4}.$$

Problem 5) Volume computation (10 points)

Did you know that there is a scaled copy of the **liberty bell** on the campus of the Harvard business school? Here we compute its volume. Find the volume of the rotationally symmetric solid if the radius $r(z)$ at height z is $r(z) = 8 - (z - 1)^3$ and the height z of the bell is between 0 and 3.



Solution:

$$\pi \int_0^3 \pi(8-(z-1)^3)^2 dz = \pi \int_{-1}^2 (8-u^3)^2 du = \pi \int_{-1}^2 64-16u^3+u^6 du = \pi(64z-16u^4/4+u^7/7)|_{-1}^3 = \pi 1053/7;$$

Problem 6) Improper integrals (10 points)

a) (5 points) Find the integral or state that it does not exist

$$\int_1^{\infty} \frac{1}{x^4} dx .$$

b) (5 points) Find the integral or state that it does not exist

$$\int_1^{\infty} \frac{1}{x^{3/2}} dx .$$

Solution:

a)


$$\int_1^{\infty} \frac{1}{x^4} dx = \frac{-x^{-3}}{3} \Big|_1^{\infty} = \frac{1}{3} .$$

b)

$$\int_1^{\infty} \frac{1}{x^{3/2}} dx = -2x^{-1/2} \Big|_1^{\infty} = 2 .$$

Problem 7) Extrema (10 points)

The Harvard stadium has a track which encloses a rectangular field of dimensions x, y . The circumference of the track is $400 = 2\pi y + 2x$ and is fixed. We want to maximize the area xy for a play field. Which x achieves this?



Solution:

Solve for $y = (200 - x)/\pi$ and plug this into the function to get

$$f(x) = xy = x(200 - x)/\pi .$$

To find the maximum of this function, we differentiate with respect to x and look where the derivative is zero:

$$f'(x) = (200 - 2x)/\pi = 0$$

showing that $x = 100$ is the maximum.

Problem 8) Integration by parts (10 points)

Find the antiderivative:

$$\int (x - 1)^4 \exp(x + 1) dx .$$

Solution:

Use the Tic-Tac-Toe integration method:

$(x - 1)^4$	$\exp(x + 1)$	
$4(x - 1)^3$	$\exp(x + 1)$	\oplus
$12(x - 1)^2$	$\exp(x + 1)$	\ominus
$24(x - 1)$	$\exp(x + 1)$	\oplus
24	$\exp(x + 1)$	\ominus
0	$\exp(x + 1)$	\oplus

Adding things up gives

$$e^{x+1}[(x - 1)^4 - 4(x - 1)^3 + 12(x - 1)^2 - 24(x - 1) + 24] .$$

Problem 9) Substitution (10 points)

- a) (3 points) Solve the integral $\int e^{x^2} 2x \, dx$.
- b) (3 points) Solve the integral $\int 2x \log(x^2) \, dx$.
- c) (4 points) Find the integral $\int e^{-2e^x} e^x \, dx$.

Solution:

These are all standard substitution problems:

- a) $e^{x^2} + c$
- b) $x^2 \log(x^2) - x^2 + c$
- c) $-e^{-2e^x} / 2 + c$

Problem 10) Partial fractions (10 points)

- a) (5 points) Find the definite integral

$$\int_1^5 \frac{1}{(x - 4)(x - 2)} \, dx .$$

- b) (5 points) Find the indefinite integral

$$\int \frac{1}{(x - 1)(x - 3)(x - 5)} \, dx .$$

Solution:

In both problems we can find the coefficients quickly with the l'Hopital method: a) $\int_1^5 \frac{1}{(x-4)(x-2)} dx = \frac{1}{2} \int_1^5 \left[\frac{1}{x-4} - \frac{1}{x-2} \right] dx = \frac{1}{2} [\log|x-4| - \log|x-2|]_1^5 = -\log(3)$. b) The factorization

$$\frac{1}{(x-1)(x-3)(x-5)} = \frac{A}{x-1} + \frac{B}{x-3} + \frac{C}{x-5}$$

can be obtained quickly from l'Hopital: $A = \lim_{x \rightarrow 1} \frac{1}{(x-3)(x-5)} = \frac{1}{8}$ and $B = \lim_{x \rightarrow 3} \frac{1}{(x-1)(x-5)} = -\frac{1}{4}$ and $C = \lim_{x \rightarrow 5} \frac{1}{(x-1)(x-3)} = \frac{1}{8}$ so that the result is

$$[\log|x-1| - 2\log|x-3| + \log|x-5|]/8 .$$


[P.S. As in the homeworks, we do not worry in a) that these are improper integrals, integrating over the logarithmic singularity. They are no problem because the integral of $\log|x|$ is $x \log|x| - x$ which has a limit 0 for $x \rightarrow 0$.]

Problem 11) Related rates (10 points)

The coordinates of a car on a freeway intersection are $x(t)$ and $y(t)$. They are related by

$$x^7 + y^7 = 2xy^2 .$$

We know $x' = 3$ at $x = 1, y = 1$. Find y' .

**Solution:**

Differentiate the relation with respect to t and solve for y' :

$$7x^6x' + 6y^2y' = 2x'y^2 + 4xyy' .$$

Therefore,

$$y' = (7x^6x' - 2y^2x') / (4xy - 6y^2) .$$

Which gives -5 .

Problem 12) Various integration problems (10 points)

Find the anti-derivatives of the following functions:

a) (2 points) $f(x) = \sin^5(x) \cos(x)$.

b) (3 points) $f(x) = \frac{1}{x^2+1} + \frac{1}{x^2-1}$.

c) (2 points) $f(x) = \sqrt{1-x^2} + \frac{1}{\sqrt{1-x^2}}$.

d) (3 points) $f(x) = \log(x) + \frac{1}{\log(x)}$.

Solution:

a) $\sin^6(x)/6 + c$

b) $\arctan(x) + \log(x-1)/2 + \log(x+1)/2 + c$

c) $\arcsin(x) + (1 + \cos(2 \arcsin(x)))/2$

d) $x \log(x) - x + \text{li}(x)$ the second integral is a nonelementary integral. Was a freebe. You got 3 points even without solving that...

Problem 13) Applications (10 points)

a) (5 points) We know the total cost $F(x) = -x^3 + 2x^2 + 4x + 1$ for the quantity x . In order to find the positive **break-even point** x satisfying $f(x) = g(x)$, where $g(x) = F(x)/x$ is the total cost and $f(x) = F'(x)$ is the marginal cost, we do - how sweet it is - find the maximum of the average cost $g(x) = F(x)/x$. Find the maximum!

b) (5 points) We know the **"velocity"**, **"acceleration"** and **"jerk"** as the first second and third derivative of position. The fourth, fifth and sixth derivatives of position as a function of time are called **"snap"**, **"crackle"** and **"pop"** according to characters used in a cereal add. Assume we know the snap $x''''(t) = t$. Find $x(t)$ satisfying $x(0) = x'(0) = x''(0) = 0, x'''(0) = 0$.



Solution:

a) We have to solve the equation $g(x) = 0$ by the strawberry theorem. Giving the equation $-x^2 + 2x + 4 + 1/x = 0$ was enough. The solution needs to be evaluated numerically, for example with Newton.

b) Integrate 4 times to get $x(t) = t^5/120$. All constants are zero.