


Lecture 8: The derivative function

We have defined the derivative $f'(x) = \frac{d}{dx}f(x)$ as a limit of $Df(x)$ for $h \rightarrow 0$. We have seen that $\frac{d}{dx}x^n = nx^{n-1}$ holds for integer n . We also know already that $\sin'(ax) = a \cos(x)$, $\cos'(ax) = -a \sin(x)$ and $\exp'(ax) = a \exp(x)$. We can already differentiate a lot of functions and evaluate the derivative $f'(x)$ at some point x . This is the slope of the curve at x .

- 1 Find the derivative $f'(x)$ of $f(x) = \sin(4x) + \cos(5x) - \sqrt{x} + 1/x + x^4$ and evaluate it at $x = 1$. **Solution:** $f'(x) = 4 \cos(4x) - 5 \sin(5x) - 1/(2\sqrt{x}) - 1/x^2 + 4x^3$. Plugging in $x = 1$ gives $-\pi - 1/2 - 1 + 4$.

The function which takes the derivative at a given point is called the **derivative function**. For example, for $f(x) = \sin(x)$, we get $f'(x) = \cos(x)$. In this lecture, we want to understand the new function and its relation with f . What does it mean if $f'(x) > 0$. What does it mean that $f'(x) < 0$. Do the roots of f tell something about f' or do the roots of f' tell something about f ?


Here is an example of a function and its derivative. Can you see the relation?



To understand the relation, it is good to distinguish intervals, where $f(x)$ is increasing or decreasing. This are the intervals where $f'(x)$ is positive or negative.

A function is called **monotonically increasing** on an interval $I = (a, b)$ if $f'(x) > 0$ for all $x \in (a, b)$. It is **monotonically decreasing** if $f'(x) < 0$ for all $x \in (a, b)$.

Lets look at the previous example again.




- 2 Can you find a function f which is bounded $|f(x)| \leq 1$ and such that $f'(x)$ is unbounded?


Given the function $f(x)$, we can define $g(x) = f'(x)$ and then take the derivative g' of g . This second derivative $f''(x)$ is called the **acceleration**. It measures the rate of change of the tangent slope. For $f(x) = x^4$, for example we have $f''(x) = 12x^2$. If $f''(x) > 0$ on some interval the function is called **concave up**, if $f''(x) < 0$, it is **concave down**.

- 3 Find a function f which has the property that its acceleration is constant equal to 6. **Solution.** We have to get a function such that its derivative is $6x$. That works for $3x^2$.
- 4 Find a function f which has the property that its acceleration is equal to its derivative. To do so, try basic functions you know and compute $f'(x)$, $f''(x)$ in each case.

After matching some functions, we look at related inverse problem called **bottle calibration problem**. We fill a circular bottle or glass with constant amount of fluid. Plot the height of the fluid in the bottle at time t . Assume the radius of the bottle is $f(z)$ at height z . Can you find a formula for the height $g(t)$ of the water? This is not so easy. But we can find the rate of change $g'(t)$. Assume for example that f is constant, then the rate of change is constant and the height of the water increases linearly like $g(t) = t$. If the bottle gets wider, then the height of the water increases slower. There is definitely a relation between the rate of change of g and f . Before we look at this more closely, lets try to match the following cases of bottles with the graphs of the functions g qualitatively.

- 5 In each of the bottles, we call g the height of the water level at time t , when filling the bottle with a constant stream of water. Can you match each bottle with the right height function?





The key is to look at $g'(t)$, the rate of change of the height function. Because $[g(t+h) - g(t)]$ times the area πf^2 is a constant times the time difference $h = dt$, we have

$$g' = \frac{1}{\pi f^2}.$$


This formula relates the derivative function of g with the thickness $f(t)$ of the bottle at height g . It tells that if f is large, then g' is small and if f is small, then g' is large.

Homework


- 1) For the following functions, determine on which intervals the function is monotonically increasing or monotonically decreasing.

- a) $f(x) = x^3 - x$ on $[-2, 2]$
- b) $f(x) = \cos(x)$ on $[-\pi, \pi]$
- c) $f(x) = x^4 - 8x^2$ on $[-4, 4]$.

- 2) Match the following functions with their derivatives. Give short explanations for each match.



- 3) Match also the following functions with their derivatives. Give short explanations documenting your reasoning in each case.



- 4) Draw for the following functions the graph of the function $f(x)$ as well as the graph of its derivative $f'(x)$. You do not have to compute the derivative analytically as a formula here since we do not have all tools yet to compute the derivatives. The derivative function you draw needs to have the right qualitative shape however.

- a) The "To whom the bell tolls" function

$$f(x) = e^{-x^2}$$

- b) The "witch of Maria Agnesi" function:

$$f(x) = \frac{1}{1+x^2}$$

- c) The three gorges function

$$f(x) = \frac{1}{x} + \frac{1}{x-1} + \frac{1}{x+1}.$$

- 5) Below you the graphs of three derivative functions $f'(x)$. In each case you are told that $f(0) = 1$. Your task is to draw the function $f(x)$ in each of the cases a), b), c). Your picture does not have to be up to scale, but your drawing should display the right features.

