

# Lecture 15: Review for first midterm

## Major points

A function is **continuous**, if whenever  $x, y$  are close, also  $f(x), f(y)$  are close. Formally, for every  $a$  there exists  $b = f(a)$  such that  $\lim_{x \rightarrow a} f(x) = b$  for every  $a$ . The intermediate value theorem:  $f(a) > 0, f(b) < 0$  implies  $f$  having a root in  $(a, b)$ .

If  $f'(x) = 0$  and  $f''(x) > 0$  then  $x$  is a **local minimum**. If  $f'(x) = 0$  and  $f''(x) < 0$  then  $x$  is a **local maximum**. To find **global minima or maxima**, compare local extrema and boundary values.

If  $f$  changes sign we have a **root**  $f = 0$ , if  $f'$  changes sign, we have a **critical point**  $f' = 0$  if  $f''$  changes sign, we have an **inflection points**. A function is **even** if  $f(-x) = f(x)$ , and **odd** if  $f(-x) = -f(x)$ . Odd functions have 0 as roots, even functions have 0 as critical point.

If  $f' > 0$  then  $f$  is increasing, if  $f' < 0$  it is decreasing. If  $f''(x) > 0$  it is **concave up**, if  $f''(x) < 0$  it is **concave down**. If  $f'(x) = 0$  then  $f$  has a horizontal tangent.

Hôpital's theorem applies for  $0/0$  or  $\infty/\infty$  situations. In that case,  $\lim_{x \rightarrow p} f(x)/g(x)$ , where  $f(p) = g(p) = 0$  or  $f(p) = g(p) = \infty$  with  $g'(p) \neq 0$  are given by  $f'(p)/g'(p)$ .

With  $Df(x) = (f(x+h) - f(x))/h$  and  $S(x) = h(f(0) + f(2h) + \dots + f((k-1)h))$  we have a **preliminary fundamental theorem of calculus**  $SDf(kh) = f(kh) - f(0)$  and  $DS(f(kh)) = f(kh)$ .

Roots of  $f(x)$  with  $f(a) < 0, f(b) > 0$  can be obtained by the dissection method by applying the **Newton map**  $T(x) = x - f(x)/f'(x)$  again and again.

## Algebra reminders

Healing:	$(a+b)(a-b) = a^2 - b^2$ or $1 + a + a^2 + a^3 + a^4 = (a^5 - 1)/(a - 1)$
Denominator:	$1/a + 1/b = (a+b)/(ab)$
Exponential:	$(e^a)^b = e^{ab}, e^a e^b = e^{a+b}, a^b = e^{b \log(a)}$
Logarithm:	$\log(ab) = \log(a) + \log(b), \log(a^b) = b \log(a)$
Trig functions:	$\cos^2(x) + \sin^2(x) = 1, \sin(2x) = 2 \sin(x) \cos(x), \cos(2x) = \cos^2(x) - \sin^2(x)$
Square roots:	$a^{1/2} = \sqrt{a}, a^{-1/2} = 1/\sqrt{a}$

## Important functions

Polynomials	$x^3 + 2x^2 + 3x + 1$	Exponential	$5e^{3x}$
Rational functions	$(x+1)/(x^3 + 2x + 1)$	Logarithm	$\log(3x)$
Trig functions	$2 \cos(3x)$	Inverse trig functions	$\arctan(x)$

## Important derivatives

$f(x)$	$f'(x)$	$f(x)$	$f'(x)$
$f(x) = x^n$	$nx^{n-1}$	$f(x) = \sin(ax)$	$a \cos(ax)$
$f(x) = e^{ax}$	$ae^{ax}$	$f(x) = \tan(x)$	$1/\cos^2(x)$
$f(x) = \cos(ax)$	$-a \sin(ax)$	$f(x) = \log(x)$	$1/x$

## Differentiation rules

Addition rule	$(f + g)' = f' + g'$ .	Quotient rule	$(f/g)' = (f'g - fg')/g^2$ .
Scaling rule	$(cf)' = cf'$ .	Chain rule	$(f(g(x)))' = f'(g(x))g'(x)$ .
Product rule	$(fg)' = f'g + fg'$ .	Easy rule	simplify before deriving

## Extremal problems

- 1 Build a fence of length  $x + 2y = 12$  and dimensions  $x$  and  $y$  with maximal area  $A = xy$ .
- 2 Find the largest area  $A = 4xy$  of a rectangle with vertices  $(x, y), (-x, y), (-x, -y), (x, -y)$  inscribed in the ellipse  $x^2 + 2y^2 = 1$ .
- 3 Which isosceles triangle of height  $h$  and base  $2x$  and area  $xh = 1$  has minimal circumference  $2x + 2\sqrt{x^2 + h^2}$ ?

To extremize  $f$  on an interval  $[a, b]$ , find all critical points inside the interval, evaluate  $f$  on the boundary  $f(a), f(b)$  and then compare the values to find the global maximum. Do not make the second derivative test at the boundary.

## Limit examples

$\lim_{x \rightarrow 0} \sin(x)/x$	l'Hopital 0/0	$\lim_{x \rightarrow 1} (x^2 - 1)/(x - 1)$	heal
$\lim_{x \rightarrow 0} (1 - \cos(x))/x^2$	l'Hopital 0/0 twice	$\lim_{x \rightarrow \infty} \exp(x)/(1 + \exp(x))$	l'Hopital
$\lim_{x \rightarrow 0} x \log(x)$	l'Hopital $\infty/\infty$	$\lim_{x \rightarrow 0} (x + 1)/(x + 5)$	no work necessary

## Important things

Summation and taking differences is at the heart of calculus

The 3 major types of discontinuities are jump, oscillation, infinity

Dissection and Newton methods are algorithms to find roots.

The fundamental theorem of trigonometry is  $\lim_{x \rightarrow 0} \sin(x)/x = 1$ .

The derivative is the limit  $Df(x) = [f(x+h) - f(x)]/h$  as  $h \rightarrow 0$ .

The rule  $D(1+h)^{x/h} = (1+h)^{x/h}$  leads to  $\exp'(x) = \exp(x)$ .

## More Examples

- 1 Is  $1/\log|x|$  continuous at  $x = 0$ . Answer: yes with  $f(0) = 0$
- 2 Is  $\log(1/|x|)$  continuous at  $x = 0$ . Answer: no.
- 3  $\lim_{x \rightarrow 1} (x^{1/3} - 1)/(x^{1/4} - 1)$ . Answer:  $4/3$ .