



FUNCTIONS OF ONE VARIABLES. A function of one variables $f(x)$ assigns to a variables x a real number $f(x)$. Example: $f(x) = x \sin(x)$. We can visualize a function by its **graph** $\{(x, y), y = f(x)\}$ which is a curve in space.




FUNCTIONS OF TWO VARIABLES. A function of two variables $f(x, y)$ assigns to two variables x, y a real number $f(x, y)$. Example: $f(x, y) = \sin(xy)$, temperature distribution on a plate.

In the same way as functions of one variable were visualized by drawing the graph $y = f(x)$ in space, we can visualize a function of three variables as a graph $\{(x, y, z) \mid z = f(x, y)\}$ in space.



FUNCTIONS OF THREE VARIABLES. A function of three variables $g(x, y, z)$ assigns to three variables x, y, z a real number $g(x, y, z)$. Example: $g(x, y, z) = \sin(xyz)$, temperature distribution in space. We can no more draw a graph of g . But we can visualize it differently by drawing surfaces $g(x, y, z) = c$, where g is constant.




SURFACES. Many surfaces can be described as a function of three variables g . The points (x, y, z) satisfying the equation $g(x, y, z) = c$ for a surface. Examples.

Graphs: $g(x, y, z) = z - f(x, y)$. If $g(x, y, z) = 0$, then $z = f(x, y)$ and the surface is a graph of a function of two variables.

Planes: $ax + by + cz = d$ is a plane orthogonal to the vector $\vec{n} = (a, b, c)$. The equation says $\vec{n} \cdot \vec{x} = d$. If a point \vec{x}_0 is on the plane, then $\vec{n} \cdot \vec{x}_0 = d$. The equation can also be written as $\vec{n} \cdot (\vec{x} - \vec{x}_0) = 0$ which means that every vector $\vec{x} - \vec{x}_0$ is orthogonal to \vec{n} .


Quadratics If $f(x, y, z) = ax^2 + by^2 + cz^2 + dxy + exz + fyz + gx + hy + kz + m$ the a surface $f(x, y, z) = 0$ is called a **quadratic**. Below are some examples.

QUARTIC



$p(x, y, z)$ degree 4 polynomial in x, y, z

SEXTIC




$p(x, y, z)$ degree 6 polynomial in x, y, z


DECIC




$p(x, y, z)$ degree 10 polynomial in x, y, z



$$(x - a)^2 + (y - b)^2 + (z - c)^2 = r^2$$



$$(x - a)^2 + (y - b)^2 - c = z$$




$$ax + by + cz = d$$


HYPERBOLOID I

CYLINDER


HYPERBOLOID II



$$(x - a)^2 + (y - b)^2 - (z - c)^2 = r^2$$



$$(x - a)^2 + (y - b)^2 = r^2$$




$$(x - a)^2 + (y - b)^2 - (z - c)^2 = -r^2$$


ELLIPSOID

HYPERBOLIC PARABOLOID


DEFORMED HYPERBOLOID



$$x^2/a^2 + y^2/b^2 + z^2/c^2 = 1$$



$$x^2 - y^2 + z = 1$$



$$x^2/a^2 + y^2/b^2 - z^2/c^2 = 1$$

TRACES. To draw surfaces, it helps to look at the **traces**, the intersections of the surfaces with the coordinate planes $x = 0$, $y = 0$ or $z = 0$. Other points to look at are the **intercepts**, the intersections of the surface with the coordinate axes. The traces are shown in the pictures of the quadrics above.

For example: for the **one-sided hyperboloid** $x^2 + y^2 - z^2 = 1$ (called HYPERBOLOID I above), the z -trace is $x^2 + y^2 = 1$, a circle, the x -traces $y^2 - z^2 = 1$ is a hyperbola as well as the y -trace $x^2 - z^2 = 1$.