

Homework 15: Tangent lines and planes

This homework is due Friday, 10/18.

- 1 a) Find the tangent plane to the surface


$$x^2 + y^2 - x^2y^2 - 2z^2 + x^2z^2 = 0$$

at the point $(x, y, z) = (-1, -2, -1)$.

- b) Find the tangent line to the curve

$$x^2 + y^2 - x^2y^2 = -23$$

at the point $(x, y) = (3, 2)$.



Solution:

a) $\nabla f = [2x - 2xy^2 + 2xz^2, 2y - 2x^2y, -2z + 2x^2z]$. At the point $-1, -2, -1$ it is $[4, 0, -2]$. The equation of the plane is $\boxed{4x - 2z = -2}$.

b) $\nabla f = [2x - 2xy^2, 2y - 2x^2y] = [-18, -32]$. The equation of the line is $\boxed{9x + 16y = 59}$, where the constant 59 was obtained by plugging in the point $(x, y) = (3, 2)$.

- 2 In each of the following four conceptual problems, please answer briefly.

a) The figure 8 curve $f(x, y) = x^4 - x^2 + y^2 = 0$ has no tangent line at $(0, 0)$. Why? Isn't it perfectly smooth function $f(x, y)$?

b) The following statement is nonsense: "the tangent plane to the graph of $f(x, y, z) = x^2 + y^2 + z^2$ at $(x, y, z) = (1, 2, 3)$ is $2x + 4y + 6z = 28$ ". Modify it to make it a true statement.

c) Let $\vec{r}(u, v) = [u, v, g(u, v)]$ and $[a, b, c] = \vec{r}_u(1, 1) \times \vec{r}_v(1, 1)$. Let $f(x, y, z) = z - g(x, y)$ and $[A, B, C] = \nabla f(1, 1, g(1, 1))$.

What is the relation between $[a, b, c]$ and $[A, B, C]$?

d) Given a closed curve $x^4 + y^8 = 2$, there is a point (x_0, y_0) where the gradient is parallel to $[3/5, 4/5]$. Are there two points?

Solution:

a) The gradient vector is zero. At points where the gradient is zero, one in general does not have a tangent line or tangent plane.

b) The tangent plane to the surface $x^2 + y^2 + z^2 = 14$ at $(1, 2, 3)$ is $2x + 4y + 6z = 28$. The graph would be an object in hyperspace!

c) The two vectors are parallel. Actually, if you compute both you see that they are the same in this situation.

d) If \vec{n} is the normal vector, then its direction can be described as $[\cos(t), \sin(t)]$. As we go around the curve, every direction is attained. Yes, there are two points as the direction can change. IN this particular example if (x_0, y_0) is the direction $[3/5, 4/5]$ then $(-x_0, -y_0)$ gives the opposite direction.

3 a) Find an equation of the tangent plane to the parametric surface

$$\vec{r}(u, v) = [u^2, v^2, uv]$$

at the point $(u, v) = (1, 1)$.

b) The surface satisfies the equation $xy - z^2 = 0$. Find the tangent plane to this surface at the same point $(x, y, z) = (1, 1, 1)$ by computing the gradient.

Solution:

a) $\vec{r}(u, v) = [u^2, v^2, uv] \Rightarrow \vec{r}(1, 1) = (1, 1, 1)$. $\vec{r}_u = [2u, 0, v]$; $\vec{r}_v = [0, 2v, u]$, so that a normal vector to the surface at the point $(1, 1, 1)$ is $\vec{r}_u(1, 1) \times \vec{r}_v(1, 1) = [2, 0, 1] \times [0, 2, 1] = [-2, -2, 4]$. The tangent plane at the point $(1, 1, 1)$ is $\boxed{-2x - 2y + 4z = d}$ where the constant d is obtained by plugging in the point $(1, 1, 1)$ and is 0. The equation is $\boxed{-2x - 2y + 4z = 0}$.

b) We have $\nabla f(x, y, z) = [y, x, -2z]$. At the point $(1, 1, 1)$ we get $[1, 1, -2]$ which is parallel to the vector obtained in a). The equation is again of the form $x + y - 2z = d$, where $d = 0$. Again $\boxed{x + y - 2z = 0}$.

- 4 Find an equation of the tangent plane and the normal line to the surface $x - z - 4 \arctan(yz) = 0$ through the point $(1 + \pi, 1, 1)$.

Solution:

Let $f(x, y, z) = x - z - 4 \arctan(yz)$. Then $x - z = 4 \arctan(yz)$ is the level surface $f(x, y, z) = 0$, and $\nabla f(x, y, z) = [1, -\frac{4z}{1 + y^2z^2}, -1 - \frac{4y}{1 + y^2z^2}]$.

a) Tangent plane: $\nabla f(1 + \pi, 1, 1) = [1, -2, -3]$ and an equation of the tangent plane $x - 2y - 3z = d$, where d is the constant obtained by plugging in the point. This is $\boxed{x - 2y - 3z = \pi - 4}$.

b) **Normal line:** The normal line has direction $[1, -2, -3]$, so that the parametrisation is $\boxed{\vec{r}(t) = [1 + \pi + t, 1 - 2t, 1 - 3t]}$.

- 5 a) Show that the ellipsoid $6x^2 + 4y^2 + 2z^2 = 18$ and the sphere $x^2 + y^2 + z^2 - 8x - 6y - 8z + 24 = 0$ are tangent to each other at the point $(1, 1, 2)$, meaning that they have the same tangent

plane at that point.

b) Find a surface different from a plane for which $x + y + 2z = 4$ is the tangent plane at the point $(1, 1, 1)$.

Solution:

a) First note that the point $(1, 1, 2)$ is on both surfaces. The ellipsoid is a level surface of $f(x, y, z) = 3x^2 + 2y^2 + z^2$ and $\nabla f(x, y, z) = [6x, 4y, 2z]$. A normal vector to the surface at $(1, 1, 2)$ is $\nabla f(1, 1, 2) = [6, 4, 4]$ and an equation of the tangent plane there is $6(x-1) + 4(y-1) + 4(z-2) = 0$ or $6x + 4y + 4z = 18$ or $3x + 2y + 2z = 9$.

The sphere is a level surface of $g(x, y, z) = x^2 + y^2 + z^2 - 8x - 6y - 8z + 24 = 0$ and $\nabla g(x, y, z) = [2x - 8, 2y - 6, 2z - 8]$. A normal vector to the sphere at $(1, 1, 2)$ is $\nabla g(1, 1, 2) = [-6, -4, -4]$ and the tangent plane is $6x + 4y + 4z = 18$.

b) An example is $x^2/2 + y^2/2 + z^2 = 2$.

Main definitions

The **gradient** $\nabla f(x, y) = [f_x(x, y), f_y(x, y)]$ or $\nabla f(x, y, z) = [f_x(x, y, z), f_y(x, y, z), f_z(x, y, z)]$ is the higher dimensional version of the first derivative $f'(x)$.

Theorem: Gradients are orthogonal to level sets.

The tangent line through (x_0, y_0) to a level curve $f(x, y) = c$ is $ax + by = d$, where $\nabla f(x_0, y_0) = [a, b]$ and d is obtained by plugging in the point. The tangent plane through (x_0, y_0, z_0) to a level surface $f(x, y, z) = C$ is $ax + by + cz = d$, where $\nabla f(x_0, y_0, z_0) = [a, b, c]$ and d is obtained by plugging in the point.

For parametrized surfaces $\vec{r}(u, v)$, the tangent plane is computed using the vectors \vec{r}_u, \vec{r}_v are velocity vectors of grid curves and so tangent to the surface. The normal is $\vec{n} = \vec{r}_u \times \vec{r}_v = [a, b, c]$ and then get $ax + by + cz = d$, where d is obtained by plugging in the point $\vec{r}(u_0, v_0)$.