

1	
2	
3	
4	
5	
6	
7	
8	
9	
10	
11	
12	
13	
14	
15	

Name:


**LINEAR ALGEBRA AND VECTOR ANALYSIS**

MATH 22A

Total:

Welcome to the final exam. Please don't get started yet. We start all together at 9:00 AM after getting reminded about some formalities. You can fill out the attendance slip already. Also, you can already enter your name into the larger box above.

- You only need this booklet and something to write. Please stow away any other material and any electronic devices. Remember the honor code.
- Please write neatly and give details. Except for problems 2 and 3 we want to see details, even if the answer should be obvious to you.
- Try to answer the question on the same page. There is additional space on the back of each page. If you must, use additional scratch paper at the end.
- If you finish a problem somewhere else, please indicate on the problem page where we can find it.
- You have 180 minutes for this 3-hourly.





**Problem 41E.2) (10 points) Each question is one point:**

a) Name the 3-dimensional analogue of the Mandelbrot set.

b) If  $A$  is a  $5 \times 4$  matrix, then  $A^T$  is a  $m \times n$  matrix. What is  $m$  and  $n$ ?

c) Write down the general formula for the arc length of a curve

$$r(t) = [x(t), y(t), z(t)]^T$$

with  $a \leq t \leq b$ .

d) Write down one possible formula for the curvature of a curve

$$r(t) = [x(t), y(t), z(t)]^T .$$

e) We have seen a parametrization of the 3-sphere invoking three angles  $\phi, \theta_1, \theta_2$ . Either write down the parametrization or recall the name of the mathematician after whom it this parametrization is named.

f) The general change of variable formula for  $\Phi : R \rightarrow G$  is  $\iiint_R f(u, v, w) \boxed{\phantom{du dv dw}} du dv dw = \iiint_G f(x, y, z) dx dy dz$ . Fill in the blank part of the formula.

g) What is the numerical value of  $\log(-i)$ ?

h) We have used the Fubini theorem to prove that  $C^2$  functions  $f(x, y)$  satisfy a partial differential equation. Please write down this important partial differential equation as well as its name. (It was used much later in the course.)

i) What is the integration factor  $|dr|$  for the parametrization

$$r(u, v) = [a \cos(u) \sin(v), b \sin(u) \sin(v), c \cos(v)]^T ?$$

j) In the first lecture, we have defined  $\sqrt{\text{tr}(A^T A)}$  as the length of a matrix. What is the length of the  $3 \times 3$  matrix which contains 1 everywhere?

**Problem 41E.3) (10 points) Each problem is 1 point:**

a) Assume that for a Morse function  $f(x, y)$  the discriminant  $D$  at a critical point  $(x_0, y_0)$  is positive and that  $f_{yy}(x_0, y_0) < 0$ . What can you say about  $f_{xx}(x_0, y_0)$ ?

b) We have proven the identity  $|dr| = |r_u \times r_v|$ , where  $r$  was a map from  $\mathbb{R}^m$  to  $\mathbb{R}^n$ . For which  $m$  and  $n$  was this identity defined?

c) Which of the following is the correct integration factor when using spherical coordinates in 4 dimensions?

$$|d\Phi| = r$$

$$|d\Phi| = (3 + \cos(\phi))$$

$$|d\Phi| = \rho^2 \sin(\phi)$$

$$|d\Phi| = \rho^3 \sin(2\phi)/2$$

d) Which of the following vector fields are gradient fields? (It could be none, one, two, three or all.)

$$F = [x, 0]^T$$

$$F = [0, x]^T$$

$$F = [x, y]^T$$

$$F = [y, x]^T$$

e) Which of the following four surfaces is a one-sheeted hyperboloid? (It could be none, one, two, three or all.)

$$x^2 + y^2 = z^2 - 1$$

$$x^2 - y^2 = 1 - z^2$$

$$x^2 + y^2 = 1 - z^2$$

$$x^2 - y^2 = z^2 + 1$$

f) Parametrize the surface  $x^2 + y^2 - z^2 = 1$  as

$$r(\theta, z) = [\dots\dots\dots, \dots\dots\dots, \dots\dots]^T .$$

g) Who was the creative person who discovered dark matter and proposed the mechanism of gravitational lensing?

h) What is the cosine of the angle between the matrices  $A, B \in M(2, 2)$ , where  $A$  is the identity matrix and  $B$  is the matrix which has 1 everywhere? You should get a concrete number.

i) We have seen the identity  $|v|^2 + |w|^2 = |v - w|^2$ , where  $v, w$  are vectors in  $\mathbb{R}^n$ . What conditions do  $v$  and  $w$  have to satisfy so that the identity holds?

j) Compute the exterior derivative  $dF$  of the differential form

$$F = e^x \sin(y) dx dy + \cos(xyz) dy dz .$$

**Problem 41E.4) (10 points):**

a) (4 points) Find the plane  $\Sigma$  which contains the three points

$$A = (3, 2, 1), \quad B = (3, 3, 2), \quad C = (4, 3, 1) .$$

b) (3 points) What is the area of the triangle  $ABC$ ?

c) (3 points) Find the distance of the origin  $O = (0, 0, 0)$  to the plane  $\Sigma$ .

**Problem 41E.5) (10 points):**

a) (8 points) Find all the critical points of the function

$$f(x, y) = x^5 - 5x + y^3 - 3y$$

and classify these points using the second derivative test.

b) (2 points) Is any of these points a global maximum or global minimum of  $f$ ?

**Problem 41E.6) (10 points):**

a) (8 points) Use the Lagrange method to find **all the maxima and all the minima** of

$$f(x, y) = x^2 + y^2$$

under the constraint

$$g(x, y) = x^4 + y^4 = 16 .$$

b) (2 points) In our formulation of Lagrange theorem, we also mentioned the case, where  $\nabla g(x, y) = [0, 0]^T$ . Why does this case not lead to a critical point here?

**Problem 41E.7) (10 points):**

a) (5 points) The hyper surface

$$S = \{f(x, y, z, w) = x^2 + y^2 + z^2 - w = 5\}$$

defines a three-dimensional manifold in  $\mathbb{R}^4$ . It is poetically called a **hyper-paraboloid**. Find the tangent plane to  $S$  at the point  $(1, 2, 1, 1)$ .

b) (5 points) What is the linear approximation  $L(x, y, z, w)$  of  $f(x, y, z, w)$  at this point  $(1, 2, 1, 1)$ ?

**Problem 41E.8) (10 points):**

Estimate the value  $f(0.1, -0.02)$  for

$$f(x, y) = 3 + x^2 + y + \cos(x + y) + \sin(xy)$$

using quadratic approximation.

**Problem 41E.9) (10 points):**

a) (8 points) We vacation in the **5-star hotel** called **MOTEL 22** in 5-dimensional space and play there ping-pong. The ball is accelerated by gravity  $r''(t) = [x(t), y(t), z(t), v(t), w(t)] = [0, 0, 0, 0, -10]^T$ . We hit the ball at  $r(0) = [4, 3, 2, 1, 2]^T$  and give it an initial velocity  $r'(0) = [5, 6, 0, 0, 3]^T$ . Find the trajectory  $r(t)$ .

b) (2 points) At which positive time  $t > 0$  does the ping-pong ball hit the **hyper ping-pong table**  $w = 0$ ? (The points in this space are labeled  $[x, y, z, v, w]$ .)

**Problem 41E.10) (10 points):**

a) (5 points) Integrate the function  $f(x, y) = (x^2 + y^2)^{22}$  over the region  $G = \{1 < x^2 + y^2 < 4, y > 0\}$ .

b) (5 points) Find the area of the region enclosed by the curve

$$r(t) = [\cos(t), \sin(t) + \cos(2t)]^T,$$

with  $0 \leq t \leq 2\pi$ .

**Problem 41E.11) (10 points):**

a) (7 points) Integrate

$$f(x, y, z) = x^2 + y^2 + z^2$$

over the solid

$$G = \{x^2 + y^2 + z^2 \leq 4, z^2 < 1\}.$$

b) (3 points) What is the volume of the same solid  $G$ ?

**Problem 41E.12) (10 points):**

a) (8 points) Compute the line integral of the vector field

$$F = [yzw + x^6, xzw + y^9, xyw - z^3, xyz + w^4]^T$$

along the path

$$r(t) = [t + \sin(t), \cos(2t), \sin(4t), \cos(7t)]^T$$

from  $t = 0$  to  $t = 2\pi$ .

b) (2 points) What is  $\int_0^{2\pi} r'(t) dt$ ?

**Problem 41E.13) (10 points):**

a) (8 points) Find the line integral of the vector field

$$F(x, y) = [3x - y, 7y + \sin(y^4)]^T$$

along the polygon  $ABCDE$  with  $A = (0, 0)$ ,  $B = (2, 0)$ ,  $C = (2, 4)$ ,  $D = (2, 6)$ ,  $E = (0, 4)$ . The path is closed. It starts at  $A$ , then reaches  $B, C, D, E$  until returning to  $A$  again.

b) (2 points) What is line integral if the curve is traced in the opposite direction?

**Problem 41E.14) (10 points):**

a) (8 points) What is the flux of the vector field

$$F(x, y, z) = [y + x^3, z + y^3, x + z^3]^T$$

through the sphere  $S = \{x^2 + y^2 + z^2 = 9\}$  oriented outwards?

b) (2 points) What is the flux of the same vector field  $F$  through the same sphere  $S$  but where  $S$  is oriented inwards?

**Problem 41E.15) (10 points):**

a) (7 points) What is the flux of the curl of the vector field

$$F(x, y, z) = [-y, x + z(x^2 + y^5), z]^T$$

through the surface

$$S = \{x^2 + y^2 + z^2 + z(x^4 + y^4 + 2\sin(x - y^2z)) = 1, z > 0\}$$

oriented upwards?

b) (3 points) The surface in a) was not closed, it did not include the bottom part

$$D = \{z = 0, x^2 + y^2 \leq 1\}.$$

Assume now that we close the bottom and orient the bottom disc  $D$  downwards. What is the flux of the curl of the same vector field  $F$  through this closed surface obtained by taking the union of  $S$  and  $D$ ?