





Lecture 10: Analysis

Analysis is the science of measure and optimization. As a collection of mathematical fields, it contains **real and complex analysis**, **functional analysis**, **harmonic analysis** and **calculus of variations**. Analysis has relations to calculus, geometry, topology, probability theory and dynamics. We will focus mostly on "the geometry of fractals" today. Examples are Julia sets which belong to the subfield of "complex analysis" of "dynamical systems". "Calculus of variations" is illustrated by the Kakeya needle set in "geometric measure theory", a glimpse of "Fourier analysis" is seen by looking at functions which have fractal graphs, "spectral theory" as part of functional analysis is represented by the "Hofstadter butterfly". As we take a tabloid approach and describe the topic with gossip about some "pop icons" in each field, consider this page the center fold page of the "Analytical Enquirer".

A **fractal** is a set with non-integer dimension. An example is the **Cantor set**, as discovered in 1875 by Henry Smith. Start with the unit interval. Cut the middle third, then cut the middle third from both parts then the middle parts of the four parts etc. The limiting set is the Cantor set. The mathematical theory of fractals belongs to **measure theory** and can also be thought of a playground for real analysis or topology. The term **fractal** had been introduced by Benoit Mandelbrot in 1975. Dimension can be defined in different ways. The simplest is the **box counting definition** which works for most household fractals: if we need n squares of length r to cover a set, then $d = -\log(n)/\log(r)$ converges to the dimension of the set with $r \rightarrow 0$. A curve of length L for example needs L/r squares of length r so that its dimension is 1. A region of area A needs A/r^2 squares of length r to be covered and its dimension is 2. The Cantor set needs to be covered with $n = 2^m$ squares of length $r = 1/3^m$. Its dimension is $-\log(n)/\log(r) = -m \log(2)/(m \log(1/3)) = \log(2)/\log(3)$.




Examples of fractals (for the first, the dimension is not yet known):

Weierstrass function	1872
Koch snowflake	1904
Sierpinski carpet	1915
Menger sponge	1926



Complex analysis extends calculus to the complex. It deals with functions $f(z)$ defined in the complex plane. Integration is done along paths. Complex analysis completes the understanding about functions. It also provides more examples of fractals by iterating functions like the **quadratic map** $f(z) = z^2 + c$:


Newton method	1879
Julia sets	1918
Mandelbrot set	1978
Mandelbar set	1989



Particularly famous are the **Douady rabbit** and the **dragon**, the **dendrite**, the **airplane**.


Calculus of variations is calculus in infinite dimensions. Taking derivatives is called taking "variations". Historically, it started with the problem to find the curve of fastest fall leading to the **Brachistochrone** $r(t) = (t - \sin(t), 1 - \cos(t))$. In calculus, we find maxima and minima of functions. In calculus of variations, we extremize on much larger spaces. Here are some examples of problems:

Brachistochrone	1696
Minimal surface	1760
Geodesics	1830
Isoperimetric problem	1838
Keakeya Needle problem	1917



Fourier theory decomposes a function into basic components of various frequencies $f(x) = a_1 \sin(x) + a_2 \sin(2x) + a_3 \sin(3x) \dots$. The numbers a_i are called Fourier coefficients. Our ear does such a decomposition, when we listen to music. By distinguish different frequencies, our ear produces a Fourier analysis.


Fourier series	1729
Fourier transform (FT)	1811
Discrete FT	Gauss?
Wavelet transform	1930



The Weierstrass function mentioned above is given as the Fourier series $\sum_n a^n \cos(\pi b^n x)$ with $0 < a < 1, ab > 1 + 3\pi/2$. The dimension of its graph is believed to be $2 + \log(a)/\log(b)$.

Spectral theory analyzes linear maps L . The **spectrum** are the real numbers E such that $L - E$ is not invertible. A Hollywood celebrity among all linear maps is the **Mathieu operator** $L(x)_n = x_{n+1} + x_{n-1} + (2 - 2 \cos(cn))x_n$: if we draw the spectrum for for each c , we see the **Hofstadter butterfly**. For fixed c the map describes the behavior of an electron in an almost periodic crystal. An other famous system is the **quantum harmonic oscillator**, $L(f) = f''(x) + f(x)$, the **vibrating drum** $L(f) = f_{xx} + f_{yy}$, where f is the amplitude of the drum and $f = 0$ on the boundary of the drum.


Hydrogen atom	1914
Hofstadter butterfly	1976
Harmonic oscillator	1900
Vibrating drum	1680



All these examples in analysis look unrelated at first. Fractal geometry ties many of them together: spectra are often fractals, minimal configurations have fractal nature, like in solid state physics or in **diffusion limited aggregation** or in other critical phenomena like **percolation** phenomena, **cracks** in solids or the formation of **lighting bolts** In Hamiltonian mechanics, minimal energy configurations are often fractals like **Mather theory**. And solutions to minimizing problems lead to fractals in a natural way like when you have the task to turn around a needle on a table by 180 degrees and minimize the area swept out by the needle. The minimal turn leads to a Kakaya set, which is a fractal. Finally, lets mention some unsolved problems in analysis: does the **Riemann zeta function** $f(z) = \sum_{n=1}^{\infty} 1/n^z$ have all nontrivial roots on the axis $Re(z) = 1/2$? This question is called the **Riemann hypothesis** and is the most important open problem in mathematics. It is an example of a question in **analytic number theory** which also illustrates how analysis has entered into number theory. Some mathematicians think that spectral theory might solve it. Also the Mandelbrot set M is not understood yet: the "holy grail" in the field of complex dynamics is the problem whether it M is locally connected. From the Hofstadter butterfly one knows that it has measure zero. What is its dimension? An other open question in spectral theory is the "can one hear the sound of a drum" problem which asks whether there are two convex drums which are not congruent but which have the same spectrum. In the area of calculus of variations, just one problem: how long is the shortest curve in space such that its convex hull (the union of all possible connections between two points on the curve) contains the unit ball.

Lecture 10: Analysis

The Cantor Set




Analysis is a large branch in mathematics. Its a bit hard to define as it reaches a lot of different areas. It also deals sometimes with rather strange objects like geometries of fractional dimension. Our goal is to understand such **fractals** as they appear in many parts of analysis: spectral theory, complex analysis, harmonic analysis, calculus of variations or functional analysis. Because these fields need some time to learn and explain, the analysis fractals looks like a nice entry point. Our story will become pictorial but there is one single formula, we want to understand:

$$\dim(X) = \frac{-\log(n)}{\log(r)}.$$


It tells that if we want to find the dimension of an object, we cover it with boxes of size $r > 0$ and count how many boxes we need. Assume this number is n . Dimension is what happens if r goes to zero. The prototype of a fractal is the **Cantor set** which was discovered in 1875 by **Henry Smith**. Start with the unit interval. Cut the middle third, then cut the middle third from both parts then the middle parts of the four parts etc. What is left in the end is the Cantor set for which the dimension is $\log(2) / \log(3)$.

The Koch Snowflake




The **Koch snowflake** is an example of a fractal, where the dimension is between 1 and 2. It was first described by the Swedish mathematician Helge von Koch (1870-1924) who described it in 1904. It is a simple model for a **snowflake**. There is a simplified version which just is defined over an interval. It is called the **Koch curve**.

The Tree of Pythagoras




The **tree of Pythagoras** is an example of a fractal with dimension value between 1 and 2. It was first described by the Swedish mathematician **Helge von Koch** (1870-1924). The Koch curve was described by him in 1904. It comes close to actual **trees**. It inspired antenna designs for small devices.

The Sierpinski Carpet



The **Sierpinski carpet** is a fractal in the plane. Its dimension is $\log(8)/\log(2)$. It was described by **Waclav Sierpinski** in 1916.

The Menger Sponge



The **Menger sponge** is a fractal in space. Its dimension is between 2 and 3. It was first described by Karl Menger (1902-1985). Its dimension is $\log(20)/\log(3)$ which is about 2.7.

The Mandelbrot set


We introduce complex numbers $z = a + ib$ and define complex multiplication

$$(a + ib)(u + iv) = au - bv + (av + bu)i$$

Now look at the map $T(z) = z^2 + c$ where c is a fixed complex number. Start with $z = i$ for example, we get $T(z) = i + c$ and $T^2(z) = T(T(z)) = (i + c)^2 + c$ etc. The **Mandelbrot set** is the set of complex numbers $c = a + ib$ for which $T^n(0)$ stays bounded. The **filled in Julia set** J_c of c is the set of z such that $T^n(z)$ stays bounded. The **Julia set** is the boundary of that set.

For example, for $c = 0$, the map is $T_0(z) = z^2$. Since $|z^n| = |z|^n$ we see that the disc $\{|z| \leq 1\}$ is the filled in Julia set for $c = 0$ and the unit circle $\{|z| = 1\}$ is the Julia set.

The following picture (Peitgen-Richter-Saupe) shows the Mandelbrot set in the c plane and a few Julia sets. The circle is shown at the bottom.



A three dimensional version of the Mandelbrot set is called the **Mandelbulb**. It uses spherical coordinates which have been introduced by Euler.