

# TEACHING MATHEMATICS WITH A HISTORICAL PERSPECTIVE

OLIVER KNILL

E-320: Teaching Math with a Historical Perspective

O. Knill, 2010-2022

## Lecture 7: Set Theory and Logic

**7.1.** Sets are fundamental building blocks of mathematics. While **logic** provides a language and the rules for doing mathematics, set theory is the building material for all mathematical structures. Set theory is not the only possible framework. More recently one has used **category theory** as a foundation. It has been seen that **Cantorian set theory** is also quite accessible for younger minds. We have the set of plants, the set of animals, the set of food for example. We can then ask about the relation between these. Are the plants which are animals are there animals which are food etc. What belongs to the set of plants and food which are not both? During the “new math” revolution the language has been introduced to younger kids. Primary school students (including me) had to learn the concept of **Venn diagrams**, long before being exposed to fractions for example. One can argue also that category theory can also be introduced early as it can be seen as some sort of graph theory. But category theory really becomes useful only if one is acquainted with a few mathematical structures. Set theory gives immediate results. By introducing the clever notion of “cardinality”, Cantor could prove that there are different scales of “infinities”. The diagonal argument can be understood long before one can grasp other frame works like calculus and has a “Wow!” effect: we can have different infinities!




FIGURE 1. The intersection  $*$  is the multiplication in the Boolean ring. The symmetric difference  $+$  is the addition.

**7.2.** One can compute with subsets of a given set  $X$  = “universe” like with numbers. There are two basic operations: the **addition**  $A + B$  of two sets is defined as the set of all points which are in exactly one of the sets. The **multiplication**  $A \cdot B$  of two sets contains all the points which are in both sets. With this **symmetric difference**  $\Delta = +$  as addition and the **intersection**  $\cdot = \cap$  as multiplication, the subsets of a given set  $X$  become a **ring**. It is called a **Boolean ring**. It has the property  $A + A = 0$  and  $A \cdot A = A$  for all sets. The **zero element** is the **empty set**  $\emptyset = \{\}$ . The additive inverse (the negative) of  $A$  is the complement  $-A$  of  $A$  in  $X$ . The multiplicative 1-element is the full universe  $X$  under consideration because  $X \cdot A = A$ . As in the ring of integers, the addition and multiplication on sets is **commutative** and multiplication does not have an inverse in general. In class and homework we will play with this ring.

**7.3.** Two sets  $A, B$  have the **same cardinality**, if there exists a **one-to-one map** from  $A$  to  $B$ . For finite sets, this means that they have the same number of elements. Sets which do not have finitely many elements are called **infinite**. Do all sets with infinitely many elements have the same cardinality? The integers  $\mathbb{Z}$  and the natural numbers  $\mathbb{N}$  for example are infinite sets which have the same cardinality:  $f(2n) = n, f(2n + 1) = -n$  establishes a bijection between  $\mathbb{N}$  and  $\mathbb{Z}$ . Also the rational numbers  $Q$  have the same cardinality than  $\mathbb{N}$ . To see this, associate a fraction  $p/q$  with an integer lattice point  $(p, q)$  in the plane, cut out the column  $q = 0$  and run the Ulam spiral on the modified plane. This gives us an explicit enumeration of the rationals. We say that sets which can be counted are called of cardinality  $\aleph_0$ .

**7.4.** Does an interval  $[0, 1]$  have the same cardinality than the real number line  $\mathbb{R}$ ? Even so an interval like  $(-\pi/2, \pi/2)$  has finite length, one can bijectively map it to the real lines with the tan map. Similarly one can see that any two intervals of positive length have the same cardinality. It was a great moment of mathematics, when **Georg Cantor** realized in 1874 that the interval  $(0, 1)$  does not have the same cardinality than the natural numbers  $\mathbb{N}$ . His argument is ingenious: assume, we could count the points  $a_1, a_2, \dots$  in  $[0, 1]$ . If  $0.a_{i1}a_{i2}a_{i3}\dots$  is the decimal expansion of  $a_i$ , define the real number  $b = 0.b_1b_2b_3\dots$ , where  $b_i = a_{ii} + 1 \bmod 10$ . Because this number  $b$  does not agree at the first decimal place with  $a_1$ , nor at the second place with  $a_2$  and so on, the number  $b$  does not appear in that enumeration of all reals. It has positive distance at least  $10^{-i}$  from the  $i$ 'th number (and any representation of the number by a decimal expansion which is equivalent). This is a contradiction. The new cardinality, the **continuum** is also denoted  $\aleph_1$ . The reals are **uncountable**. This gives elegant proofs like the existence of **transcendental number**, numbers which are not algebraic, the root of any polynomial with integer coefficients: algebraic numbers can be counted.

**7.5.** Similarly as one could establish a bijection between the natural numbers  $N$  and the integers  $Z$ , there is a bijection  $f$  between the interval  $I$  and the unit square: if  $x = 0.x_1x_2x_3\dots$  is the decimal expansion of  $x$  then  $f(x) = (0.x_1x_3x_5\dots, 0.x_2x_4x_6\dots)$  is a bijection between  $I$  and  $I \times I$ . Are there cardinalities above  $\aleph_1$ ? Cantor answered also this question. He showed that for an infinite set, the set  $2^X$  of all subsets of the set  $X$  has a larger cardinality than the set  $X$  itself. How does one see this? Assume there is a bijection  $x \rightarrow A(x)$  which maps each point to a set  $A(x)$ . Now look at the set  $B = \{x \mid x \notin A(x)\}$  and let  $b$  be the point in  $X$  which corresponds to  $B$ . If  $y \in B$ , then  $y \notin B(y)$ . On the other hand, if  $y \notin B$ , then  $y \in B$ . The set  $B$  does appear in the "enumeration"  $x \rightarrow A(x)$  of all sets. The set of all subsets of  $\mathbb{N}$  has the same cardinality than the continuum:  $A \rightarrow \sum_{j \in A} 1/2^j$  provides a map from  $P(\mathbb{N})$  to  $[0, 1]$ . The set of all **finite subsets** of  $\mathbb{N}$  however can be counted. The set of all subsets of the real numbers has cardinality  $\aleph_2$ , etc.

**7.6.** Is there a cardinality between  $\aleph_0$  and  $\aleph_1$ ? In other words, is there a set which can not be counted and which is strictly smaller than the continuum in the sense that one can not find a bijection between it and  $\mathbb{R}$ ? This was the first of the 23 problems posed by Hilbert in 1900. The answer is surprising: one has a choice. One can accept either the "yes" or the "no" as a new axiom. In both cases, mathematics is still fine. The nonexistence of a cardinality between  $\aleph_0$  and  $\aleph_1$  is called the **continuum hypothesis** and is usually abbreviated CH. It is independent of the other axioms making up mathematics. This was the work of **Kurt Gödel** in 1940 and **Paul Cohen** in 1963. For most mathematical questions, it does not matter whether one accepts CH or not. The story of exploring the consistency and completeness of axiom systems of all of mathematics is exciting. Euclid gave axioms for Euclidean geometry, Hilbert's goal was much more ambitious, to find a set of axiom systems for all of mathematics. The challenge to prove Euclid's 5'th postulate is paralleled by the quest to prove the CH. But the later is much more fundamental and striking because it deals with **all of mathematics** and not only with a particular field of geometry. Here are the **Zermelo-Frenkel Axioms** (ZFC) including the Axiom of choice

(C) as established by **Ernst Zermelo** in 1908 and **Adolf Fraenkel** and **Thoralf Skolem** in 1922.

### 7.7.

<b>Extension</b>	If two sets have the same elements, they are the same.
<b>Image</b>	Given a function and a set, then the image of the function is a set too.
<b>Pairing</b>	For any two sets, there exists a set which contains both sets.
<b>Property</b>	For any property, there exists a set for which each element has the property.
<b>Union</b>	Given a set of sets, there exists a set which is the union of these sets.
<b>Power</b>	Given a set, there exists the set of all subsets of this set.
<b>Infinity</b>	There exists an infinite set.
<b>Regularity</b>	Every nonempty set has an element which has no intersection with the set.
<b>Choice</b>	Any set of nonempty sets leads to a set which contains an element from each.

**7.8.** The **axiom of choice (C)** has a nonconstructive nature which can lead to seemingly paradoxical results like the **Banach Tarski paradox**: one can cut the unit ball into 5 pieces, rotate and translate the pieces to assemble two identical balls of the same size than the original ball. Gödel and Cohen showed that the axiom of choice is logically independent of the other axioms ZF. Other axioms in ZF have been shown to be independent, like the **axiom of infinity**. A **finitist** would refute this axiom and work without it. It is surprising what one can do with finite sets. The **axiom of regularity** excludes Russellian sets like the set  $X$  of all sets which do not contain themselves. The **Russell paradox** is: does  $X$  contain  $X$ ? It has also been popularized as the **Barber riddle**: a barber in a town only shaves the people who do not shave themselves. Does the barber shave himself?

**7.9.** A complete axiom system for mathematics is not possible because of **Gödel's theorems** of 1931. They deal with **mathematical theories**. They are assumed to be sufficiently strong meaning that one can do at least basic arithmetic in them and call it simply **a theory**:

**First incompleteness theorem:**

In any theory there are true statements which can not be proved within the theory.

**Second incompleteness theorem:**

In any theory, the consistency of the theory can not be proven within the theory.

The proof uses an encoding of mathematical sentences which allows to state liar paradoxical statement "this sentence can not be proved". While the later is an odd recreational entertainment gag, it is the core for a theorem which makes striking statements about mathematics. These theorems are not limitations of mathematics; they illustrate how vast it is. Wouldn't it be terrible if we could build axiom system and enumerate mechanically all possible truths from it?

**7.10.** The work of **George Cantor** and **Kurt Gödel** changed the way we think and teach about mathematics. In both cases, the mathematics community needed time to absorb the implications of the revolutions. Hilbert said about Cantor "Nobody will drive us from the paradise that Cantor has created for us". Cantor clarified the term "cardinality" is, showed that certain infinities like that the cardinality of points in the plane or points in space are the same and most importantly showed that different infinities exist. Gödel's theorems show that mathematics and knowledge in general can not be exhausted by listing a sequence of basic truths from which everything follows. Whenever we make such a list, there are statements which are independent of the system. It would be a mistake to take this as a limitation of mathematics, in contrary it shows that mathematics is inexhaustible: there is always something more to explore.

**7.11.** We first demonstrate that one can compute with sets like with numbers. There is an addition, the symmetric difference and a multiplication, the intersection. With these two operations, we prove the familiar rules of arithmetic

$$A + B = B + A, A \cdot B = B \cdot A, A \cdot (B + C) = A \cdot B + A \cdot C$$

hold. This is called a **Boolean algebra**. There is a set which plays the role of 0. Which one is it? There is also a set which plays the role of 1. Which one is it? One can calculate with sets as with numbers. They form a "Boolean ring".

**Addition:**  $A + B = A \Delta B$  with the zero element  $\emptyset$

**Multiplication:**  $A \cdot B = A \cap B$  with the one element  $\Omega$ .

All the rules of the real numbers apply but there are additional consequences which appear a bit strange  $A + A = 0$  and  $A^2 = A \cdot A = A$ . This means that  $A$  is its own additive inverse. We will visualize the laws of this algebra with **Venn diagrams**.

**7.12. Hilbert's hotel** is located on route 8. It has countably many rooms numbered  $1, 2, 3, \dots$ . The hotel is fully booked. As a newcomer arrives. David, the hotel manager is mortified. David has an idea and moves guest in room  $i$  to room  $i + 1$  and gives the newcomer the first room 1. An other day, the hotel is empty but a large group arrives. They are the "fractions" on their way to a cardinal match with the "squares". Can David accommodate them? He thinks hard and finally manages.

In the summer, the "reals" appear. David is not there but has George, the apprentice is in the office. The group consists of all real numbers between 0 and 1. Can George accommodate them? As much as he tries to shift and renumber, he can not do it.

**7.13.** Paradoxa like the **Liars paradox** "I never tell the truth", the **barbers paradox** "the barber is the person who shaves everybody who does not shave himself" the **surprise exam problem** "it is impossible to make a surprise exam problem", the **heap problem** "take a grain away from a heap keeps it a heap", the **biographer's problem** "who needs one year to write one day of his biography", Here is an other, the **Berry paradox** which comes somehow close to the Goedel numbering: **The smallest integer not definable in less than 11 words.** The problem is that this number is defined in 10 words. This looks like a stupid example but it illustrates that there are properties of numbers like "the shortest way to describe the number" which is not computable.

**7.14.** The **axiom of choice (C)** has a nonconstructive nature which can lead to seemingly paradoxical results like the **Banach Tarski paradox**: one can cut the unit ball into 5 pieces, rotate and translate the pieces to assemble two identical balls of the same size than the original ball. Cohen showed that the axiom of choice is logically independent of the other axioms ZF.

## Work problems

- 7.15.** 1) We have defined addition as  $A + B = A \Delta B$  and **multiplication** as  $A \cdot B = A \cap B$ .
- How can we write the union of two sets  $A \cup B$  using addition and multiplication?
  - Verify that  $A = -A$ . The set  $A$  is the unique set which satisfies  $A + A = \emptyset$ .
  - Draw the Venn diagram of  $A - B = A + (-B)$ .
  - How can we write the set difference  $A \setminus B$  using addition and multiplication. (Note that this is not the difference  $A - B$ ).
  - Draw a diagram which illustrates the associativity property  $A + (B + C) = (A + B) \cdot A + C$  for addition.
  - Draw a diagram which illustrates the associativity property  $A \cdot (B \cdot C) = (A \cdot B) \cdot C$  for multiplication.
  - Draw a diagram which illustrates the distributive property  $A \cdot (B + C) = A \cdot B + A \cdot C$ .
  - Which sets have an inverse  $A^{-1}$  in the sense that  $A \cdot A^{-1} = 1 = X$ ?

**7.16.** 2) The **Russell paradox** defines the set  $X$  of all sets which do not contain themselves as elements.

- a) Verify that if  $X$  contains itself as an element then  $X$  contains itself as an element.
- b) Verify that if  $X$  does not contain itself as an element then  $X$  does not contain itself as an element.
- c) A barber in Cambridge only shaves people who do not shave themselves. Does the barber shave himself? There is an elegant (non mathematical) solution to this paradox. Can you find it?
- d) The first version of the Liar Paradox seems have been found in the fourth century BC by Greek philosopher **Eubulides**, successor of Euclid. Somebody tells "I'm always lying". Does the person tell the truth? The paradox is attributed to the Cretan philosopher **Epimenides** who said "All Cretans are liars."

**7.17.** 3) We explore some cardinality questions. We look at a continuous map from the unit interval onto the unit square. And explore whether it is a bijection.

- a) What is the cardinality of the prime numbers?
- b) What is the cardinality of the set of linear functions  $ax + b$ , where  $a, b$  are real numbers?
- c) A number is called **algebraic** if it is a solution to a polynomial equation

$$p(x) = 0 ,$$

where the polynomial has integer coefficients Can we count all the polynomials? This would establish that one can count all the algebraic numbers and therefore that there are numbers which are not algebraic.