

This is part 1 (of 2) of the homework for the third week. It is due July 15 at the beginning of class.

## SUMMARY.


- $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$ ,  $t \in [a, b]$  **curve** in space.
- $t \mapsto \vec{r}(t)$  is called **parametric representation** of the curve.
- $\vec{r}'(t) = \langle x'(t), y'(t), z'(t) \rangle$  **velocity**,  $\vec{T}(t) = \vec{r}'(t)/|\vec{r}'(t)|$  **unit tangent vector**.
- $\vec{r}''(t) = \langle x''(t), y''(t), z''(t) \rangle$  **acceleration**.
- $\vec{R}(t) = \int_a^t \vec{r}(s) ds = (\int_a^t x(s) ds, \int_a^t y(s) ds, \int_a^t z(s) ds) + (C_1, C_2, C_3)$  **anti-derivative**, satisfies  $\vec{R}'(t) = \vec{r}(t)$ .  $(C_1, C_2, C_3)$  are arbitrary constants.
- $s(t)$  function,  $\vec{r}(t)$  curve,  $\frac{d}{dt}r(s(t)) = r'(s(t))s'(t)$ . **scalar chain rule**.

## Homework Problems

1) (4 points)

Sketch the plane curve  $\vec{r}(t) = \langle x(t), y(t) \rangle = \langle t^3, t^2 \rangle$  for  $t \in [-1, 1]$  by plotting the points for different values of  $t$ . Calculate its velocity  $\vec{r}'(t)$  as well as its acceleration  $\vec{r}''(t)$  at the point  $t = 2$ .

## Solution:



The velocity is  $\vec{r}'(t) = \langle 3t^2, 2t \rangle$ . The acceleration is  $\vec{r}''(t) = \langle 6t, 2 \rangle$ . At the time  $t = 2$  we have  $\vec{r}'(2) = \langle 12, 4 \rangle$  and  $\vec{r}''(2) = \langle 12, 2 \rangle$ .

2) (4 points) A device in a car measures the acceleration  $\vec{r}''(t) = \langle \cos(t), -\cos(3t) \rangle$  at time  $t$ . Assume that the car is at the origin  $(0, 0)$  at time  $t = 0$  and has zero speed at  $t = 0$ , what is its position  $\vec{r}(t)$  at time  $t$ ?

## Solution:

$\vec{r}'(t) = \langle \sin(t), -\sin(3t)/3 \rangle + (C_1, C_2)$ . Because the car has zero speed at time  $t = 0$ , we have  $C_1 = C_2 = 0$ .


From  $\vec{r}'(t) = \langle \sin(t), -\sin(3t)/3 \rangle$ , we obtain  $\vec{r}(t) = \langle -\cos(t), \cos(3t)/9 \rangle + (C_1, C_2)$ . Because  $\vec{r}(0) = (0, 0)$ , we have  $C_1 = 1, C_2 = -1/9$ .  $\vec{r}(t) = \langle -\cos(t) + 1, \cos(3t)/9 - 1/9 \rangle$ .

3) (4 points)

The curve  $\vec{r}(t) = \langle x(t), y(t) \rangle = \langle \int_0^t \sin(s^2) ds, \int_0^t \cos(s^2) ds \rangle$  is called the **cornu spiral** or **clothoid**. You see the curve in the picture.

a) (2) Find the speed of the curve and determine, at which times  $t$  is the velocity vector  $\vec{r}'(t)$  horizontal?

b) (2) What happens with the acceleration  $\vec{r}''(t)$  and the jerk vector  $\vec{r}'''(t)$  when  $t$  goes to  $\infty$ ?



## Solution:

a) The speed is constant 1. The velocity vector is  $\langle \sin(t^2), \cos(t^2) \rangle$ . It is horizontal if  $\cos(t^2) = 0$  which happens for  $t = \sqrt{\pi/2 + \pi k}$ .

b) The acceleration  $\vec{r}''(t) = \langle 2t \cos(t^2), -2t \sin(t^2) \rangle$  and the jerk  $\vec{r}'''(t) = \langle 2 \cos^2(t^2) + 4t^2 \sin(t^2), 2 \sin^2(t^2) + 4t^2 \cos(t^2) \rangle$  oscillate faster and faster too but also grow in amplitude.

4) (4 points)

a) (2) Verify that the curve  $\vec{r}(t) = \langle t \cos(t), 2t \sin(t), t^2 \rangle$  is located on the elliptic paraboloid

$$z = x^2 + y^2/4.$$

b) (2) Use this fact to sketch the curve.

## Solution:

a) Just plug in  $x(t)^2 + y(t)^2 = z$ .

b)



5) (4 points) Find the parameterization  $\vec{r}(t) = \langle x(t), y(t), z(t) \rangle$  of the curve obtained by intersecting the elliptical  $x^2/9 + y^2/4 = 1$  with the surface  $z = xy$ .

a) Write down the formula for the velocity vector  $\vec{r}'(t)$ .

b) If  $s(t) = t^2$ . Find  $\frac{d}{dt}\vec{r}(s(t))$  using the chain rule.

**Solution:**

We find first  $x(t) = 3 \cos(t)$ ,  $y(t) = 2 \sin(t)$  using the first equation. Then get  $z(t) = x(t)y(t) = 6 \cos(t) \sin(t)$ .

$\vec{r}(t) = (x(t), y(t), z(t)) = (3 \cos(t), 2 \sin(t), 6 \cos(t) \sin(t)/2)$ . The velocity vector is  $\vec{r}'(t) = (x'(t), y'(t), z'(t)) = (-3 \sin(t), 2 \cos(t), 6 \cos^2(t) - 6 \sin^2(t))$ .


b)  $d/dtr(s(t)) = r'(s^2)2t = (-3 \sin(t), 2 \cos(t), 6 \cos^2(t) - 6 \sin^2(t))2t$ .

## Challenge Problems

(Solutions to these problems are **not** turned in with the homework.)

- 1) A closed curve in space is called a **knot**. Consider the space curve  $\vec{r}(t) = (\sin(3t), \cos(4t), \cos(5t))$ . Find the smallest interval  $[a, b]$  such that this curve is a knot. Sketch the curve.

**Hint.** You might also use technology: `ParametricPlot3D[{Sin[3t], Cos[4t], Cos[5t]}, {t, 0, 2Pi}]` in Mathematica.



- 2) How could one verify that it is not possible to deform the knot  $\vec{r}(t) = (\sin(3t), \cos(4t), \cos(5t))$  into the trivial knot  $\vec{r}(t) = (\cos(t), \sin(t), 0)$  in such a way that during the deformation, the curve can never self-intersect?

**Hint.** Look at the possible types of closed curves which don't intersect the knot. How many different types are there for the trivial knot or for the given knot?