

|       |
|-------|
| Name: |
|-------|

- Start by writing your name in the above box.
- Try to answer each question on the same page as the question is asked. If needed, use the back or the next empty page for work. If you need additional paper, write your name on it.
- Do not detach pages from this exam packet or unstaple the packet.
- Except for problems 1,2 and 6, give details.
- Please write neatly. Answers which are illegible for the grader can not be given credit.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have exactly 90 minutes to complete your work.

|        |  |     |
|--------|--|-----|
| 1      |  | 20  |
| 2      |  | 10  |
| 3      |  | 10  |
| 4      |  | 10  |
| 5      |  | 10  |
| 6      |  | 10  |
| 7      |  | 10  |
| 8      |  | 10  |
| 9      |  | 10  |
| Total: |  | 100 |


Problem 1) (20 points) No justifications are needed.

- 1) ☐ T ☐ F      The integral  $\int_a^b |\vec{T}'(t)| dt$  measures the arc length of a curve.
- 2) ☐ T ☐ F      The length of the vector projection  $P_{\vec{v}}(\vec{w})$  is the absolute value of the scalar projection  $\text{comp}_{\vec{v}}(\vec{w})$ .
- 3) ☐ T ☐ F      The velocity vector of  $\vec{r}(t) = [t^5, t^3, t^7]^T$  at time  $t = 1$  is  $[1, 1, 1]^T$ .
- 4) ☐ T ☐ F      If  $\vec{v} \times \vec{w} = \vec{v} \times \vec{w}$  then  $\vec{v}, \vec{w}$  are parallel.
- 5) ☐ T ☐ F      The vector  $[1, 2, 3]^T$  is perpendicular to the line  $[t, 2t, 3t]^T$ .
- 6) ☐ T ☐ F      The point given in spherical coordinates as  $(\rho, \phi, \theta) = (1, \pi, \pi)$  is the same point than the point  $(\rho, \phi, \theta) = (1, \pi, 0)$ .
- 7) ☐ T ☐ F      The parametrized curve  $\vec{r}(t) = [3 \cos(t), 0, 5 \sin(t)]^T$  is an ellipse.
- 8) ☐ T ☐ F      The curvature of a line is 1 at every point of the line.
- 9) ☐ T ☐ F      If  $|\vec{v} \times \vec{w}| = \vec{v} \cdot \vec{w}$  then either  $\vec{v}$  is parallel to  $\vec{w}$  or perpendicular to  $\vec{w}$ .
- 10) ☐ T ☐ F      If the dot product between two unit vectors is 1, then the two vectors are the same.
- 11) ☐ T ☐ F      We have  $|(\vec{u} \times \vec{v}) \times \vec{w}| \leq |\vec{u}||\vec{v}||\vec{w}|$  for all vectors  $\vec{v}, \vec{w}, \vec{u}$ .
- 12) ☐ T ☐ F      The curvature of a curve  $\vec{r}(t)$  is given by  $\kappa(t) = |\vec{T}'(t)|/|\vec{r}''(t)|$ .
- 13) ☐ T ☐ F      The arc length of the curve  $[\sin(t^2), \cos(t^2)]^T$  from  $t = 0$  to  $t = 1$  is equal to 1.
- 14) ☐ T ☐ F      Given three planes in space, there is a point which is on two planes at least.
- 15) ☐ T ☐ F      The curve  $\vec{r}(t) = [t^2, t, t^4]^T$  is the intersection curve of the two cylindrical paraboloids  $x = y^2$  and  $z = x^2$ .
- 16) ☐ T ☐ F      The lines  $\vec{r}_1(t) = [5 + t, 3 - t, 2 + t]^T$  and  $\vec{r}_2(t) = [5 - t, 3 + t, 2t]^T$  intersect perpendicularly.
- 17) ☐ T ☐ F      You can intersect a two sheeted hyperboloid with a plane to get a parabola.
- 18) ☐ T ☐ F      The vector  $[12/13, 3/13, 4/13]^T$  is a direction.
- 19) ☐ T ☐ F       $\vec{v} \times (\vec{u} \times \vec{v}) = \vec{0}$  for all vectors  $\vec{u}, \vec{v}$ .
- 20) ☐ T ☐ F      If  $f(x, y) = x^2 y^2$ , then  $f_{xx} = f_{yy}$ .


Total

Problem 2) (10 points) No justifications are needed in this problem.


a) (2 points) Match the contour plots. Enter O, if there is no match.



I




II




III

| Function $g(x, y, z) =$ | Enter O,I,II or III |
|-------------------------|---------------------|
| $x + 2y$                |                     |
| $x^2 + 2y^2 + 3z^2$     |                     |
| $x^2 - 2y^2$            |                     |
| $x^2 + 2y^2$            |                     |
| $x^2 + 2y$              |                     |


b) (2 points) Match the graphs of the functions  $f(x, y)$ . Enter O, if there is no match.



I




II



III

| Function $f(x, y) =$   | Enter O,I,II or III |
|------------------------|---------------------|
| $\cos(x^2 - y^2)$      |                     |
| $ xy $                 |                     |
| $x^2 \exp(-x^2 - y^2)$ |                     |
| $x^4$                  |                     |
| $\exp(-x)$             |                     |


c) (2 points) Match the plane curves with their parametrizations  $\vec{r}(t)$ . Enter O, if there is no match.



I




II




III

| Parametrization $\vec{r}(t) =$             | Enter O, I,II or III |
|--------------------------------------------|----------------------|
| $\vec{r}(t) = [t^2, 2t^2]^T$               |                      |
| $\vec{r}(t) = [t^2, t^4 + 1]^T$            |                      |
| $\vec{r}(t) = [\cos(\sqrt{t}), \sin(t)]^T$ |                      |
| $\vec{r}(t) = [t, t^3]^T$                  |                      |
| $\vec{r}(t) = [\exp( t ), t]^T$            |                      |


d) (2 points) Match functions  $g$  with level surface  $g(x, y, z) = 1$ . Enter O, if no match.



I




II




III

| Function $g(x, y, z) = 1$ | Enter O, I,II or III |
|---------------------------|----------------------|
| $x^2 + y^4 = 1$           |                      |
| $x^4 - y^4 + z^4 = 1$     |                      |
| $y^4 = 1$                 |                      |
| $x^2 + y^2 + z^4 = 1$     |                      |
| $xy - z = 1$              |                      |


e) (2 points) Match the parametrization. Enter O, where no match.



I



II



III

| $\vec{r}(u, v) =$             | Enter O,I,II or III |
|-------------------------------|---------------------|
| $[u, v, u^2v]^T$              |                     |
| $[\cos v, uv, v]^T$           |                     |
| $[u \cos(v), u \sin(v), u]^T$ |                     |
| $[u \cos(v), u \sin(v), v]^T$ |                     |
| $[\cos v, u \sin(v), v]^T$    |                     |

Problem 3) (10 points)

Routine "toy" problems:

- a) (2 points)  $([1, 1, 1]^T \cdot [2, 1, 2]^T) + |[4, 3, 2]^T|^2$ .
- b) (2 points)  $|([2, 3, 4]^T \times [2, 3, 5]^T) + [0, 6, 0]^T|$ .
- c) (2 points)  $[4, 2, 3]^T \cdot ([3, 2, 1]^T \times [1, 1, 1]^T)$ .
- d) (2 points)  $\vec{P}_{\vec{w}}(\vec{v})$  with  $\vec{v} = [1, 4, 1]^T$  and  $\vec{w} = [9, 18, 9]^T$ .
- e) (2 points) The angle  $\alpha$  between  $[0, 3, 3]^T, [2, 0, 2]^T$ .



Problem 4) (10 points)

There are four **Kepler-Poinsot solids**. They are non-convex platonic solids. The first is centered at  $A = (1, -1, 1)$ , the second at  $B = (2, 2, 0)$ , the third at  $C = (4, 5, 6)$  the fourth at  $P = (3, 3, 3)$ . Find the distance from  $P$  to the plane through  $A, B, C$ .




Problem 5) (10 points)

a) (7 points) A **cloth hanger** is padded around a curve which has the form

$$\vec{r}(t) = [t^2 \cos(t), t^2 \sin(t), t^3/3]^T$$

with  $-1 \leq t \leq 1$ . Find the arc length from  $t = -1$  to  $t = 1$ .

b) (3 points) As the picture indicates, the curvature appears problematic at  $t = 0$ . Verify, that the curvature does not exist there by writing down a formula for the curvature and indicating that it can not be evaluated at  $t = 0$ .



Problem 6) (10 points)

No justifications are needed in this problem. You can assume that  $\vec{r}(t)$  is a space curve, and that the vectors  $\vec{v}, \vec{w}, \vec{r}', \vec{r}'', \vec{T}, \vec{T}', \vec{N}, \vec{B}$  are non-zero, where  $\vec{N}$  is the normal vector and  $\vec{B}$  the bi-normal vector. All these vectors  $\vec{r}, \vec{T}, \vec{B}, \vec{N}$  and its derivatives are evaluated at a fixed time, like  $t = 0$ .

| First vector                         | Second vector            | always parallel | always perpendicular | depends |
|--------------------------------------|--------------------------|-----------------|----------------------|---------|
| $\vec{v} \times \vec{v}$             | $\vec{v}$                |                 |                      |         |
| $\vec{w} + \vec{v}$                  | $\vec{v} - \vec{w}$      |                 |                      |         |
| $\vec{v} \times \vec{w}$             | $\vec{w} \times \vec{v}$ |                 |                      |         |
| $(\vec{v} + \vec{w}) \times \vec{w}$ | $\vec{v} \times \vec{w}$ |                 |                      |         |
| $\text{Proj}_{\vec{v}}(\vec{w})$     | $\vec{v}$                |                 |                      |         |
| $\text{Proj}_{\vec{v}}(\vec{w})$     | $\vec{w}$                |                 |                      |         |
| $\vec{T}$                            | $\vec{r}'$               |                 |                      |         |
| $\vec{T}$                            | $\vec{T}'$               |                 |                      |         |
| $\vec{T}'$                           | $\vec{r}''$              |                 |                      |         |
| $\vec{B}$                            | $\vec{N}$                |                 |                      |         |

Problem 7) (10 points)

**111 Huntington** is a tall building in Boston which is in the shadow of the Prudential and Hancock tower. But its probably the most interesting building mathematically. It is only 169 meters high (compared with Hancock with 241) but look at the beauty of the dome! Inside this structure there are two metal connection lines

$$\vec{r}_1(t) = [t, 3t, -t]^T$$

and

$$\vec{r}_2(t) = [1 + 2t, t, t]^T .$$

Find their distance.



Problem 8) (10 points)

Very soon, Apple is expected to come out with an **iWatch**. Lets build one ourselves. Simplicity is the rule: it consists only of a band, and a "home button". And because Apple is known for fancy packaging, we strapped around a one sheeted hyperboloid. For each of the following surfaces, find a parametrization of the form

$$\vec{r}(u, v) = [x(u, v), y(u, v), z(u, v)]^T .$$

a) (3 points) "Band" cylinder


$$x^2 + y^2 = 9 .$$

b) (4 points) "Button" ellipsoid

$$10(x - 3)^2 + y^2 + z^2 = 1/4 .$$

c) (3 points) "Package" 1-sheeted hyperboloid

$$x^2 + y^2 - z^2 - 1/2 = 0 .$$



Problem 9) (10 points)

This June 2014, the Swiss extreme sports women **Géraldine Fasnacht** jumped with a wing-suit from the Matterhorn (a mountain in Switzerland). When flying with a wingsuit, there is the gravitational force, the force from the wind and a force from the wing. Assume

$$\vec{r}''(t) = [1, t, \exp(-t) - 10]^T$$

and  $\vec{r}(0) = [0, 0, 4500]^T$  and  $\vec{r}'(0) = [0, 2, 0]^T$ . Find the path  $\vec{r}(t)$ .

