

7/21/2022 SECOND HOURLY Practice 5      Maths 21a, O.Knill, Summer 2022

**"I affirm my awareness of the standards of the Harvard College Honor Code."**

Name:

- Start by writing your name in the above box.
- Try to answer each question on the same page as the question is asked. If needed, use the back or the next empty page for work. If you need additional paper, write your name on it.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader can not be given credit.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have exactly 90 minutes to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
Total:		100


Problem 1) True/False questions (20 points). No justifications needed.

Mark for each of the 20 questions the correct letter. No justifications are needed.

- 1)  T  F The directional derivative of  $f(x, y)$  in the direction  $[1, 0]$  is  $f_x(x, y)$ .
- 2)  T  F The surface area of the unit sphere  $x^2 + y^2 + z^2 = 1$  is  $4\pi$ .
- 3)  T  F The point  $(0, 0)$  is a critical point of  $f(x, y) = x^5y^4$ .
- 4)  T  F The function  $f(x, y) = x^2 - y^2$  has a global minimum under the constraint  $y = 0$ .
- 5)  T  F The gradient  $\nabla f$  of the function  $f(x, y)$  with graph  $z = 3x + y$  is  $[3, 1, -1]$ .
- 6)  T  F If  $(0, 0)$  is a critical point for  $f$  and the second directional derivative  $D_{\vec{v}}D_{\vec{v}}f(0, 0)$  is positive for all unit vectors  $\vec{v}$ , then  $(0, 0)$  is a local minimum.
- 7)  T  F If  $(0, 0)$  is a local maximum for  $f$ , then  $f_{xy}(0, 0) = 0$ .
- 8)  T  F For  $\vec{u} = [1, 1]/\sqrt{2}$ , we have  $D_{\vec{u}}f = f_{xy}$ .
- 9)  T  F The chain rule assures that  $\frac{d}{dt}H(x(t), y(t)) = H_x(x(t), y(t))x'(t) + H_y(x(t), y(t))y'(t)$ .
- 10)  T  F The function  $f = g^2$  under the constraint  $g(x, y) = x^2 + y^2 = 1$  has never a finite set of minima.
- 11)  T  F The function  $u(x, y) = x^2 - y^2$  solves the partial differential equation  $u_x^2 - u_y^2 = 0$ .
- 12)  T  F For every point  $(x, y)$  (not necessarily a critical point), there exists a direction  $\vec{v}$  for which  $D_{\vec{v}}f(0, 0) = 0$ .
- 13)  T  F The identity  $f_{xxx} = f_{yyy}$  holds for all smooth functions  $f(x, y)$ .
- 14)  T  F The integral  $\int_0^1 \int_0^{x^2} 1 \, dydx + \int_0^1 \int_0^{\sqrt{y}} 1 \, dx dy = 1$ .
- 15)  T  F The directional derivative satisfies  $D_{\vec{v}}f = (f_{xx}f_{yy} - f_{xy}^2) = \nabla f(x, y) \cdot \vec{v}$ .
- 16)  T  F Fubini's theorem assures that  $\int_0^1 \int_0^x f(x, y) \, dydx = \int_0^1 \int_0^y f(x, y) \, dx dy$ .
- 17)  T  F When computing the surface area of the Gabriel trumpet given by  $r = 1/z, z \geq 1$ , we got the integral  $\int_0^{2\pi} \int_1^\infty \frac{1}{z} \sqrt{1 + z^{-4}} \, dz d\theta$ .
- 18)  T  F In class, we were able to compute the integral  $\int_{-\infty}^\infty e^{-x^2} \, dx = \pi$ .
- 19)  T  F The gradient vector to  $z = x^2 + y^2$  at  $(1, 1, 4)$  is  $[2x, 2y]$ .
- 20)  T  F The integral  $\iint_{x^2+y^2 \leq 1} |f(x, y)| \, dx dy$  computes the surface area of the surface  $z = f(x, y), x^2 + y^2 \leq 1$ .

Problem 2) (10 points) No justifications are needed

a) (6 points) Match the following regions with their area computation.



Enter A-F	Area Integral
	$\int_0^{2\pi} \int_0^{1+\sin(5\theta)} r \, dr d\theta$
	$\int_0^1 \int_0^{1-x^2} 1 \, dy dx$
	$\int_0^{2\pi} \int_0^{2+\sin(5\theta)} r \, dr d\theta$
	$\int_0^1 \int_{\sqrt{y}}^1 1 \, dx dy$
	$\int_0^{2\pi} \int_0^{ \sin(5\theta) } r \, dr d\theta$
	$\int_0^1 \int_{x^2}^1 1 \, dy dx$

b) (4 points) In the Book "In Pursuit of the Unknown", the English mathematician **Ian Stewart** covers 17 equations. Some of the are partial differential equations. Which of the following 4 equations appears in the sticky list seen in the picture? There is just one.



Fill in a)-d)	Name
	Transport
	Burgers
	Heat
	Wave

Problem 3) (10 points) (No justifications are needed.)

**X-alps challenge** is a cool alpine race which took place 2 weeks ago. The athletes had to cross the alps several times either by foot or paraglider from Innsbruck to Monaco. For the 5th time, the best was the Swiss competitor **Chrigel Maurer**. He covered 2272 kilometers in 11 days. Participants have first to answer a theoretical question:




a) (2 points) Assume  $g(x, y, z) = x + 2y + z$  is the amount of "thermal uplift" at location  $(x, y, z)$  and you are at  $(0, 0, 0)$  you want to go into direction, in which the uplift increase is largest. In which direction  $\vec{v}/|\vec{v}|$  do you go?

- |                          |                             |                           |  |
|--------------------------|-----------------------------|---------------------------|--|
| A) $\vec{v} = [1, 2, 1]$ | B) $\vec{v} = [-1, -2, -1]$ | C) $\vec{v} = [2, -1, 0]$ | Check A)-C): <input style="width: 100%;" type="text"/> |
|--------------------------|-----------------------------|---------------------------|--|

b) (8 points) Now let's look at the terrain. In each part, pick the correct point in A – K. There is a possible match so that each letter appears exactly once.

	Choose one A-H
A point where $f_x = 0$ and $f_y > 0$	<input style="width: 100%;" type="text"/>
A point where $f_y = 0$ and $f_x > 0$ and $f_{yy} = 0$	<input style="width: 100%;" type="text"/>
A point where $ \nabla f $ is maximal and $f_x > 0$	<input style="width: 100%;" type="text"/>
A point where $f_x = 0$ and $f_y < 0$	<input style="width: 100%;" type="text"/>
A local minimum	<input style="width: 100%;" type="text"/>
A global maximum, the Matterhorn	<input style="width: 100%;" type="text"/>
A saddle point	<input style="width: 100%;" type="text"/>
A local minimum under the constraint $x = 0$	<input style="width: 100%;" type="text"/>



Problem 4) (10 points)

**New England houses** often feature “two slope roofs”. If the length of each roof part is 1 and the angle of the upper roof is  $x$  and the angle of the lower  $y$ , then the attic room gained under the roof is

$$f(x, y) = \cos(x) + \cos(y) .$$

Assume the architect has the constraint to build it so that

$$g(x, y) = \sin(x) + \sin(y) = 1 .$$

Find the optimal  $x, y$  using the Lagrange method. We are only interested in solutions where  $x, y$  are acute angles so that there is exactly one solution. Find it.



Problem 5) (10 points)

We want to design an **US mailbox** for which the cost functional

$$f(x, y) = 4xy - (3\pi + 4)y^2 - 4x$$

is extremal. [ The  $4xy$  the surface area of the box part counting positive because it stores mail, the cylinder material and leg material parts counts negative. ]

While it turns out that we can not find a minimum or maximum for  $f$ , we want to use the second derivative test to find and classify all parameters  $(x, y)$  which are critical points of  $f$ .



Problem 6) (10 points)

Oliver got a **lemon tree** this summer. It is seen on the picture to the right. One of the leaves is parametrized by


$$\vec{r}(x, y) = [x, y, y + x^2]$$

over the triangle  $G : 0 \leq y \leq 1, y \leq x \leq 1$ .

a) (5 points) Verify that the surface area simplifies to

$$\int_0^1 \int_y^1 \sqrt{2 + 4x^2} \, dx dy .$$

b) (5 points) Solve this integral! And make sure to use some of the sour power to slice that lemon.



Problem 7) (10 points)

One of the creative problems in the Mathematica project will be to create a **pasta** of your own.

a) (5 points) Find the tangent plane to the perfect **X-macaroni**


$$f(x, y, z) = x^2y^4 + y^2x^4 + 4z^2 = 6$$

at the point  $(1, 1, 1)$ .

b) (5 points) To taste the X-pasta, we cut it at  $z = 0$  to get the curve

$$g(x, y) = x^2y^4 + y^2x^4 = 6 .$$

Find the tangent line at  $(1, \sqrt{2})$  to this curve.



Problem 8) (10 points)

While writing this exam, Oliver drank from an **orange soda can** of radius 1 and height 6. If the can is empty or full, the center of mass is in the middle of the can. After having tasted the juice and knowing the juice to be 4 times heavier than the can, the center of mass has gone down: there must exist a minimal value by Rolle's theorem. We simplify the problem by assuming the can is rectangular (which just changes constants). Lets find the center function  $f(h)$  which depends on the **juice level height**  $0 \leq h \leq 6$ .

$$f(h) = \frac{\int_0^1 \int_0^6 y dy dx + 4 \int_0^1 \int_0^h y dy dx}{\int_0^1 \int_0^6 1 dy dx + 4 \int_0^1 \int_0^h 1 dy dx}$$

- a) (4 points) Compute each of the four double integral in this expression.
- b) (3 points) What is  $f(h)$ ? You don't need to simplify except if you want to make it easier for c).
- c) (3 points) Evaluate the  $f(h)$  values for  $h = 0, h = 3$  and  $h = 6$  to check that  $f(0) = f(6)$  is indeed in the middle of the can and that  $f(3)$  is smaller.



P.S. ignore the product placement for 4 brands on this picture. You also don't have to find the minimum  $h$ . It turns out to be the golden ratio  $(\sqrt{5} - 1)/2$  times the height. Golden juice + Golden ratio = Golden summer!

Problem 9) (10 points)


- a) (5 points) Find the volume below the graph of the function  $f(x, y) = x^4 + y^4$  and above the square  $G$  given by  $-1 \leq x \leq 1, -1 \leq y \leq 1$ . In other words, find

$$\int \int_G f(x, y) dx dy .$$

- b) (5 points) 2017 was the year of the **fidget spinner**! What is the moment of inertia

$$\int \int_G x^2 + y^2 dx dy$$

of the **fidget spinner region**  $G$  given in polar coordinates as  $1 \leq r \leq 3 + \sin(3\theta)$ ?



Problem 10) (10 points)

a) (4 points) Write down the double integral for the surface area of

$$\vec{r}(x, y) = \langle 2x, y, x^3/3 + y \rangle$$

with  $0 \leq x \leq 2$  and  $0 \leq y \leq x^3$ .

b) (6 points) Find the surface area.

