

**"I affirm my awareness of the standards of the Harvard College Honor Code."**

Name:

- Start by writing your name in the above box.
- Try to answer each question on the same page as the question is asked. If needed, use the back or the next empty page for work. If you need additional paper, write your name on it.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader can not be given credit.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have exactly 90 minutes to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
Total:		120




Problem 1) (20 points) No justifications are needed.

- 1)  T  F For any vector  $\vec{v} = [a, b, c]$  we have  $|[a, b, c]| = [|a|, |b|, |c|]$ .
- 2)  T  F The curvature of a curve at a point is independent of the parametrization.
- 3)  T  F It is possible to intersect a cylinder with a plane and get a hyperbola.
- 4)  T  F If  $\vec{T}, \vec{N}, \vec{B}$  is a TNB frame then  $\vec{N} = \vec{B} \times \vec{T}$
- 5)  T  F The intersection between two spheres of radius 1 and 2 is either empty, a point, a circle.
- 6)  T  F The set of points in space which satisfy  $x^2 - y^2 = 1$  form a hyperbola.
- 7)  T  F The length of the sum of two vectors in space is always larger or equal than the sum of the lengths of the vectors.
- 8)  T  F For any three vectors  $\vec{u}, \vec{v}, \vec{w}$ , the identity  $\vec{u} \times (\vec{v} + \vec{w}) = (\vec{u} \times \vec{v}) + \vec{u} \times \vec{w}$  holds.
- 9)  T  F The set of points which satisfy  $-x^2 - 2x + y^2 + z^2 = 1$  define a cone.
- 10)  T  F If  $A, B, C$  are three points space which are not contained in a common line, then  $\vec{AB} \times \vec{AC}$  is a vector orthogonal to the plane containing  $A, B, C$ .
- 11)  T  F The line  $\vec{r}(t) = [t, 5t, 4t]$  hits the plane  $x + 5y + 4z = 100$  at a right angle.
- 12)  T  F The surface given in  $(r, \theta, z)$  coordinates as  $r = \sin(\theta)$  is a paraboloid.
- 13)  T  F If  $\vec{v} \times \vec{w} = \vec{w} \times \vec{v}$ , then  $\vec{v}$  and  $\vec{w}$  are parallel (in the sense that there exists a constant  $c$  such that  $\vec{v} = c\vec{w}$ ).
- 14)  T  F If  $|\vec{x} \times \vec{v}| = 0$  for all vectors  $\vec{v}$ , then  $\vec{x} = \vec{0}$ .
- 15)  T  F If  $\vec{u}$  and  $\vec{v}$  are orthogonal, then  $(\vec{u} \times \vec{v}) \times \vec{u}$  is parallel to  $\vec{v}$ .
- 16)  T  F Every vector contained in the line  $\vec{r}(t) = [4 + 2t, 2 + 3t, 3 + 4t]$  is parallel to the vector  $(4, 2, 3)$ .
- 17)  T  F If in spherical coordinates a point is given by  $(\rho, \theta, \phi) = (1/2, 3\pi/2, \pi/2)$ , then its rectangular coordinates are  $(x, y, z) = (0, -1/2, 0)$ .
- 18)  T  F The set of points which satisfy  $x^2 - 4x + 2y^2 + 3z^2 = -3$  is an ellipsoid.
- 19)  T  F If  $\vec{u} \cdot (\vec{v} \times \vec{w}) = 0$ , then all three vectors  $\vec{u}, \vec{v}, \vec{w}$  are in the same plane.
- 20)  T  F The set of points in  $\mathbb{R}^3$  which have distance 1 from the curve  $\vec{r}(t) = [3 \cos(t), 3 \sin(t), 0]$  form a torus (doughnut)



Problem 2) (10 points) No justifications are needed in this problem.

In each sub-problem, each of the numbers 0,1,2,3 each occur exactly once.

a) (2 points) Match the graphs of the functions  $f(x, y)$ . Enter 0 if there is no match.

1		2		3	
Function $f(x, y) =$					0,1,2, or 3
$xy(x^2 - y^2)$					
$e^{-x^2-y^2}(x^2 + y^2)$					
$1/(x^2 + y^4 + 1)$					
$ \sin(x + y) $					

b) (2 points) Match the surfaces  $g(x, y, z) = c$ . Enter 0 if there is no match.

1		2		3	
Function $g(x, y, z) =$					0,1,2, or 3
$1000 * \sin(xyz) = 1$					
$x^6 + z^6 = 1$					
$z - (x^2 - y^2)x = 0$					
$x - y^2 = 1$					


c) (2 points) Match the space curves with the parametrizations. Enter 0 if there is no match.

1		2		3	
Parametrization $\vec{r}(t) =$					0,1,2, or 3
$[t \cos(t), t \sin(t), \sin(t)]$					
$[\cos(3t), 0, 3 \sin(3t)]$					
$[\exp(t), 2 \exp(t), 3 \exp(t)]$					
$[t, 0, t \sin(t)]$					

d) (2 points) Match the functions  $g$  with contour plots in the  $xy$ -plane. Enter 0 if there is no match.

1		2		3	
Function $g(x, y) =$					0,1,2, or 3
$x^2 - y^2$					
$(x + y)^4$					
$x^3 - y$					
$\cos(3x) + \sin(3y)$					

e) (2 points) Match the parametrized surfaces. Enter 0 if there is no match.

1		2		3	
Parametrization $\vec{r}(u, v) =$					0-3
$[\cos(v), \sin(v), u]$					
$[u^2 \cos(v), u^2 \sin(v), u]$					
$[\sin(v) \cos(u), \cos(v), 2 \sin(v) \sin(u)]$					
$[u^6 + v^6, u^3, v^3]$					

Problem 3) (10 points)

We perform some computations with the vectors  $\vec{v} = [2, 2, 3]$  and  $\vec{w} = [1, 2, 2]$ .


- a) (2 points) Find the cross product  $\vec{v} \times \vec{w}$ ?
- b) (2 points) Construct a unit vector in the same direction than  $\vec{v} \times \vec{w}$ .
- c) (2 points) Find  $\cos(\alpha)$  for the angle  $\alpha$  between  $\vec{v}, \vec{w}$ .
- d) (2 points) What is the vector projection  $\vec{P}_{\vec{w}}(\vec{v})$  of  $\vec{v}$  onto  $\vec{w}$ ?
- e) (2 points) Compute  $(\vec{v} + \vec{w}) \cdot (\vec{v} \times \vec{w})$ .

Problem 4) (10 points)

- a) (2 points) Parametrize the plane containing the points  $(1, 0, 0), (0, 5, 0), (0, 0, 2)$  using parameters  $s, t$ :
- b) (3 points) Now find the equation  $ax + by + cz = d$  of that plane in a).
- c) (5 points) Finally find the distance between that plane defined in a) and the point  $P = (3, 3, 3)$ .

Problem 5) (10 points)

A **prismatic polyhedron** with 8 vertices contains the 4 points  $A = (2, 0, 0), B = (1, \sqrt{3}, 0)$  and  $C = (-1, \sqrt{3}, 0), D = (0, 0, \sqrt{3})$ . Find the distance of the line containing the points  $A, B$  and the line containing the points  $C, D$ .



Problem 6) (10 points)

a) (5 points) Find the arc length of the path

$$\vec{r}(t) = \left[ \frac{t^3}{3}, t^2, 2t \right]$$

with  $-1 \leq t \leq 1$ .

b) (5 points) Compute  $\vec{v} = \vec{r}'(1)$ ,  $\vec{w} = \vec{r}''(1)$  and express the curvature of the curve at  $\vec{r}(1)$  in terms of  $\vec{v}$  and  $\vec{w}$ .

Problem 7) (10 points)

On Mars, the sum of gravitational acceleration and wind force is  $\vec{r}''(t) = [0, \sin(t), -4]$ . The **Mars helicopter Ingenuity** has been rising up to 5 meters. There had been a small stone pebble stuck on one of the legs. It falls down from the position  $\vec{r}(0) = [2, 1, 5]$  with velocity  $\vec{r}'(0) = [1, 0, 0]$  subject to the acceleration given above.

a) (6 points) Determine the path of the pebble.

b) (4 points) At which time does the pebble hit the ground?

Problem 8) (10 points)

We denote with  $|ABC|$  the **area** of a triangle  $ABC$  defined by three points  $A, B, C$ . Take  $A = (2, 0, 0)$ ,  $B = (0, 3, 0)$ ,  $C = (0, 0, 1)$  as well the point  $O = (0, 0, 0)$ . Verify in this case the **3D Pythagoras theorem**

$$|ABC|^2 = |ABO|^2 + |BCO|^2 + |ACO|^2 .$$

$$|ABC|^2 =$$

$$|ABO|^2 =$$

$$|BCO|^2 =$$

$$|ACO|^2 =$$

Problem 9) (10 points) No justifications are needed.

a) (2 points) Parametrize the surface  $3x + 2y + 4z = 12$ .

$$\vec{r}(s, t) = \left[ \boxed{\phantom{000}}, \boxed{\phantom{000}}, \boxed{\phantom{000}} \right]$$

b) (2 points) Parametrize the surface  $y^2 + (z - 1)^2 = 9$  using an angle  $\theta$  in the  $yz$ -plane.

$$\vec{r}(\theta, x) = \left[ \boxed{\phantom{000}}, \boxed{\phantom{000}}, \boxed{\phantom{000}} \right]$$

c) (2 points) Parametrize the surface  $x^2 + 2x + y^2 + z^2/9 = 0$ .

$$\vec{r}(\theta, \phi) = \left[ \boxed{\phantom{000}}, \boxed{\phantom{000}}, \boxed{\phantom{000}} \right]$$

d) (2 points) Parametrize the surface  $y = x^4 - z^4$ .

$$\vec{r}(x, z) = \left[ \boxed{\phantom{000}}, \boxed{\phantom{000}}, \boxed{\phantom{000}} \right]$$

e) (2 points) Parametrize the surface obtained by taking the helix  $r(\vec{t}) = [\cos(t), \sin(t), t]$  and connect each point  $r(\vec{t})$  with the projection onto the  $z$ -axes.

$$\vec{r}(t, s) = \left[ \boxed{\phantom{000}}, \boxed{\phantom{000}}, \boxed{\phantom{000}} \right]$$

