

"I affirm my awareness of the standards of the Harvard College Honor Code."

Name:

- Start by writing your name in the above box.
- Try to answer each question on the same page as the question is asked. If needed, use the back or the next empty page for work. If you need additional paper, write your name on it.
- Do not detach pages from this exam packet or unstaple the packet.
- Please write neatly. Answers which are illegible for the grader can not be given credit.
- No notes, books, calculators, computers, or other electronic aids can be allowed.
- You have exactly 90 minutes to complete your work.

1		20
2		10
3		10
4		10
5		10
6		10
7		10
8		10
9		10
10		10
11		10
Total:		120

Problem 1) (20 points) No justifications are needed.


- 1) T F The plane $x = 3$ does intersect the yz -plane.
- 2) T F The curve $\vec{r}(t) = [1 + 2t, t, 1 + t]$ intersects the z -axis in a point.
- 3) T F The Cauchy-Schwartz inequality states $|\vec{v} \cdot \vec{w}| \leq |\vec{v}|$ for any two vectors \vec{v}, \vec{w} .
- 4) T F The curvature of a circle $\vec{r}(t) = [\cos(2t), 0, \sin(2t)]$ is equal to $1/2$ everywhere.
- 5) T F The surface $y^2 - x + z^2 = 2$ is an elliptic paraboloid.
- 6) T F The angle between the vectors $0\vec{A}$ and $0\vec{B}$ is positive if the distance between the points A and B is positive.
- 7) T F Let $\vec{j} = [0, 1, 0]$. There is a vector \vec{v} for which the vector projection of \vec{v} onto \vec{j} is equal to $-\vec{j}$.
- 8) T F Two particles with path $\vec{r}_1(t) = [0, t, -t]$ and $\vec{r}_2(t) = [1 - t, t - 1, 0]$ do collide.
- 9) T F In spherical coordinates the surface $\rho^2 \sin^2(\phi) - \rho^2 \cos^2(\phi) = 1$ is a one-sheeted hyperboloid.
- 10) T F If $|\vec{u} \times \vec{v}| = 1$, for unit vectors \vec{u}, \vec{v} , then \vec{u}, \vec{v} are orthogonal.
- 11) T F The curve $r^2 \cos^2(\theta) - r^2 \sin^2(\theta) = 1$ in polar coordinates is a hyperbola.
- 12) T F If the arc length of a curve connecting A with B is 0, then $A = B$.
- 13) T F The surface parametrized as $\vec{r}(y, z) = [y, z, y^2 - z^2]$ is a hyperbolic paraboloid.
- 14) T F The velocity vector and the acceleration are always either parallel or perpendicular.
- 15) T F It is possible that a plane and a one-sheeted hyperboloid intersects in two crossing lines.
- 16) T F The function $f(x, y) = \log(x^2 + y^2)$ contains as domain all points except the origin $(0, 0)$.
- 17) T F The normal vector \vec{N} and the unit tangent vector \vec{T} are perpendicular if \vec{T}, \vec{T}' are both not zero vectors.
- 18) T F The distance between two non-parallel lines in three dimensional space can be zero.
- 19) T F If $\vec{i}, \vec{j}, \vec{k}$ denote the unit vectors in the x, y and z axis, then $\vec{i} \cdot (\vec{j} \times \vec{k}) = 1$.
- 20) T F For any two lines L, M , there are points P on L and Q on M such that $d(P, Q) = 2d(L, M)$, where $d(L, M)$ is the distance between the lines.

Problem 2) (10 points) No justifications are needed in this problem.


In each sub-problem, each of the numbers 0,1,2,3 each occur exactly once.

a) (2 points) Match the surfaces $g(x, y, z) = c$. Enter 0 if there is no match.


1



2



3




Function $g(x, y, z) =$	0,1,2, or 3
$x - y^2 + z = 0$	
$x^2 + y^2 - z^4 = 1$	
$z - e^{-x^2-y^2} = 0$	
$y^2 + z^3 = 1$	

b) (2 points) Match the graphs of the functions $f(x, y)$. Enter 0 if there is no match.


1



2



3




Function $f(x, y) =$	0,1,2, or 3
$\sqrt{ 1 + x^2 - y^2 }$	
$\sin(x^2 + y^2)$	
$ x - y - x + y $	
$y^2 \sin(x^4)$	

c) (2 points) Match the space curves with the parametrizations. Enter 0 if there is no match.


1



2




3



Parametrization $\vec{r}(t) =$	0,1,2, or 3
$[t, t, t]$	
$[\cos(2t), 0, \cos(2t)]$	
$[0, \cos(2t), \sin(2t)]$	
$[t, \sin(t), 0]$	

d) (2 points) Match the functions g with contour plots in the xy -plane. Enter 0 if there is no match.

1



2




3




Function $g(x, y) =$	0,1,2, or 3
$\cos(2x) + \sin(2y)$	
$(x - y)^2$	
$y^2 - x$	
$(2x^2 + 7y^2)^2$	

e) (2 points) Match the parametrized surfaces. Enter 0 if there is no match.

1



2



3



Parametrization $\vec{r}(u, v) =$	0-3
$[u, u^2 + v^2, v]$	
$[\sin(v) \cos(u), \sin(v) \sin(u), \cos(v)]$	
$[u^2 - v^2, u, v]$	
$[u \cos(v), u \sin(v), u]$	

$$\vec{r}(s, t) =$$

b) (5 points) Find the distance between $4x + 4y + 2z = 0$ and the point $(2, 2, 2)$.

$$\text{distance} =$$

Problem 5) (10 points)

a) (5 points) Find a parametrization $\vec{r}(t)$ of the intersection of the planes

$$x + y + z = 1, \quad 2x - y + 2z = 2 .$$

$$\vec{r}(t) =$$

b) (5 points) Find the distance between that line computed in a) and $P = (1, 1, 1)$.

$$\text{distance} =$$

Problem 6) (10 points)

a) (5 points) Find the arc length of the path

$$\vec{r}(t) = \left[\frac{3t^2}{2}, \frac{4t^2}{2}, \frac{5t^3}{3} \right]$$

with $0 \leq t \leq 1$.

Length =

b) (5 points) Find the curvature of $\vec{r}(t)$ at $t = 1$ using the vectors $\vec{v} = \vec{r}'(1)$, $\vec{w} = \vec{r}''(1)$.

$\kappa(\vec{r}(1)) =$

Problem 7) (10 points)

a) (5 points) Given the **jerk**

$$\vec{r}'''(t) = [0, 0, -12]$$

with $\vec{r}(0) = [0, 0, 3]$, $\vec{r}'(0) = [4, 0, 0]$, $\vec{r}''(0) = [0, 2, 0]$, find $\vec{r}(t)$ and especially $\vec{r}(10)$.

$$\vec{r}(10) =$$

b) (5 points) Compute the unit tangent vector \vec{T} of the TNB-frame to the curve $\vec{r}(t)$ at $t = 0$.

$$\vec{T}(0) =$$

Problem 8) (10 points)

We experiment with a **paper air plane**. The wing tips are $C = (0, 3, 4)$ and $D = (0, -3, 4)$ the front is $A = (3, 0, 0)$ the back is $B = (-3, 0, 0)$.

a) (5 points) The sum of the areas of the triangles ABC and ABD is the total wing area. Find the wing area.

